

X-disease phytoplasma and Little Cherry Virus 2 Proficiency test Protocols

Proficiency testing is conducted by WSU with cooperating labs in order to help labs maintain the highest quality outputs for growers. WSU is not a certifying agency and can only provide testing as a service to the cherry industry.

Steps:

- 1. WSU Prosser will obtain samples from known negative and known positive trees (12 samples).
- 2. Samples are pooled, homogenized and a subsample tested by WSU Prosser.
- 3. Subsamples of pooled samples are provided to participating labs which contain tissue to be tested with accompanying excel submission sheet.
- 4. For consistency both labs will process the same tissue. Tissue type will depend on timing of analysis, harvest timing = fruit stems, after harvest pooled leaf/bark samples.
- 5. Extraction and analysis conducted using standardized protocols (Appendix 1-3).
- 6. Sample results from participating labs are provided to WSU Prosser Dr Scott Harper in excel spreadsheet format with column 1 sample name, column 2 LChV2 positive/negative (designate as P/N), column 3 LChV2 Ct value, column 4 XDP positive/negative (designate as P/N), column 5 XDP Ct value.
- 7. Labs will also provide accompanying run data to allow for Dr. Harper to check that appropriate protocols were used.
- 8. To pass the panel, all negative samples must be correctly identified, and 85% of results of positives must fall within a range of 3 cycles (Ct values).

Appendix 1 Extraction Protocols



PCNW-VDL Request No.
 Performed by:
Date:

Nucleic Acid Extraction Type: Total RNA

Method Approved for use on: Cherry (Prunus avium & P. dulcis)

Peach & Nectarine (P. persica)

Plum (*P. domestica*) Apple (*Malus* sp.) Pear (*Pyrus* sp.)

Reagents Needed: Qiagen RNeasy Plant Mini Kit (Qiagen Cat. 74904)

Polyvinylpyrrolidone-40 (Sigma Cat. PVP40) Sodium Metabisulfite (Sigma Cat. 31448) Antifoam A Concentrate (Sigma Cat. A6582)

20% N-Laurylsarcosine Sodium Salt Solution (Sigma Cat. L5125)

Before Beginning the Extraction:

- Prepare modified RLC buffer (RLC with 2.5% w/v PVP-40, 0.1% v/v sodium metabisulfite, and 0.2% v/v Antifoam A).
- 2. Prepare RPE buffer by adding four volumes of 96-100% ethanol to the stock solution.

Assay Protocol:

- 1. Weigh 400 mg of scraped bark or stem, petiole, and/or leaf midrib tissue into a grinding bag.
- 2. Homogenize tissue in 2 mL of modified RLC buffer, then transfer 700 μ l of the homogenate into a 1.5 ml microcentrifuge tube.
- 3. Add 84 μ l of N-laurylsarcosine solution and incubate at 65 °C for 10 minutes.
- 4. Centrifuge at 16,100 x g for 4 minutes, then transfer the supernatant to a QIAshredder (purple) column and proceed with steps 5-10 <u>OR</u> transfer the supernatant to a 2 ml tube and follow the Qiacube process.
- 5. Centrifuge at 16,100 x g for 2 minutes, then transfer the supernatant to a new tube and add 0.5 volumes of 96-100% ethanol. Mix and transfer a maximum of 700 μ l to an RNeasy spin column (pink).
- 6. Centrifuge at 8,000 x g for 1 minute, and discard the flow-through. Add 700 μ l of RW1 buffer and centrifuge at 8,000 x g for 1 minute. Again, discard the flow-through.
- 7. Add 500 μ l of RPE buffer to the column and centrifuge at 8,000 x g for 1 minute, then discard the flow-through.
- 8. Add another 500 μ l of RPE buffer to the column and centrifuge at 8,000 x g for 1 minute, then discard the flow-through.
- 9. Place the column in a new 2 ml collection tube and centrifuge dry for 1 minute at 16,100 x g.
- 10. Place the column in a clean 1.5 or 2 ml microcentrifuge tube and add 100 μ l of RNAse-free H₂O. Incubate at room temperature for 2 minutes, then centrifuge at 8,000 x g for 1 minute. Discard the column and store the samples.

Appendix 2 Little Cherry Virus 2



 CPCNW-VDL Request No.
Performed by:
Date:

Pathogen: Little cherry virus 2 (Ampelovirus)

Assay Type: Real-Time PCR

Assay Source: Harper et al. (unpublished)

Primer Name	Orientation	Sequence (5'3')	Binding Site	Fluorophore
LChV2-RT-F	+	GCGGAGGGTCTGGAGACTT	6769-6787	
LChV2-RT-R	-	AGGCCAGGAGACAACAACATC	6832-6850	FAM
LChV2-RT-P	+	TGAGTCTTTCAAGGACAC	6789-6806	

Controls to be used:

1. Positive Little cherry virus-2 positive tissue

2. Negative Virus-free Prunus tissue

3. No-Template Water

Reagents to be used: Invitrogen SuperScript III Platinum One-Step qRT-PCR Kit (11732-020)

PCR Hood Number:	qPCR Machine Number:	
2x Master Mix Lot Number:	Date Opened	
Enzyme Lot Number:	Date Opened	_
Number of reactions needed:	Add 1 additional reaction to	account for wastaa

Reagent	Concentration	1x	Master Mix
2x Reaction Mix	N/A	10	
Forward Primer (10 μ M)	400 nM	0.8	
Reverse Primer (10 μM)	400 nM	0.8	
Probe (10 μM)	100 nM	0.2	
SSIII/Taq enzyme mix	N/A	0.4	
RNAse-free H₂O	N/A	5.8	
RNA template	N/A	2	
Total Volume:		20	

Thermocycling conditions
50 °C for 15 minutes
94 °C for 2 minutes
40x 94 °C for 10 seconds
60 °C for 45 seconds

Please attach the real-time results spreadsheet to the back of this worksheet, add any comments in the space below, and attach the completed worksheet to the back of the diagnostic request form.

Appendix 3 X-disease phytoplasma



	VDL Request No	/-VDL Request No	PO
ned by:	Performed by:	Performed by:	
Date:	Date:	Date:	

Pathogen: Phytoplasma **Assay Type:** Real-Time PCR

Assay Source: Ito and Suzuki (2017) PLoS ONE 12(9): e0185427

Primer Name	Orientation	Sequence (5'3')	Binding Site	Fluorophore
UPH-F (F)	+	CGTACGAAGTATGAAACTTAAAGGA	280281-280299	
D-UPHr2 (R)	-	CGACAACCATGCACCACCTGIIIIICTGATAACC	280442-280423	FAM
UPH-Pb (P)	+	TGACGGGACTCCGCACAAGCG	280302-280322	

Controls to be used:

1. Positive Phytoplasma positive tissue

2. Negative Phytoplasma-free Vitis or Prunus tissue

3. No-Template Water

Reagents to be used: Applied Biosystems TaqMan Universal PCR Master Mix (4304437)

PCR Hood Number:	qPCRMachine Number:	
2x Master Mix Lot Number:	Date Opened	

Number of reactions needed: ______ Add 1 additional reaction to account for wastage

Reagent	Concentration	1x	Master Mix
2x Reaction Mix	N/A	10	
Forward Primer (10 μ M)	300 nM	0.6	
Reverse Primer (10 μM)	300 nM	0.6	
Probe (10 μM)	100 nM	0.2	
DNA template	N/A	2	
RNAse-free H₂O	N/A	6.6	
	Total Volume:	20	

Thermocycling conditions
50 °C for 2 minutes
95 °C for 10 minutes
40x 95 °C for 15 seconds
60 °C for 60 seconds

Please attach the real-time results spreadsheet to the back of this worksheet, add any comments in the space below, and attach the completed worksheet to the back of the diagnostic request form.