AGRICULTURAL EXTENSION LABORATORY

REPORT OF PLANT TISSUE ANALYSIS FERTILITY ASSAY

County: Submitted by: MERCED SCHEUERMAN

R. MEYER, L. BROWN, R. BURAU AND R. VOSS

No.

P-9429-D

Date submitted: 9-15-76

Date reported: 7-15-77

Identification:

JOE ALVERNAZ

CROP: SWEET POTATO

Sample No.	Alfred and	Description			211	NO ₃ -N	PO ₄ -P	SO ₄ -S	N	P	К	Ca	Mg	Na	CI	В	Zn	BR DRY BASIS
l	AE	and		TRA		ppm	ppm	ppm	%	%	%	%	%	%	%	ppm	ppm	PPM 375
1. 2 3 4	3B 4A 4B				auktine duing		State	Strong St	07 > 09-07	The gardinate	2 C O O O O O O O O O O O O O O O O O O	asports of the organical and the second seco	segots altwo	n. Your cooperation	320 318 372			
	dried and acid					8		vor v vor v		5,000	3,000	000.8	, K. 0.50	9,000		below which	in comple	
	a bround a					280	000			1000	1000	2000		1800		the crop	bro gail	
	omple of the disconnection of					300	8							3		519	returning	
	ordxe br					1.0		0	-	0	64 82	0 -		S4 6		utay be re	1 10	
	se dos sa de							54						1		educed.	de " se	
	te bro a					10.		10		73 73 * *	100 mg		3.	5 5 5			liw atset	

SO₄-S <u>Sulfate-Sulfur</u> — The inorganic sulfur fraction in the dried and ground sample, expressed as ppm sulfur in the dry plant material.

All other elements determined as total elemental content and expressed as percent or parts per million of dry plant material.

Critical Concentrations**

CROP	PLANT PART	SAMPLING TIME	NO ₃ -N	PO ₄ -P	SO ₄ -S ppm	K %	B	Zn
Alfalfa	midstem	1/10 bloom		700	250*	0.75	25*	
Sugar Beets	petiole of recently	anytime during	1000	750	250*	1.0		10*
Ladino Clover	matured leaf	growing season	***	600		0.8		
Sub Clover	7 2 3 18			600	170			
Grape s	petiole opposite flower cluster	bloom time	T.S. var. only 226			0.8	25	15

Approximate Adequate Nutrient Levels***

R ₁ A		July 1	10,000	1000	6	15*
0 (6)	petiole of recently matured leaf	July 15	5,000	1000		15*
Cotton (San Joa-		August 1	2,000	1000		15*
quin Valley)		August 15	1,000	1000	2.5	15*
		days from planting				
Potato	petiole of 4th ma- ture leaf from top	< 70	8,000	1200	9	
Foluto		70-90	6,000	800	7	
		over 90	3,000	500	4	
Townsto process of	whole plant top	thinning or transplant	1,500	1500	1.5	10*
Tomato process-	petiole of recently	set	6,000	1800	2.5	10*
ing crop		1st color	2,000			10*
	matured leaf	1st pick	300	700	0.75	10*

Level in blades of recently matured leaves.

These values are based on present information. Your cooperation in completing and returning "followup" sheets will add greatly to the development of more and better interpretation data on a wider range of crops.

^{**} The nutrient concentration below which plant growth begins to decrease.

^{***} The approximate nutrient level at various growth stages below which the crop yield may be reduced.

C hatternamentally

White the state of the s

COOPERATIVE EXTENSION UNIVERSITY OF CALIFORNIA

Report No: <u>P-9429-D</u> Received <u>SEP 151976</u>

LABORATORY WORK REQUEST

	and in the control of									
	County MERCEd	Submitted by Robert W. Scheuerman								
	Date sampled 9-13-76	Send report copies to R.D. MEYER, A.L. Brown R. VOSS, & R.G. BURAN								
	Identification	R, VOSS, & R.G. BURAM								
	Identification	Living STON, Ca								
		T R(Required for Water)								
	Crop SWEET POTATOE Soil type	Drainage								
	Irrigation method	Water source								
	No. of samples Soil	, water plant, feed, misc. (circle one only)								
	Plant part sampled: Petiole, blade, mid-rib, grain, whole tops, mid-stem: LEAUES									
	Analyses requested Bloom									
	Project No. or problem description									
	Sample No. Description (for plants, include	e age, stage of growth, or cutting)								
P. Tolk										
13	A									
2.3	B									
3. A	1.4									
	10									
4.4										
		0 1 01 1011								
-		Keter to 19-1061								
		Gr Root Samples								
		destroyed by wice								
		7-12-77								
British and Applications described		J-6								
Spinish requirement of the later										
0										