# **Health Care Resources in Northeast Indiana**

# A Hope Center

3630 Hobson Rd. 260-422-3544



3702 S. Clinton St. 260-969-4357

Columbia City 260-244-7779

### A Mother's Hope

260-376-0097 www.amothershopewfw.org

## **Allen County STD Clinic**

4813 New Haven Ave. 260-449-7504

## **Compassion Pregnancy Center**

Angola– 2107 N. Wayne St. 260-668-7773

LaGrange– 221 S. Detroit St. 260-463-7377

#### **Catholic Charities**

915 S. Clinton St. 260-422-5625

## **Helping Hands**

Bluffton-116 E. Dustman Rd. 260-824-4263

### **Life and Family Services**

Kendallville– 201 S. Park Ave. 260-347-5720

#### Matthew 25 Health & Dental

413 E. Jefferson Blvd. 260-426-3250

### **Neighborhood Health Clinic**

1717 S Calhoun St. 260-458-2641

### **The Hope Clinic**

Decatur 870 S. 11th St. 260-728-4191

Berne– 165 W. Water St. 260-589-3561

#### **Women's Care Center**

4600 W. Jefferson Blvd. 260-203-5476

921 W. Coliseum Blvd. 260-483-8918 '

2910 Lake Ave 260-424-9377



We can Help, Know your Options.

## Women who have an abortion

- Are 3x more likely to commit suicide. (1)
- Have an 81% increased risk of mental health problems. (1)
- Have a 70% higher rate of deliberate self harm than women who give birth. (1)
- Have a 110% increased risk of alcohol abuse. (1)

# **Teenagers who have an abortion**

- Are up to 10x more likely to attempt suicide than their counterparts who have not had an abortion. (1)
- Risk of breast cancer increases by 30% among women who chose abortion after giving birth once. (2)
- Risk of breast cancer increases by 50% for women who chose abortion before giving birth. (2)
- 64% of women who have an abortion report feeling coerced or forced into the decision. If you are being coerced call the Center Against Forced Abortions at (210)614-7157. (3)

# **Fetal Development**



- A human's heart starts beating 18 to 21 days after conception. (4)
- Babies have brainwaves 6 weeks after conception. (4)
- By 18 weeks after conception your baby can hear. (5)
- During an abortion, the cervix is forcibly dilated and subjected to injury, and this damage to the cervix may cause a woman to later have a premature delivery. (6)
  - (1) http://healthresearchfunding.org/19-shocking-post-abortion-depression-statistics/
  - (2) J. Brind et al., "Induced abortion as an independent risk factor for breast cancer: a comprehensive review and meta-analysis," Journal of Epidemiology & Community Health 50 (1996): 481-496.
  - (3) http://www.theunchoice.com/articles/howcommoniscoercion.htm
  - (4) http://www.ehd.org/
  - (5) http://www.healthline.com/health/pregnancy/when-can-a-fetus-hear
  - (6) http://www.bcpinstitute.org/FactSheets/FS-induced\_abortion.pdf