

PRODUCT MONOGRAPH

MICRO Se
(Selenious Acid Injection USP)

Trace Element

Sandoz Canada Inc.
145 Jules-Léger
Boucherville PQ

Date of Preparation
June 12, 2006

Control # 106397

PRODUCT MONOGRAPH

MICRO Se

Selenious Acid Injection USP

(40 µg/mL)

Trace Element

CLINICAL PHARMACOLOGY

Selenium is an essential component of glutathione peroxidase, an enzyme which helps prevent oxidative damage to cells caused by peroxides and free radicals.

The absorption, retention and distribution of selenium within the body, and the amounts, forms and routes of excretion vary with the chemical forms and amounts of the element ingested and with the dietary levels of other elements such as arsenic and mercury.

Selenium is absorbed from the small intestine and distributed widely to tissues including liver, skin, muscle, kidney, lung, brain, testis, ovary, heart, spleen, thyroid, pancreas, dental enamel and fingernails.

Blood levels have been shown to vary geographically due to selenium levels of soil and food. Reported values of selenium levels in whole blood vary from 150 ng/ml in selenium deficient areas to 3200 ng/ml in a seleniferous zone (in China). Studies in North America have reported whole blood levels of 70 to 229 ng/ml and plasma levels of 78 to 157 ng/ml.

Excretion occurs mainly via the kidneys. However a small amount of the endogenous stores is lost through the feces.

Keshan disease, a cardiomyopathy of children and young women in China, is the only clinical condition that has been firmly linked to selenium deficiency. Plasma selenium levels have been shown to decrease during total parenteral nutrition (TPN), however symptoms of selenium deficiency are not seen in all TPN patients. Several factors may influence the development of selenium deficiency including geographical location, nutritional and clinical status, excessive GI losses, age, volume of fluid administered and duration of selenium-deficient TPN.

Several cases of cardiomyopathy have occurred in patients receiving total parenteral nutrition. Muscle pain and weakness have been reported during total parenteral nutrition and have responded to selenium supplementation.

Selenium supplementation during TPN helps prevent development of the following deficiency symptoms: cardiomyopathy, muscle pain and weakness.

INDICATIONS

MICRO Se (Selenious Acid Injection USP) is indicated as a supplement to intravenous solutions given for TPN. Its administration in TPN solutions helps to maintain plasma selenium levels, and to prevent depletion of endogenous stores of selenium and subsequent deficiency symptoms.

WARNINGS

MICRO Se is a hypotonic solution which should be administered in admixtures only.

If toxicity symptoms occur due to selenium, discontinue supplementation of TPN solutions immediately.

Do not give undiluted **MICRO Se** by direct injection into a peripheral vein because of the potential of infusion phlebitis.

PRECAUTIONS

The possibility of selenium retention should be considered in patients with renal dysfunction and/or gastrointestinal malfunction since selenium is eliminated in the urine and to a smaller extent in the feces.

ADVERSE REACTIONS

The amount of selenium present in **MICRO Se** is small. Symptoms of toxicity from selenium are unlikely to occur at the recommended dosage level.

OVERDOSAGE

Chronic selenium toxicity due either to occupation-related exposure, high selenium content in food/water or oral supplements, resembles arsenic toxicity. Hair loss, white horizontal streaking on fingernails, paronychia, fatigue, irritability, hyperreflexia, nausea, vomiting, garlic odor on breath, and metallic taste characterize toxicity. Muscle tenderness, tremor, lightheadedness, and facial flushing are observed in selenite poisoning. Serum selenium levels are elevated but do not correlate well with symptoms. Blood chemistries, hematology, and liver and renal function tests are usually normal.

Acute selenious acid ingestions are almost invariably fatal. Stupor, respiratory depression, hypotension, and death can result several hours postingestion. Severe hypotension develops secondary both to decreased contractility from a toxic cardiomyopathy and to inappropriately low peripheral vascular resistance. Laboratory abnormalities include thrombocytopenia, moderate hepatorenal dysfunction, and elevated serum creatine kinase levels.

The electrocardiogram may demonstrate ST elevations and T-wave changes characteristic of myocardial infarction. The urinary excretion of selenium is rapid. Terminal respiratory

failure developed after a selenious acid ingestion (15 mL gun bluing solution) despite the use of an extracorporeal membrane oxygenator. Death occurred on the 18th hospital day. The plasma selenium level reached 285 μ g/mL on the first hospital day and returned to normal levels by day 4.

There are no antidotes to selenious acid toxicity; treatment is expectant (cardiopulmonary monitoring in an intensive care setting) and supportive (intravenous infusion, supplemental oxygen and ventilation as needed).

DOSAGE AND ADMINISTRATION

Dosage:

The suggested dosage ranges are:

Adults

For the metabolically stable adult receiving TPN, the suggested additive dosage level is 20 to 40 μ g selenium per day.

Pediatrics

For pediatric patients the suggested dosage level is 3 μ g selenium per kg per day.

Administration:

Routine monitoring of selenium levels is suggested as a guideline for administration. The range for selenium blood levels has been reported as plasma, 78 to 157 ng/ml; whole blood, 70 to 229 ng/ml.

MICRO Se should be aseptically added to the TPN solution under the laminar flow hood. The selenious acid present in **MICRO Se** is physically compatible with electrolytes and vitamins usually present in the amino acid/dextrose solution used for TPN.

PHARMACEUTICAL INFORMATION

DRUG SUBSTANCE

Proper Name: Selenious Acid

Chemical Name: Selenium di-oxide, monohydrated

Molecular Formula: H_2SeO_3

Molecular Weight: 128.97

Description: Selenious Acid is a colourless or white crystal, efforescent in dry air and hygroscopic in moist air. It is insoluble in water and in alcohol.

COMPOSITION

Each mL of **MICRO Se** contains 65.36 µg of Selenious acid (equivalent to 40 µg Selenium). Nitric acid is used to adjust the pH.

DILUTION FOR INTRAVENOUS USE

Aseptic addition of **MICRO Se** to the amino acid/dextrose component of a TPN solution under a laminar flow hood is recommended. After dilution, the solution must be used within 24 hours. Visually inspect parenteral drug products for particulate matter and discoloration prior to administration whenever container and solution permit.

STABILITY AND STORAGE RECOMMENDATIONS

Store at controlled room temperature not exceeding 28°C. Do not permit to freeze.

AVAILABILITY

One strength, 40 µg selenium/mL, is available in single dose 10 mL clear glass vials in boxes of 10.

MICRO Se is for intravenous use after dilution only.

REFERENCES

1. Underwood EJ. Trace elements in human and animal nutrition. 4th ed. New Academic Press 1977: 302-346.
2. Grieg PD, Baker JP, Jeejeebhoy KN. Metabolic effects of total parenteral nutrition. *Ann Rev Nutr* 1982; 2: 179-199.
3. Combs GF, and Combs SB Ch 8. Selenium in human nutrition and health. In: The role of selenium in nutrition. Academic Press 1986; 327-399.
4. Levander OA, Burk RF. Report on the 1986 A.S.P.E.N. Research workshop on selenium in clinical nutrition. *JEPN* 1986; 10: 545-549.
5. Van Rij AM, Thomson CD, McKenzie JM and Robinson MF. Selenium deficiency in total parenteral nutrition. *Am J Clin Nutr* 1979; 32: 2076-2085.
6. Phillips GD, and Garnys VP. Parenteral administration of trace elements to critically ill patients. *Anaesth Intens Care* 1981; 9: 221-225.
7. Johnson RA, Baker SS, Fallon JT, Maynard EP, Ruskin JN, Wen Z, Ge K, and Cohen HJ. An accidental case of cardiomyopathy and selenium deficiency. *N Eng J Med* 1981; 304: 1210-1212.
8. Ellenhorn MJ, and Barceloux, eds. *Medical Toxicology. Diagnosis and treatment of human poisoning.* Elsevier Science Publishing Co. Inc., New York 1988; 1059-1060.
9. Phillips GD, and Odgers CL. Parenteral Nutrition: Current status and concepts. *Drug* 1982; 23: 276-323.
10. Ricour C, Gros J, Maziere B, Comar D. Trace elements in children on total parenteral nutrition (TPN). *Acta. Chir Scand. Suppl.* 1975; 466: 22-23.