





The Basics

Scientific name: Acer


macrophyllum

Family: Sapindaceae

Height: 50-65 ft (tallest - 88') **Diameter**: 2-5 ft (largest - 97") **Crown**: 50+ ft (largest - 104')

Age: ~200 years
Notable features:

- Produces edible flowers and
 - sap
- Support many epiphytes (mosses, liverworts, ferns)
 - Up to 1 ton!
- Largest maple species in North America (+ largest leaves)



3



WSU Extension Forestry

Identification

Leaves: large (8-12"), lustrous dark green above, pale beneath, 3-5 toothed lobes, petiole yields milky sap when detached, turn golden yellow in fall

Leaf/twig arrangement: Opposite

Bark: Grey-green, smooth when young, turning brown and furrowed with age, often covered in epiphytes

Flowers: male and female flowers in same greenish-yellow drooping clusters, develop into hairy double-samaras

Habitat/range: California to BC, moist soils with adequate drainage (floodplains, streambanks, drainage slopes), below 3000' elevation









- Produces seed at roughly 10 years old
- Monecious separate male and female flowers but on the same tree
- Flower in early spring
- Important pollinator species for bees, flies, and beetles
- Flowers are edible (ex: maple flower fritters)



7




WSU Extension Forestry

- Produces seed at roughly 10 years old
- Monecious separate male and female flowers but on the same tree
- Flower in early spring
- Important pollinator species for bees, flies, and beetles
- Flowers are edible (ex: maple flower fritters)
- Prolific seed producer
- Seeds are double samaras, descend like "helicopters"
- Seeds viable for one year





- Prolific "stump sprouter", common regeneration postharvest or fire
- Generally treated via herbicide post-harvest or else
- Rapid juvenile growth, outcompetes conifers
- Significant potential for coppice management
 - · Can produce dozens of stems
 - To maintain vigorous growth, select 3-5 stems to keep, remove the rest (no herbicide necessary)



9



WSU Extension Forestry

- Moderately shade tolerant
- Prefers at least small gaps in the canopy for regeneration and recruitment
- Can be a "seedling banker"
- Wide-spreading canopy can take up significant growing space
- Long-life and ability to regenerate in moderate shade makes it an important component of old growth forests







Sooty bark disease

- Caused by fungus Cryptostroma corticale
- "Sooty bark" refers to black mats of fungal spores
- Effects multiple maples species, including ornamentals
- Discovered in WA in 1968 but recent outbreaks linked to severe droughts
- Damage is typically restricted to stressed/weakened trees
- Spores can cause respiratory irritation
- https://foresthealth.org/sbd/

Damage and Disease Agents



11




WSU Extension Forestry

Bigleaf maple decline

- Observed since 2010
- Symptoms:
 - Smaller leaves
 - Chlorotic (yellowish) leaves and/or scorch
 - Crown thinning
 - Crown/branch dieback
 - Heavy seed corps
- Studies tied decline to drought/heat and stressful environments (forest edges, urban areas, landscapes, roads)
- No single pest or disease responsible










Ecological Role and Importance

- Typically grows in mixed stands but can be dominant canopy species
- Associated species: Doug-fir, grand fir, red alder, Sitka spruce, western redcedar, western hemlock, Pacific dogwood, serviceberry, salal, red huckleberry, salmonberry, sword fern, vine maple
- Can be present at any seral stage, but typically a mid or late successional species
- Important riparian species
 - Reduces soil and stream bank erosion
 - Enhances riparian habitats
 - Shade, woody debris, and nutrient inputs




15



WSU Extension Forestry

Ecological Role and Importance


- Importance to wildlife:
 - Favored browse species by deer and elk
 - Beaver feed on bark and use stems for dams and lodges
 - Birds and rodents feed on maple seeds
 - Nesting/cavity habitat
 - Preferred cavity tree for pileated woodpeckers (Hartwig et al. 2004)
 - Important pollinator species (early spring bloom)





Ecological Role and Importance

- Better Be-leaf it:
 - · Bigleaf maple "microsites"
 - Soils below maples have higher cationexchange capacity and concentrations of C, Ca, K, Mg, and N/nitrates (Hamdan & Schmidt 2012, Turk et al. 2007, Boyle et al. 1990)
 - Supporting stream habitat
 - Large amounts of detritus inputs into streams (including downed branches) tied to increased macroinvertebrate habitat (Kennedy et al. 2005)
 - Dense leaf litter layer provides habitat for insects and microfauna




17



WSU Extension Forestry

Ecological Role and Importance

- Heaven for Epiphytes:
 - Likely connected to the moisture retention of bigleaf maple bark
 - Common species: licorice fern, club moss, mountain fern moss, and several lichens (Clodonia, Nephroma, Crocynia)
 - Species hosted will vary depending on location on tree and microenvironment, can be indicators of air quality
 - Dense epiphyte communities on branches can serve as rooting medium for other plants and a food source for arthropods
 - Sometimes harvested for botanical products





Indigenous Uses:

- Bark used for rope
- Wood for utensils, instruments, and canoe paddles (called the "paddle tree")
- Leaves were used to wrap food and other items
- · Bark infusions to treat sore throats
- Shoots for basket-weaving
- Fuelwood
- Little or no evidence of sap collection ...
- *maples were often burned or pruned to force sprouting



19

WSU Extension Forestry

Uses/Products

Characteristics

- Moderate weight and hardness
- Fine grained
- Easily machined (planning, shaping, boring, turning)
- · Not suitable for outdoor uses

Wood products

- Veneer
- Furniture
- Paneling, flooring
- Plywood
- Musical instruments
- Molding
- Pulpwood/fuelwood
- Generally low-value wood except for . . .



WSU Extension Forestry **Agroforestry Opportunities** • Bigleaf Maple Syrup Produces sweet sap with 1-1.5% sugar · Can be processed into syrup and other sugar products • Sap season from Dec-March • UW/OSU/WSU research and outreach efforts underway Join the Sapsucker program! • forestry.wsu.edu/sapsuckers Other Forest Farming Opportunities • Light shade creates opportunities to grow "forest crops" Ex: hazelnuts, osoberry, currants, service berry, Oregon grape, cascara,

salal, sword fern, devil's club



- Not commonly planted
- Site selection can grow on a wide variety of soils, grows best on deep well-drained soils with abundant moisture (river terraces, flood plains, seepage areas)
- Site preparation None needed for vegetation reproduction, control needed if planting or in shaded areas
- Seedlings can be difficult to come by but relatively easy to propagate from seed
 - Collect and plant in fall, partial shade
- Recommended density 10-12' spacing (436-303 trees/acre)
- Underplanting in existing canopies
 - · Can establish in light shade
 - Small gaps can create rapid growth
- Browse protection!





23



WSU Extension Forestry

Management - Thinning and Harvesting

- Pre-commercial thinning and/or stump sprout management necessary to maintain growth
 - Likely around age 5-15
- If well-managed and on high quality sites, regeneration harvest with sawlog production could occur as early as 40-50 years
 - Very uncommon to manage for maple
- Volume per acre around 4500-5000 ft3 at 45-50 years
- Can be managed via clear cut, single-tree, small group, or potentially shelterwood systems



