



**ADULT AMBULATORY INFUSION ORDER**

**3/2023**

**Blood Transfusion**

**NAME:**  
**BIRTHDATE:**  
**INSURANCE:**  
**PROVIDER NAME:**  
**CLINIC NAME and Phone number:**

Patient identification

Weight: \_\_\_\_\_ kg Height: \_\_\_\_\_ cm Allergies: \_\_\_\_\_ Diagnosis Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_

**GUIDELINES FOR ORDERING:**

- 1) Patient name and date of birth must be on EACH page of this order form. This form will be returned if any left blank.
- 2) Send FACE SHEET and complete copy of INSURANCE information. Send H&P or most recent chart note, must be within the past 12 months.
- 3) Send a copy of related lab results if not available for review in Epic.

**Frequency of treatment: (must check one)**

- This is a ONE TIME order only, based on lab results included with this fax or available in Epic.
- These are STANDING ORDERS for PRN transfusions valid for one year from signature date for when patient's labs meet parameters to transfuse. Provider will fax lab results each time for our review if unavailable in Epic.

**Pre-Procedure: PARQ orders (must check one)**

- Patient consents to transfusion ONLY OF SELECTED blood products. Informed consent (PARQ) discussion completed.
- Patient consents to all blood products. Informed consent (PARQ) discussion completed.
- Patient consents to all blood products. PARQ discussion previously completed/documented.

**LABS:**

- ✓ Type and Screen (standing orders will be placed if PRN transfusions selected above).

**Transfusion Premeds: (select all that apply, to be given 30 minutes prior to transfusion)**

- Acetaminophen (TYLENOL) tablet, 650 mg, oral.
- Loratadine (CLARITIN) tablet, 10 mg, oral.
- OR
- DiphenhydrAMINE (BENADRYL) tablet, 25 mg, oral.
- OR
- DiphenhydrAMINE (BENADRYL) iv injection, 25 mg.

**Blood: (must fill in each blank to be a valid order)**

- ✓ **Red Blood Cells**-Transfuse \_\_\_\_\_ units (two-unit max), duration two hours per unit.  
Transfuse for Hgb equal to or less than \_\_\_\_\_.

**Platelets: (must fill in each blank to be a valid order)**

- ✓ **Platelets**-Transfuse amount \_\_\_\_\_ units, duration one hour per unit.  
Transfuse for Platelet level equal to or less than \_\_\_\_\_.

**Transfusion PRN Medications:**

- Furosemide (LASIX) injection \_\_\_\_ mg. Intravenous, once. When to give: \_\_\_\_\_.



ADULT AMBULATORY INFUSION ORDER

3/2023

Blood Transfusion

NAME:  
BIRTHDATE:  
INSURANCE:  
PROVIDER NAME:  
CLINIC NAME and Phone number:

Patient identification

**Standard included Nursing Orders:**

- ✓ Sodium chloride 0.9% (NS) infusion, at 10-20 ml/hr. Intravenous, CONTINUOUS PRN.
- ✓ VITAL SIGNS – Per policy.
- ✓ Insert peripheral IV if no IV access.
- ✓ May use/access CVC line and give 500 units (5ml) heparin injection for central line care.
- ✓ Hypersensitivity reaction order set. If hypersensitivity or infusion reactions develop, temporarily hold the infusion, initiate infusion reaction orders, and notify provider immediately.

**HYPERSENSITIVITY INFUSION REACTION ORDERS**

**STOP infusion if:**

**MILD REACTION:** Localized pruritus, rhinitis, localized rash, fever greater than 38 degrees C.

1. Stop infusion and maintain IV access with NS line.
2. Notify MD of symptoms and infusion reaction orders initiated.

**MODERATE REACTION:** Generalized pruritus, generalized flushing, rash, back pain, mild dyspnea, chills, hypotension with SBP less than 90 mmHg or greater than 20% decrease in SBP.

1. Stop infusion, but do not discard infusion bag/bottle unless instructed by MD that therapy is to be discontinued.
2. Maintain IV access with NS line.
3. Place patients in a supine position with lower extremities elevated unless respiratory distress or nausea/vomiting.
4. Maintain airway.

**SEVERE REACTION:** Respiratory distress (bronchospasm, stridor, wheezing, persistent cough, respiratory depression, cardiac arrhythmia, chest pain, generalized urticaria (hives), syncope, gastrointestinal symptoms (abdominal pain, N/V, diarrhea), hypotension SBP less than 80 mmHg, face/throat/tongue swelling, shock, loss of consciousness.

1. Stop infusion, but do not discard bag/bottle unless instructed by MD that therapy is to be discontinued.
2. Stay with patient and have another nurse notify MD of symptoms and infusion reaction orders initiated.
3. Place patients in supine position with lower extremities elevated unless respiratory distress or nausea/vomiting.
4. Maintain airway.
5. Maintain IV access with NS line.
6. Inpatients: Activate code blue for loss of blood pressure, pulse, or consciousness.

**MEDICATIONS:**

- Acetaminophen (TYLENOL) tablet 650 mg, Oral, EVERY 4 HOURS PRN, Pain, Other, Mild reaction. Starting when released.
- Diphenhydramine (Benadryl) injection, 25 mg, Intravenous, EVERY 15 MIN PRN, Moderate or severe reaction, may repeat x1. Starting when released, For 2 doses.
- Methylprednisolone sodium succinate (solu-MEDROL) 62.5mg/ml injection, 125 mg, Intravenous, ONCE PRN, Severe reaction. Starting when released, For 1 dose. Mix with 2 mL provided diluent to make 62.5 mg/mL.
- Famotidine (PEPCID) injection 20 mg, Intravenous, ONCE PRN, Moderate reaction, or anaphylaxis, starting when released. Dilute 2 mL of famotidine with 8 mL of normal saline to a final concentration of 2 mg/mL. Administer ordered dose over a period of at least 2 minutes.



**ADULT AMBULATORY INFUSION ORDER**

**3/2023**

**Blood Transfusion**

**NAME:**  
**BIRTHDATE:**  
**INSURANCE:**  
**PROVIDER NAME:**  
**CLINIC NAME and Phone number:**

Patient identification

- Albuterol 90 mcg/puff inhaler 2 puff, Inhalation, ONCE PRN, Severe reaction p. Starting when released, For 1 dose. Shake well. Use with spacer.
- Start oxygen. For *MODERATE* Reaction: Start 2 liters O2 per nasal cannula as needed for SpO2 less than 93%. For *SEVERE* Reaction: Start continuous pulse oximetry. Administer 6 to 8 liters per minute (LPM) via face mask, or up to 100% oxygen to maintain oxygen saturation greater than 90%.
- EPINEPHrine 1 mg/mL injection 0.3 mg. Intramuscular, EVERY 5 MIN PRN, Anaphylaxis. Starting when released, For 3 doses. Administer in the anterior lateral thigh using 1 - 1.5-inch needle for airway obstruction symptoms and/or hypotension.
- Sodium chloride 0.9% (NS) infusion 500ml. At 500 mL/hr., Intravenous, PRN, Moderate reaction, or anaphylaxis. Starting when released. Run wide open to gravity.

**By signing below, I represent the following:**

I am responsible for the care of the patient (*who is identified at the top of this form*).

I hold an active, unrestricted license to practice medicine in Oregon.

**My physician license Number is #** \_\_\_\_\_ and I am acting within my scope of practice and authorized by law to order infusion of the medications and blood products described above for the patient identified on this form.

**Provider's printed name:** \_\_\_\_\_

**Provider's signature:** \_\_\_\_\_

**Date:** \_\_\_\_\_

**Please complete the following intake questions.**

**Mobility status**

- Independent
- Walker
- Wheelchair
- Fully dependent

**Bariatric needs**

- Yes, briefly describe \_\_\_\_\_
- No

**Transportation concerns**

- Yes, briefly describe \_\_\_\_\_
- No

**Interpreter Service required**

- Yes, language \_\_\_\_\_
- No

**Outpatient Infusion Services Intake Team:**

Please check the appropriate box for the patient's preferred infusion center location:

- PORTLAND Phone-503-215-6046 Fax-503-487-3582
- WILLAMETTE FALLS Phone-503-215-6046 Fax-503-487-3582
- MEDFORD Phone-541-732-7000 Fax-541-732-3939
- HOOD RIVER Phone-541-387-6445 Call for fax #
- SEASIDE Phone-503-717-7650 Fax-503-470-3146
- NEWBERG Phone-503-537-6040 Fax-503-205-8720