



ADULT AMBULATORY INFUSION ORDER

3/2023

Infliximab (REMICADE & Biosimilars) Infusion

NAME:
BIRTHDATE:
INSURANCE:
PROVIDER NAME:
CLINIC NAME and Phone number:

Patient identification

Weight: _____ kg Height: _____ cm Allergies: _____ Diagnosis Code: _____

Height & weight are required for a complete order. Treatment Start Date: _____

****If treatment not started within 90 days, new orders with updated date/signature will be required. Otherwise, orders are valid for 1 year from signature date.****

GUIDELINES FOR ORDERING:

- 1) Patient name and date of birth must be on EACH page of this order form. This form will be returned if any left blank.
- 2) Send FACE SHEET and complete copy of INSURANCE information. Send H&P or most recent chart note, must be within the past 12 months.
- 3) Send a copy of related lab results if not available for review in Epic.

MEDICATION:

- Infliximab** in sodium chloride 0.9% 250 mL infusion, iv. Patient will be given biosimilar substitute (Renflexis, Avsola, Inflectra, Remicade) according to insurance coverage/preference. If insurance coverage allows, patient will receive drug based on pharmacy availability & the Providence Preferred Products List.
- Only check this box if it is NOT okay to substitute for insurance/Providence preferred product. Specific product required: _____, **specific reason why this is required:** _____.

REQUIRED Pre-screening labs: (Results must be available prior to initiation of therapy, attach copy if not available in Epic)

- QuantiFERON Gold TB (must be result prior to initiation of therapy).

Recommended Pre-screening labs: Send any applicable results from list below.

- Hepatitis C Antibody
- Hepatitis B surface antigen
- Hepatitis B core antibody total
- Hepatitis B surface antibody
- HIV

Optional LABS to be done at infusion appointments: (select all that apply)

- Comprehensive Metabolic Set, Routine, every _____
- CBC with differential, Routine, every _____
- Sedimentation Rate, Routine, every _____
- Quantiferon Gold, Routine, annually.
- Other: _____

Optional PRE-MEDICATIONS: (Select all that apply, to administer 30 minutes prior to infusion.)

- DiphenhydrAMINE (BENADRYL) tablet, 25 mg, oral, every visit. *Give either loratadine or diphenhydrAMINE, not both.*

OR

- Loratadine (CLARITIN) tablet, 10 mg, oral, every visit. *Give either loratadine or diphenhydrAMINE, not both.*
- Acetaminophen (TYLENOL) tablet, 650 mg, oral, every visit
- Other: _____



ADULT AMBULATORY INFUSION ORDER

3/2023

Infliximab (REMICADE & Biosimilars) Infusion

NAME:
BIRTHDATE:
INSURANCE:
PROVIDER NAME:
CLINIC NAME and Phone number:

Patient identification

DOSE & FREQUENCY:

Induction AND Maintenance doses:

Give _____mg/kg. To infuse on weeks 0, 2, and 6. After week 6, maintenance doses to be given every _____ weeks.

OR

Maintenance doses ONLY: *To continue treatment with ongoing maintenance orders. If infusions received elsewhere, **date of last infusion here:** _____.*

Give _____mg/kg every _____ weeks.

AS NEEDED MEDICATIONS:

- ✓ Sodium chloride 0.9% (NS) bolus 250 ml, Flush before and after IV medication(s) and as needed with NS.

Standard included Nursing Orders:

- ✓ Administer over a period of not less than 2 hours. Use 0.22-micron filter for administration. Rate for first infusion: 10 mL/hr. for 15 min, then 20 mL/hr. for 15 min, then 40 mL/hr. for 15 min, then 80 mL/hr. for 15 min, then 150 mL/hr. for 15 min, then 250 mL/hr.
- ✓ FIRST VISIT: Confirm that prescreening tests are normal or have been addressed by Licensed Independent Practitioner. Notify provider before starting infusion if patient has signs of active infection or history of reaction to previous infusion that has not been addressed by LIP.
- ✓ Remind patient to avoid live vaccines.
- ✓ Insert peripheral IV if no iv access.
- ✓ May use/access CVC line and give 500 units (5ml) heparin injection for central line care.
- ✓ If hypersensitivity or infusion reactions develop, temporarily hold the infusion, and notify provider immediately. Include hypersensitivity reaction order set. (See attached form).

HYPERSENSITIVITY INFUSION REACTION ORDERS

STOP infusion if:

MILD REACTION: Localized pruritus, rhinitis, localized rash, fever greater than 38 degrees C.

1. Stop infusion and maintain IV access with NS line.
2. Notify MD of symptoms and infusion reaction orders initiated.

MODERATE REACTION: Generalized pruritus, generalized flushing, rash, back pain, mild dyspnea, chills, hypotension with SBP less than 90 mmHg or greater than 20% decrease in SBP.

1. Stop infusion, but do not discard infusion bag/bottle unless instructed by MD that therapy is to be discontinued.
2. Maintain IV access with NS line.
3. Place patients in a supine position with lower extremities elevated unless respiratory distress or nausea/vomiting.
4. Maintain airway.

SEVERE REACTION: Respiratory distress (bronchospasm, stridor, wheezing, persistent cough, respiratory depression, cardiac arrhythmia, chest pain, generalized urticaria (hives), syncope, gastrointestinal symptoms



ADULT AMBULATORY INFUSION ORDER

3/2023

Infliximab (REMICADE & Biosimilars) Infusion

NAME:
BIRTHDATE:
INSURANCE:
PROVIDER NAME:
CLINIC NAME and Phone number:

Patient identification

(abdominal pain, N/V, diarrhea), hypotension SBP less than 80 mmHg, face/throat/tongue swelling, shock, loss of consciousness.

1. Stop infusion, but do not discard bag/bottle unless instructed by MD that therapy is to be discontinued.
2. Stay with patient and have another nurse notify MD of symptoms and infusion reaction orders initiated.
3. Place patients in supine position with lower extremities elevated unless respiratory distress or nausea/vomiting.
4. Maintain airway.
5. Maintain IV access with NS line.
6. Inpatients: Activate code blue for loss of blood pressure, pulse, or consciousness.

MEDICATIONS:

- Acetaminophen (TYLENOL) tablet 650 mg, Oral, EVERY 4 HOURS PRN, Pain, Other, Mild reaction. Starting when released.
- DiphenhydrAMINE (Benadryl) injection, 25 mg, Intravenous, EVERY 15 MIN PRN, Moderate or severe reaction, may repeat x1. Starting when released, For 2 doses.
- Methylprednisolone sodium succinate (solu-MEDROL) 62.5mg/ml injection, 125 mg, Intravenous, ONCE PRN, Severe reaction. Starting when released, For 1 dose. Mix with 2 mL provided diluent to make 62.5 mg/mL.
- Famotidine (PEPCID) injection 20 mg, Intravenous, ONCE PRN, Moderate reaction, or anaphylaxis, starting when released. Dilute 2 mL of famotidine with 8 mL of normal saline to a final concentration of 2 mg/mL. Administer ordered dose over a period of at least 2 minutes.
- Albuterol 90 mcg/puff inhaler 2 puff, Inhalation, ONCE PRN, Severe reaction p. Starting when released, For 1 dose. Shake well. Use with spacer.
- Start oxygen. For *MODERATE* Reaction: Start 2 liters O2 per nasal cannula as needed for SpO2 less than 93%. For *SEVERE* Reaction: Start continuous pulse oximetry. Administer 6 to 8 liters per minute (LPM) via face mask, or up to 100% oxygen to maintain oxygen saturation greater than 90%.
- EPINEPHrine 1 mg/mL injection 0.3 mg. Intramuscular, EVERY 5 MIN PRN, Anaphylaxis. Starting when released, For 3 doses. Administer in the anterior lateral thigh using 1 - 1.5-inch needle for airway obstruction symptoms and/or hypotension.
- Sodium chloride 0.9% (NS) infusion 500ml. At 500 mL/hr., Intravenous, PRN, Moderate reaction, or anaphylaxis. Starting when released. Run wide open to gravity.

By signing below, I represent the following:

I am responsible for the care of the patient (*who is identified at the top of this form*).

I hold an active, unrestricted license to practice medicine in Oregon.

My physician license Number is # _____ **and I am acting within my scope of practice and authorized by law to order infusion of the medications and blood products described above for the patient identified on this form.**

Provider's printed name: _____

Provider's signature: _____

Date: _____



ADULT AMBULATORY INFUSION ORDER

3/2023

Infliximab (REMICADE & Biosimilars) Infusion

NAME:
BIRTHDATE:
INSURANCE:
PROVIDER NAME:
CLINIC NAME and Phone number:

Patient identification

Please complete the following intake questions.

Mobility status

- Independent
- Walker
- Wheelchair
- Fully dependent

Bariatric needs

- Yes, briefly describe _____
- No

Transportation concerns

- Yes, briefly describe _____
- No

Interpreter Service required

- Yes, language _____
- No

Outpatient Infusion Services Intake Team:

Please check the appropriate box for the patient's preferred infusion center location:

- | | |
|---|-------------------------------------|
| <input type="checkbox"/> PORTLAND | Phone-503-215-6046 Fax-503-487-3582 |
| <input type="checkbox"/> WILLAMETTE FALLS | Phone-503-215-6046 Fax-503-487-3582 |
| <input type="checkbox"/> MEDFORD | Phone-541-732-7000 Fax-541-732-3939 |
| <input type="checkbox"/> HOOD RIVER | Phone-541-387-6445 Call for fax # |
| <input type="checkbox"/> SEASIDE | Phone-503-717-7650 Fax-503-470-3146 |
| <input type="checkbox"/> NEWBERG | Phone-503-537-6040 Fax-503-205-8720 |