

Policy and Procedure	
MEDICAL PRIOR AUTHORIZATION POLICY AND CRITERIA ORPTCNEU029.12/20	NEUROMUSCULAR DRUGS Botulinum Toxin Botox® (onabotulinumtoxinA) Dysport® (abobotulinumtoxinA) Jeuveau® (prabotulinumtoxinA-xvfs) Myobloc® (rimabotulinumtoxinB) Xeomin® (incobotulinumtoxinA)
Effective Date: 2/1/2021  Robert Gluckman, M.D. Chief Medical Officer	Review/Revised Date: 05/19, 08/19, 12/19, 03/20, 05/20, 07/20 (jls), 9/20 (ss), 11/20 (JLS)
	P&T Committee Meeting Date: 06/19, 08/19, 10/19, 12/19, 02/20, 06/20, 08/20, 12/20
	Original Effective Date: 09/19
	Approved by: Oregon Region Pharmacy and Therapeutics Committee
	Page: 1 of 9

SCOPE:

Providence Health Plan, Providence Health Assurance, Providence Plan Partners, and Ayin Health Solutions as applicable (referred to individually as “Company” and collectively as “Companies”).

APPLIES TO:

Commercial
Medicaid

POLICY CRITERIA:

COVERED USES:

All Food and Drug Administration (FDA) approved and selected medically accepted indications not otherwise excluded from the benefit, as outlined below.

Coverage for Medicaid is limited to a condition that has been designated a covered line item number by the Oregon Health Services Commission listed on the Prioritized List of Health Care Services.

CRITERIA:

OnabotulinumtoxinA (Botox®) may be covered for the following indications when criteria are met:

1. Chronic migraine headaches in adults when all of the following is met:
 - a. Documentation of at least 15 headache days per month with headaches lasting 4 hours or longer
 - b. Documentation of trial and failure, intolerance, or contraindication to at least TWO of the following classes used for migraine prevention. Trial and

failure is defined as inadequate response following a minimum three (3) months of consistent use.

- i. Antidepressants (e.g., amitriptyline, venlafaxine)
 - ii. Beta-blockers (e.g., metoprolol, propranolol, timolol)
 - iii. Antiepileptics (e.g., divalproex, valproate, topiramate)
 - c. Combination therapy with Calcitonin Gene-Related Peptide (CGRP) Receptor Antagonists (e.g., Aimovig®, Emgality®, Ajovy®) is considered **investigational and is not covered**
2. Upper and lower limb spasticity in patients at least 2 years of age
3. Cervical dystonia in adults
4. Strabismus and blepharospasm associated with dystonia in patients at least 12 years of age
5. Severe axillary hyperhidrosis in adults after documented trial and failure, intolerance or contraindication to topical agents
 - a. Note: The safety and effectiveness of onabotulinumtoxinA for hyperhidrosis in other body areas have not been established.
6. Overactive bladder in adults with:
 - a. Symptoms of urge urinary incontinence, urgency, and frequency
 - b. Documented trial and failure, intolerance, or contraindication to at least one month of anticholinergic medication (e.g., oxybutynin, tolterodine)
7. Urinary incontinence in adults:
 - a. Due to detrusor over activity related to a neurologic condition (e.g., spinal cord injury, multiple sclerosis)
 - b. Documented trial and failure, intolerance, or contraindication at least one month of anticholinergic medication (e.g., oxybutynin, tolterodine)
8. Excessive salivation due to advanced Parkinson's disease
9. Hemifacial spasm
10. Chronic anal fissure when all of the following is met:
 - a. Prescribed by, or in consultation with, a gastroenterologist or colorectal surgeon
 - b. Documentation of trial and failure, intolerance, or contraindication to at least 6 weeks of therapy with either topical nitrates or topical calcium channel blockers
 - c. One of the following:
 - i. Documentation that the patient is not a good candidate for surgery or appropriate medical rationale is provided for avoiding surgery
 - ii. Botox® is to be used in conjunction with fissurotomy
 - d. The use of Botox® in combination with sphincterotomy or anal advancement flap is considered **experimental and investigational** and will not be covered

**MEDICAL PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCNEU029**

**NEUROMUSCULAR DRUGS
Botulinum Toxin**

AbobotulinumtoxinA (Dysport®) may be covered for the following indications:

1. Spasticity in adults
2. Cervical dystonia in adults
3. Upper and lower limb spasticity in patients at least 2 years of age
4. Blepharospasm in adults

IncobotulinumtoxinA (Xeomin®) may be covered for the following indications:

1. Chronic sialorrhea in adult patients
2. Upper limb spasticity in patients at least two years of age
3. Cervical dystonia in adults
4. Blepharospasm in adults

RimabotulinumtoxinB (Myobloc®) may be covered for the following indications:

1. Cervical dystonia in adults
2. Chronic sialorrhea in adult patients

EXCLUSION CRITERIA:

- When the above criteria are not met, botulinum toxin is considered **investigational and not covered**.
- Botulinum toxin is considered **cosmetic and is not covered** for the treatment of glabellar lines and/or fine wrinkles on the face.
 - PrabotulinumtoxinA (Jeuveau®) will **not be covered** as it is only FDA approved for the treatment of glabellar lines and/or fine wrinkles on the face.

REQUIRED MEDICAL INFORMATION:

For initiation of treatment, a prior authorization form and relevant chart notes documenting medical rationale are required and for continuation of therapy, ongoing documentation of successful response to the medication may be necessary.

AGE RESTRICTIONS: N/A

PRESCRIBER RESTRICTIONS: N/A

COVERAGE DURATION:

Initial authorization and reauthorization will be approved for one year

Requests for a non-FDA approved (off-label) indication requires the proposed indication be listed in either the American Hospital Formulary System (AHFS), Drugdex, or the National Comprehensive Cancer Network (NCCN) and is considered subject to evaluation of the prescriber's medical rationale, formulary alternatives, the available published evidence-based research and whether the proposed use is determined to be experimental/investigational.

**MEDICAL PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCNEU029**

**NEUROMUSCULAR DRUGS
Botulinum Toxin**

Coverage for Medicaid is limited to a condition that has been designated a covered line item number by the Oregon Health Services Commission listed on the Prioritized List of Health Care Services.

Coverage decisions are made on the basis of individualized determinations of medical necessity and the experimental or investigational character of the treatment in the individual case

INTRODUCTION:

Botulinum toxin injections are used to treat various focal muscle spastic disorders and excessive muscle contractions such as dystonias, spasms, twitches, etc. These drugs produce a presynaptic neuromuscular blockade by preventing the release of acetylcholine from the nerve endings. The resulting chemical denervation of muscle produces local paresis or paralysis and allows individual muscles to be weakened selectively. Botulinum toxins have the advantage of being potent neuromuscular blocking agents with good selectivity and duration of action.

Botulinum toxins types A and B are neurotoxins produced by Clostridium Botulinum. Botulinum Toxin Type A and Botulinum Toxin Type B have many similarities and as experience has been gained, medical consensus has gradually developed that the two toxins have similar, but not identical, properties. Each botulinum toxin product is pharmacologically and clinically distinct, and therefore, not interchangeable with any other botulinum toxin product. As a result, approved indications for the two toxins differ.

The rationale for treatment is to create temporary paralysis of sufficient depth and duration that the injected muscles become slightly atrophied and stretched. The antagonist muscle shortens simultaneously taking up the slack created by agonist paralysis. After several weeks enervation to the injected muscle returns. The safety and efficacy of long term use of Botox, Myobloc, Dysport or Xeomin is unknown.

FDA APPROVED INDICATIONS:

Botox® (onabotulinumtoxinA)

- Bladder Dysfunction
- Chronic Migraine
- Adult upper and lower limb spasticity
- Pediatric upper limb spasticity (2 to 17 years of age)
- Pediatric lower limb spasticity (2 to 17 years of age), excluding spasticity caused by cerebral palsy
- Cervical dystonia
- Primary axillary hyperhidrosis
 - Limitations of use:

- Safety and effectiveness for hyperhidrosis in other body areas have not been established. Weakness of hand muscles and blepharoptosis may occur in patients who receive treatment for palmar hyperhidrosis and facial hyperhidrosis, respectively. Patients should be evaluated for potential causes of secondary hyperhidrosis (e.g., hyperthyroidism) to avoid symptomatic treatment of hyperhidrosis without the diagnosis and/or treatment of the underlying disease
- Safety and effectiveness have not been established for the treatment of axillary hyperhidrosis in pediatric patients under age 18
- Blepharospasm and strabismus

Dysport® (abobotulinumtoxinA)

- Glabellar Lines
- Cervical dystonia
- Spasticity in adults
- Lower limb spasticity in pediatric patients 2 years of age and older
- Upper limb spasticity in pediatric patients 2 years of age and older, excluding spasticity caused by cerebral palsy

Myobloc® (rimabotulinumtoxinB)

- Cervical dystonia
- Chronic sialorrhea in adults

Xeomin® (incobotulinumtoxinA)

- Chronic Sialorrhea in adult patients
- Glabellar Lines
- Cervical dystonia in adult patients
- Upper limb spasticity in adult patients
- Upper limb spasticity in pediatric patients 2 to 17 years of age, excluding spasticity caused by cerebral palsy
- Blepharospasm in adult patients

Jeuveau® (prabotulinumtoxinA-xvfs)

- Glabellar Lines

POSITION STATEMENT:

The safety and efficacy of long term use of Botox, Myobloc, Dysport or Xeomin is unknown.

The policy criteria was developed based on medically accepted indications for the specific products. Medically accepted refers to FDA approved or generally recognized as efficacious by certain drug compendia references (e.g., DrugDex, AHFS).

Migraine Headache Prophylaxis

The [2018 American Headache Society \(AHS\) Consensus Statement](#) uses the International Classification of Headache Disorders definition of chronic migraines as follows:

- A. Migraine-like or tension-type-like headache on ≥ 15 days/month for >3 months that fulfill criteria B and C
- B. Occurring in a patient who has had at least 5 attacks fulfilling criteria B-D for migraine without aura and/or criteria B and
- C. C for migraine with aura
- D. On ≥ 8 days/month for >3 months, fulfilling any of the following:
 1. Criteria C and D migraine without aura
 2. Criteria B and C for migraine with aura
 3. Believed by the patient to be migraine at onset and relieved by a triptan or ergot derivative
- E. Not better accounted for by another diagnosis

Furthermore, the AHS recommends initiation of prophylactic treatment when any of the following occurs:

- Attacks significantly interfere with patients' daily routines despite acute treatment
- Frequent attacks (≥ 4 monthly headache days)
- Contraindication to, failure, or overuse of acute treatments, with overuse defined as:
 - 10 or more days per month for ergot derivatives, triptans, opioids, combination analgesics, and a combination of drugs from different classes that are not individually overused
 - 15 or more days per month for non-opioid analgesics, acetaminophen, and nonsteroidal anti-inflammatory drugs
- Adverse events with acute treatments
- Patient preference

Treatments with established efficacy for migraine prophylaxis include antiepileptic drugs (*i.e.*, divalproex sodium, valproate sodium, topiramate), beta-blockers (*i.e.*, metoprolol, propranolol, and timolol), triptans (*i.e.*, frovatriptan for short-term prevention of menstrual migraines), onabotulinumtoxinA, and calcitonin gene-related peptide (CGRP) receptor antagonists [*i.e.*, erenumab (Aimovig®), fremanezumab (Ajovy®), and galcanezumab (Emgality®)].

CGRP receptor antagonists are a newer class of medications indicated for migraine prophylaxis. The clinical trials for these agents excluded patients that were currently using botulinum toxin. There is currently no clinical information to support use of combination therapy. The AHS statement includes recommendations that adding on injectable CGRP therapy to oral preventive therapies is appropriate given the lack of drug-drug interactions.

Chronic Anal Fissure

The [2014 American College of Gastroenterology guidelines](#) outline the diagnosis and management of chronic anal fissure. They define anal fissure as an “ulcer-like, longitudinal tear in the midline of the anal canal” and define chronic as lasting more than 8-12 weeks with edema and fibrosis.

Treatment typically consists of topical agents targeting the relief of spasms, such as topical nitrates, such as 0.2% nitroglycerin ointment applied twice for 6-8 weeks, and topical calcium channel blockers (CCBs), such as 2% diltiazem applied twice daily for 6–8 weeks. While these therapies are minimally invasive and typically inexpensive, the rate of healing are considered only marginally better than placebo for nitrates, to have “insufficient data to conclude whether [topical CCBs] are superior to placebo.” Botulinum toxin has healing rates superior than placebo (60-80%) and patients may be retreated on relapse with similar healing rates. Patients considered medical refractory to these treatments should be referred for surgical evaluation. Lateral internal sphincterotomy (LIS) is preferred over manual anal dilation due to better outcomes and less incontinence adverse effects. However, there is still a risk of incontinence, so topical and injectable therapies continue to be used, despite better efficacy with LIS. A systematic review found that LIS was statistically significantly better in healing rates and recurrence rates than Botox®

Recommendation: “Gastroenterologists and other providers should refer patients who do not respond to conservative or pharmacologic treatment for local injections of botulinum toxin (strong recommendation, low quality of evidence) or surgical internal anal sphincterotomy (strong recommendation, high quality of evidence).”

CPT/HCPCS Codes

All Lines of Business Except Medicare	
Prior Authorization Required	
31513	Laryngoscopy, indirect; with vocal cord injection
31570	Laryngoscopy, direct, with injection into vocal cord(s), therapeutic
31571	Laryngoscopy, direct, with injection into vocal cord(s), therapeutic; with operating microscope or telescope

**MEDICAL PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCNEU029**

**NEUROMUSCULAR DRUGS
Botulinum Toxin**

31573	Laryngoscopy, flexible; with therapeutic injection(s) (eg, chemodenervation agent or corticosteroid, injected percutaneous, transoral, or via endoscope channel), unilateral
43192	Esophagoscopy, rigid, transoral; with directed submucosal injection(s), any substance
43201	Esophagoscopy, flexible, transoral; with directed submucosal injection(s), any substance
43236	Esophagogastroduodenoscopy, flexible, transoral; with directed submucosal injection(s), any substance
46505	Chemodenervation of internal anal sphincter
52287	Cystourethroscopy, with injection(s) for chemodenervation of the bladder
64611	Chemodenervation of parotid and submandibular salivary glands, bilateral
64612	Chemodenervation of muscle(s); muscle(s) innervated by facial nerve, unilateral (eg, for blepharospasm, hemifacial spasm)
64615	Chemodenervation of muscle(s); muscle(s) innervated by facial, trigeminal, cervical spinal and accessory nerves, bilateral (eg, for chronic migraine)
64616	Chemodenervation of muscle(s); neck muscle(s), excluding muscles of the larynx, unilateral (eg, for cervical dystonia, spasmodic torticollis)
64617	Chemodenervation of muscle(s); larynx, unilateral, percutaneous (eg, for spasmodic dysphonia), includes guidance by needle electromyography, when performed
64642	Chemodenervation of one extremity; 1-4 muscle(s)
64643	Chemodenervation of one extremity; each additional extremity, 1-4 muscle(s) (List separately in addition to code for primary procedure)
64644	Chemodenervation of one extremity; 5 or more muscles
64645	Chemodenervation of one extremity; each additional extremity, 5 or more muscles (List separately in addition to code for primary procedure)
64646	Chemodenervation of trunk muscle(s); 1-5 muscle(s)
64647	Chemodenervation of trunk muscle(s); 6 or more muscles
64650	Chemodenervation of eccrine glands; both axillae
64653	Chemodenervation of eccrine glands; other area(s) (eg, scalp, face, neck), per day
67345	Chemodenervation of extraocular muscle
95873	Electrical stimulation for guidance in conjunction with chemodenervation (List separately in addition to code for primary procedure)
95874	Needle electromyography for guidance in conjunction with chemodenervation (List separately in addition to code for primary procedure)
J0585	Injection, onabotulinumtoxina, 1 unit
J0586	Injection, abobotulinumtoxina, 5 units
J0587	Injection, rimabotulinumtoxinb, 100 units
J0588	Injection, incobotulinumtoxin a, 1 unit
J3590	Injection, prabotulinumtoxin a (dump code)
S2340	Chemodenervation of abductor muscle(s) of vocal cord
S2341	Chemodenervation of adductor muscle(s) of vocal cord

**MEDICAL PRIOR AUTHORIZATION
POLICY AND CRITERIA
ORPTCNEU029**

**NEUROMUSCULAR DRUGS
Botulinum Toxin**

Unlisted Codes

All unlisted codes will be reviewed for medical necessity, correct coding, and pricing at the claim level. If an unlisted code is billed related to services addressed in this policy then **prior-authorization is required.**

31599	Unlisted procedure, larynx
43499	Unlisted procedure, esophagus
64999	Unlisted procedure, nervous system

REFERENCE/RESOURCES:

1. Botox package insert. Irvine, CA; Allergan Pharmaceuticals; 2018 Sept.
2. Dysport package insert.
3. Xeomin package insert. Franksville, WI; Maerz Aesthetics; 2020 Sept.
4. Myobloc package insert. Louisville, KY; Solstice Neurosciences; 2019 Oct.
5. DRUGDEX® System [Internet database]. Greenwood Village, Colo: Thomson Reuters (Healthcare) Inc. Updated periodically.
6. American Headache Society. The American Headache Society Position Statement On Integrating New Migraine Treatments Into Clinical Practice. Available at <https://headachejournal.onlinelibrary.wiley.com/doi/epdf/10.1111/head.13456> (Accessed May 7, 2020)
7. American College of Gastroenterology (ACG). ACG Clinical Guideline: Management of Benign Anorectal Disorders. Available at https://journals.lww.com/ajg/Fulltext/2014/08000/ACG_Clinical_Guideline_Management_of_Benign.7.aspx (Accessed July 16, 2020)
8. Shao WJ Li GC, Zhang ZK. Systematic review and meta-analysis of randomized controlled trials comparing botulinum toxin injection with lateral internal sphincterotomy for chronic anal fissure *Int J Colorectal Dis*. 2009 Sep;24(9):995-1000.
9. Sileri P, Stolfi VM, Franceschilli L et al. Conservative and surgical treatment of chronic anal fissure: prospective longer term results. *J Gastrointest Surg*. 2010 May;14(5):773-80.