Engelman, R., D. Pauly, D. Zeller, R.G. Prinn, J.K. Pinnegar and N.V.C. Polunin. 2008. Introduction: Climate, people, fisheries and aquatic ecosystems, p. 1-15 *In:* N.V.C. Plunin (ed.) *Aquatic Ecosystems: Trends and Global Prospects*. Cambridge University Press, New York.

1 • Introduction: Climate, people, fisheries and aquatic ecosystems

ROBERT ENGELMAN, DANIEL PAULY, DIRK ZELLER, RONALD G. PRINN, JOHN K. PINNEGAR AND NICHOLAS V. C. POLUNIN



INTRODUCTION TO THE BOOK

Evidence of human damage to natural resources and the environment is long-standing even in the sea, but some 50 years ago awareness of human degradation of natural environments around the globe grew substantially. This concern was expressed above all in the creation of protected areas, organizations and agencies focused on nature conservation. The extent to which wilderness areas everywhere were contaminated or otherwise influenced by human agency was beginning to be generally recognized, and some people predicted these impacts would only grow in future. But few predicted the extent and number of global environmental changes that are now occurring. The limits to growth of the world's economies, individually or

Aquatic Ecosystems, ed. N. V. C. Polunin. Published by Cambridge University Press. © Foundation for Environmental Conservation 2008.

collectively, were little questioned in ecological contexts. Uncertainties surrounding how economics and ecology might relate to each other were little explored, and questions such as 'At what points will economies become constrained by the decline and loss of natural ecological goods and services?' persist to this day.

Water is increasingly seen as a constraint on sustainable human development and focus of potential human conflict. Nevertheless, it has been rare for the provision of this good, and related ecological services such as fish production and waste disposal, to be viewed holistically in the context of the natural environment. Also, there have been no comprehensive reviews of status and trends encompassing all aquatic environments (Clark et al. 2006), from the fresh and terrestrial saline to those of the deepest seas. Certain fresh waters and saline lakes have been radically altered by humans, but the extent and nature of these impacts vary among ecosystem types and geographical locations. The seas and oceans retain a greater complement of resource-development frontiers than does the land, for example in mining and fisheries (e.g. Berkes et al. 2006); their wilderness value remains substantial. Clearly, the extent of human disturbance is increasing across environment types and locations, vet there is little comprehensive scientific appraisal of the magnitude and nature of this permeation.

It is important at this time to consider what is happening ecologically to the world's aquatic environments, and where possible project these forward to an appropriate time horizon at an appropriate ecological scale. The terminology surrounding ecological units is large and application can be problematic especially where, as here, freshwater and marine environments are to be bridged. Thus the term 'biome' is discerning of terrestrial environments but not with respect to those of the sea, while 'ecoregions' (e.g. Bailey 1998) omit the continental shelves. It was important in the design of this book to focus on a number of natural environments that would be comprehensive yet feasible in terms of relevant data and expertise. One aim of this chapter is to consider alternative comprehensive categorizations of aquatic environments as a basis for this book.

Natural processes and ecological services rely on notions of system functioning that have long been integral to the concept of the 'ecosystem'. Other recognized types of ecological unit such as the biotope exist, but the term 'ecosystem' is the most relevant, and most widely used and recognized, and is thus the unit of choice for this book. What was also needed in the design of this book was a time horizon sufficiently far in the future to provoke thoughtful and imaginative projections across all the ecosystems, but within a time-frame for which some detailed scenarios such as of human population growth and global ambient temperatures existed. The year 2025 was chosen as an appropriate compromise between those two constraints. This book aims to assemble the views of expert ecologists on the status of all ecosystems across the aquatic realm, review contemporary changes and their drivers, and consider likely outcomes at the 2025 time horizon. Major drivers considered are climate change and the direct impacts of human population growth and economic development.

Climate is usefully defined as the average of the weather experienced over a 10-20-year period. Temperature and rainfall changes are typical measures of change that can be expressed at local, regional, national or global scales. Global warming or cooling can be driven by any imbalance between the solar energy the Earth receives from the Sun and the energy it radiates back to space as invisible infrared light. The greenhouse effect is a warming influence caused by the presence of gases and clouds in the atmosphere, which are very efficient absorbers and radiators of this infrared light. The greenhouse effect is opposed by substances at the Earth's surface (such as snow and desert sand) and in the atmosphere (such as clouds and white aerosols) that efficiently reflect sunlight back into space (albedo) and are thus a cooling influence. Global climate changes and their possible consequences are and will remain a major area of debate within and among most nations. International negotiations in the Framework Convention on Climate Change and its Kyoto Protocol are contentious in particular because the components of the problem are many, the science is uncertain and the whole issue has a major bearing on other global problems such as poverty, inequality and economic development. Another aim of this introductory chapter is to consider uncertainties in climate predictions and review warming, sea-level rise and related trends as they might affect the world's aquatic environments.

The human population is approaching the 7 billion mark, increasing by c.1.2% per year. Mapping and photography of the Earth's night lights (Weier 2000) show how unevenly human beings are settled (Deichmann *et al.* 2001). Especially dense populations characterize northern South Asia, East Asia, Europe, eastern North America and south-western Africa. More than half of the world's population lives within 60 km of the coast, and this figure could reach 75% by the year 2020 (Roberts & Hawkins 1999). In addition, human economic wealth and output are

often concentrated close to major rivers or estuaries, especially in the temperate regions of the northern hemisphere (Cohen 1997; Nicholls & Small 2002). Various human interactions with critical natural resources and the environment illustrate that the human population over the past several decades has reached levels that significantly modify aquatic ecosystems. While some of this change is economically and socially beneficial, some is ecologically detrimental (e.g. biodiversity loss; see below). A third aim of this chapter is to introduce a human backdrop for assessing global changes in aquatic ecosystems; this will be done in particular by reviewing some of the consequences of human population growth for water, water supply and its natural sources.

Fisheries have been recognized as a major driver of change in the aquatic realm. They constitute a major impact at the interfaces among human population growth, economic development and environmental changes, both anthropogenic and natural. Modern industrial fisheries resulted from the economic and technological development of Europe, North America and Japan over a century ago. Frontiers in the development of global fishery resources persist in the use of Southern Ocean and deep-sea or reef resources. What are the current trends in global fisheries, and how do these relate to what is happening in the aquatic ecosystems involved? The intention in this chapter is also to introduce marine fisheries as an important example of human intrusion into the oceans and seas; this is followed by an overview of global trends in aquatic biodiversity and specific extinctions.

COMPLEXITIES OF CLIMATE CHANGE AND ITS CONSEQUENCES

Global climate has remained relatively stable (temperature changes of <1 °C over a century) during the last 10000 years. However, society now faces potentially rapid changes because human activities have altered the atmosphere's composition and changed the Earth's radiation balance (IPCC [Intergovernmental Panel on Climate Change] 1996).

The most important greenhouse gas is water vapour, which typically remains for a week or so in the atmosphere. Concerns about global warming, however, revolve around much longer-lived greenhouse gases, especially carbon dioxide but also other long-lived gases (methane, nitrous oxide, chlorofluorocarbons), concentrations of which have increased substantially since c.1750. Carbon dioxide (CO₂) has risen by about 30%, methane (CH₄) by more than

100% and nitrous oxide (N₂O) by about 15%. These gases are now at higher concentrations than at any time in the past 160 000 years (IPCC 1996). The combustion of fossil fuels and, to a lesser extent, changes in land use account for anthropogenic CO₂ emissions. Agriculture is responsible for nearly 50% of human-generated CH₄ emissions and about 70% of anthropogenic N₂O emissions.

When the concentration of a particular greenhouse gas increases, it tends to lower the flow of infrared energy to space and increases the flow of infrared energy down toward the surface. The Earth is then receiving more energy than it radiates to space. This 'radiative forcing' warms the surface and the lower atmosphere; however, the rate of surface warming is slowed by uptake of heat by the world's oceans. The greenhouse effect as quantified by this radiative forcing is real and the physics relatively well understood. What is more uncertain, and the cause of much of the scientific debate, is the response of the global system that determines climate to this radiative forcing. Feedbacks in the system can either amplify or dampen the response in ways that are still only partially understood.

The IPCC was jointly established by the World Meteorological Organization and the United Nations Environment Programme (UNEP) in 1988, in order to (1) assess available scientific information on climate change, (2) assess the environmental and socioeconomic impacts of climate change, and (3) formulate response strategies. The Integrated Global System Model (IGSM) is one means that the IPCC has at its disposal to analyse carefully the scientific and economic implications of proposed mitigation policies (Fig. 1.1). The IGSM consists of a set of coupled submodels of economic development and associated emissions, natural biogeochemical cycles, climate and natural ecosystems (Prinn et al. 1999; MIT [Massachusetts Institute of Technology] 2003). It attempts to include each of the major areas in the natural and social sciences that are amenable to quantitative analysis and are relevant to the issue of climate change (Schneider 1992; Prinn & Hartley 1992; IPCC 2001a, b, c). A major challenge inherent in global climate modelling is to decide what is important and what is unimportant. In the IGSM, the coupled atmospheric chemistry and climate model (Fig. 1.1) is driven by a combination of anthropogenic and natural emissions. The essential components of this model are chemistry, atmospheric circulation and ocean circulation, each of which, by itself, can require enormous computer resources. The atmospheric chemistry is modelled in sufficient detail to capture its sensitivity to climate and different mixes of emissions, and to address the




Fig. 1.1. Schematic illustration of the framework and processes of the MIT Integrated Global System Model (IGSM). Feedbacks between the component models that are currently included or proposed for inclusion in the next generation are shown as solid and dashed lines respectively (Prinn *et al.* 1999; MIT 2003).

relationship among policies proposed for control of emissions related to air pollution, aerosols and greenhouse gases (Wang *et al.* 1998). But linking complex models together leads to many challenges, illustrated by the failure of coupled ocean–atmosphere models (including IGSM) to accurately simulate current climate without arbitrary adjustments. The IGSM coupled chemistry–climate model outputs drive a terrestrial ecosystems model that predicts vegetation changes, land CO_2 fluxes and soil composition (Xiao *et al.* 1998), and these feed back to the climate model, chemistry model and natural emissions model. The effects of changes in land cover and surface albedo on climate, and the effects of changes in climate and ecosystems on agriculture and anthropogenic emissions are also included. Hundreds of runs of the IGSM have been used to quantitatively assess uncertainty in climate projections (Webster *et al.* 2002, 2003). One study shows a global median surface temperature rise from 1990 to 2100 of $2.4 \,^{\circ}$ C, with a 95% confidence interval of $1.0-4.9 \,^{\circ}$ C (Webster *et al.* 2003). For comparison, IPCC (2001*a*) reported a range for the global mean surface temperature rise by 2100 of $1.4-5.8 \,^{\circ}$ C, but did not provide likelihood estimates for this key finding although it did do so for others. The implications of the projected level of climate warming for the aquatic realm are unclear, but will vary with type of ecosystem. For example, small surface freshwater bodies will typically vary readily with climate, while in the open ocean, ambient temperature will lag that of the atmosphere by decades or more.

When the probability distribution functions for the mean global surface temperature and sea-level increases between 1990 and 2100 are compared between no-policy and applied-policy scenarios (designed to simulate strict regulations leading to stabilization of atmospheric CO_2 concentrations at about twice pre-industrial levels), a median of only 1.6 °C and 95% range of only 0.8–3.2 °C is forecast (Fig. 1.2). There is a 50% chance of warming exceeding 2.4 °C in the no-policy case and a 1-in-7 chance in the policy case. Implementing the policy therefore lowers the probability of large amounts of warming to a substantial degree.

To better appreciate the risks of the no-policy scenario, it is important to examine the latitudinal distribution of the projected warming. In common with other climate models, the computed temperature increases in polar regions are much greater than those in equatorial regions (no-policy case: Fig. 1.3). Polar regions contain vulnerable ecosystems with large carbon storage (e.g. Chapter 8), and the Greenland and Antarctic ice sheets with large water storage (e.g. Chapter 21). Release of some of this stored carbon and water is of significant concern. A 1-in-40 chance of warming by 8–12 °C or greater in polar regions for the nopolicy case (Fig. 1.3) is particularly worrisome. The policy scenario lowers the polar warming in the 1-in-40 calculation to 5–7 °C (Webster *et al.* 2003).

Similar significant reductions in the probability of large and risky amounts of sea-level rise due to the hypothetical policy are also evident (Fig. 1.2). Emissions reductions are predicted to lower the chance of exceeding an extreme climate outcome but not to eliminate the risk entirely, and analysis of the reduction in probability is an important policy consideration. Future climate assessments would better serve the policy process by including formal analysis of




Fig. 1.2. Probability density function (PDF) for the change in global mean: (a) surface temperature and (b) sea-level rise from 1990 to 2100 estimated as a best fit to 250 simulations using latin hypercube sampling from input PDFs for uncertain variables. The solid line shows the PDF resulting from no explicit emissions restrictions and the dashed line is the PDF under hypothetical emissions policy leading to steady levels of atmospheric CO₂ of about twice pre-industrial values (Webster *et al.* 2003). The IPCC (2001*a*, *b*, *c*) upper estimate is beyond the 95% confidence limit. Based on this distribution there is a 12% chance that the temperature change in 2100 would be less than the IPCC lower estimate.

uncertainty for key projections, with an explicit description of the methods used (Allen *et al.* 2001; Reilly *et al.* 2001).

Sea-level rise

Global mean sea level has risen by 10–25 cm over the last 100 years, and although there has been no detectable change in the rate of sea-level rise over the course of the century, it would appear that this has been significantly higher than the rate averaged over the last several thousand years (IPCC




Fig. 1.3. Zonal mean temperature change in surface warming by latitude band between 1990 and 2100 in the case assuming no explicit policy in Fig. 1.2. There is a 1-in-40 chance of being above or below the upper and lower curves and a 1-in-2 chance of being above or below the middle curve respectively (Webster *et al.* 2003).

2001*a*, *b*). It is likely that the rise in sea level has been largely due to the increase in global temperature over the last 100 years. The possible factors include thermal expansion of the oceans and melting of glaciers, ice caps and ice sheets. Changes in surface water and groundwater storage (related to changes in human activities and changes to drainage basins) may also have affected sea level.

Model simulations have predicted 2–7 cm of the reported sea-level rise over the last 100 years (10–25 cm) and attributed it to thermal expansion of the oceans, yet many of the world's mountain glaciers have retreated substantially over this time period and may have accounted for 2–5 cm of the observed rise. However, most of the non-oceanic water on Earth resides in great ice sheets, namely those of Antarctica and Greenland, and most of their volume lies on land above sea level. Loss of only a small fraction of this volume could have a significant effect on sea level (Warrick *et al.* 1996). In Antarctica, discharges of enormous icebergs and recent break-ups of Antarctic Peninsula ice shelves have focused public attention on the possibility of 'collapse' of this ice reservoir within the next century, with major potential impacts on sea level.

According to the IPCC (2001*a*, *b*) 'best estimate' climate change scenario, global sea level is projected to rise by 20 cm by the year 2050 (within a range of uncertainty of 7–39 cm) and by 49 cm by 2100 (within a range of uncertainty of 20–86 cm); more than half of this is attributed to oceanic thermal expansion.

Lakes, streams and wetlands

Inland aquatic ecosystems are expected to be greatly influenced by climate change through altered water temperatures, flow regimes and water levels (e.g. Kaczmarek *et al.* 1996; Öquist *et al.* 1996). Water-level declines may be severe in lakes and streams in dry evaporative drainages and in basins with small catchments, whilst the distribution of wetlands is likely to shift with changes in temperature and precipitation (IPCC 2001*a*; Chapters 2–10). Wetlands temporarily store runoff water, thereby reducing floodwater peaks and protecting downstream areas, consequently a reduction of wetland area due to climate change could severely hamper flood-control efforts in some regions (e.g. Kaczmarek *et al.* 1996; Öquist *et al.* 1996; Chapters 9 and 10).

Coastal systems and small islands

Coastal systems are economically and ecologically important. Climate warming, sea-level rise and increases in storms and storm-surge frequencies may result in erosion of shores and associated habitats, altered tidal ranges in rivers and bays, changes in sediment and nutrient transport, and increased coastal flooding (see Bijlsma *et al.* 1996). Some coastal ecosystems are particularly at risk, namely saltmarshes, mangroves, coral reefs and atolls, and river deltas (Chapters 11–13 and 16). Changes in these ecosystems are likely to negatively affect tourism, fisheries and biodiversity (IPCC 2001*b*). Many small island countries could lose a significant part of their land area with a sea-level rise of 0.5–1 m, the Maldives for instance having average altitudes of only 1–1.5 m (Bijlsma *et al.* 1996).

Oceans

Climate change could lead to altered ocean circulation (e.g. weakening of the Atlantic thermohaline circulation) and wave climates, and reductions in sea-ice cover (IPCC 2001*a*, *b*, *c*; ACIA [Arctic Climate Impact Assessment] 2004). As a result, nutrient availability, biological productivity, the structure and functions of marine ecosystems, and heat and carbon storage capacity may be affected with important feedbacks to the global climate system (see Ittekkot *et al.* 1996). These changes would have implications for coastal regions, fisheries, tourism and recreation, transport and offshore structures.

Most CO_2 released into the atmosphere as a result of burning fossil fuels will eventually be absorbed by the ocean. As the amount of CO_2 in the atmosphere rises, more of the gas reacts with seawater to produce bicarbonate and hydrogen ions that increase the acidity of the surface layer. Ocean pH was around 8.3 after the last ice age and 8.2 before CO_2 emissions took off in the industrial era. Ocean pH is now 8.1 and if atmospheric CO_2 exceeds 1900 ppm by the year 2300, pH at the ocean surface could fall to 7.4 (Caldeira & Wickett 2003). There is limited understanding of the effect increased acidity might have on marine biota, but coral reefs, calcareous plankton and other organisms, skeletons or shells of which contain calcium carbonate, may be substantially affected.

PEOPLE AND WATER

Perhaps the best-documented human interaction with aquatic ecosystems involves the use of fresh water (Engelman & LeRoy 1993; Engelman et al. 2000). Precious little of the world's water is salt-free, and only a small proportion of this fresh water is accessible to human beings. Water use has increased for many centuries, but in many major watersheds (the key geographic unit of interest for freshwater use), only in the past few centuries has the scale reached the point where natural variations in water supply have begun to collide with growing human use. This is most evident in western Asia and eastern and southern Africa, but to varying degrees such problems of availability are also present elsewhere. Humans already use more than half of the renewable fresh water that is readily accessible (Postel et al. 1996), and human population growth is currently the dominant factor in the increase of water withdrawals worldwide.

The world's urban population is currently growing at four times the rate of the rural population. Between 1990 and 2025, the number of people living in urban areas is projected to double to more than 5 billion out of a total 8 billion (Table 1.1). An estimated 90% of the increase will occur in developing countries and at the current pace, over 60 million people are added to urban populations each year, placing great strain on local governments to provide even the most basic services. Of all urban inhabitants in developing countries, 25–50% live in impoverished slums and squatter settlements, with little or no access to adequate water, sanitation or refuse collection (UN [United Nations] 1997).

Human water demand for agricultural irrigation (an increasing proportion of food production comes from irrigated cropland), industrial and household uses almost invariably takes precedence over environmental needs. In

 Table 1.1. World population by year and UN projection

 variant

Year	Low-variant projection	Medium-variant projection	High-variant projection
2010	6 843 645 000	6 906 558 000	6 967 407 000
2025	7 568 539 000	8 010 509 000	8 450 822 000
2050	7 791 545 000	9 191 287 000	10 756 366 000

Source: UNPD (2006).

all but the most remote and best-protected areas, there is constant pressure on water resources. By hydrological benchmarks of water stress ($c.1000-1700 \text{ m}^3$ per person per year) and scarcity (<1000 m³ per person per year) (Falkenmark & Widstrand 1992; Engelman & LeRoy 1993), the numbers of human beings living in these two high-risk categories is growing much faster than population growth generally. One-third of humanity lives in countries experiencing moderate to high water stress (WRI [World Resources Institute] 1998), and a billion people could face severe water shortages by 2025 (Potts 2000).

Apart from fisheries, which are considered below, several other interactions between human population and the environment are especially salient for aquatic ecosystems. In terms of food security, much is made of agricultural intensification as the antidote to forest loss. When more food can be produced on the same land, there is less need to convert relatively wild land to farmland. This can, of course, benefit wetlands, which historically have been lost to farmland more frequently than to settlement, although this ratio is changing as coasts and river valleys become more densely populated (Chapters 9–13).

Water is the greatest transportation network for chemical wastes of all types. Humanity now exceeds nature as a fixer and producer of nitrogen compounds (Smil 1997), and studies of rivers have shown that nitrogen loads are highly correlated with human population density along river banks (Cole *et al.* 1993). Intensive farming can also exacerbate soil erosion if farmers do not have the resources to manage their soil properly, and the world's rivers offload much of the resulting silt onto the continental shelves and beyond. Conversion of forest to farmland is another source of soil erosion, also closely correlated with population density and growth.

Introduction of alien species is an indirect result of human population growth and economic and technological development, because human beings uniquely face few barriers to their movement over the Earth's surface. From Nile perch to zebra mussels, the list of introduced aquatic species altering aquatic ecosystems is lengthening. In addition, marine life is at growing risk from a range of diseases, the spread of which is being hastened by global warming, global transport and pollution (Harvell *et al.* 1999, 2002). Well-documented cases include crab-eater seals in Antarctica infected with distemper by sled-dogs, sardines in Australia infected with herpes virus caught from imported frozen pilchards, and sea-fans in the Caribbean killed by a soil-borne fungus.

The further interaction between human populations and economics offers an additional obstacle to the conservation or restoration of aquatic ecosystems. When ecosystems are degraded, a common action of last resort is to attempt some sort of protection, where human use and access are restricted. But the process of setting land and/or fishing space aside becomes more difficult as population density increases. People may bid up the price of land to the point where conservation efforts become financially impossible (Cincotta & Engelman 2000). Land conservation organizations have increasingly focused on wetlands, coasts and other aquatic ecosystems, knowing they are in a race with time to buy key parcels of land or convince governments to protect them. Too rarely, however, do conservationists acknowledge that continued population growth limits their prospects for long-term success, simply because land will become too valuable to serve environmental rather than economic interests.

Market forces and global economics can also greatly influence the need for and development of port and shipping facilities. Historically, estuaries and saltmarshes have provided cheap sources of land for development. As commodities continue to be traded on global markets and ships become ever bigger, the demand for coastal land for development will increase and much of it will become degraded (Pinnegar *et al.* 2006).

While these interactions between human population and aquatic ecosystems paint a bleak picture, there are also reasons for hope. One of the most positive trends is that the human population is seen by most demographers as unlikely to double again. One way to explain this is that women all over the world are having fewer children than ever before, and seemingly want to have even fewer in the future. Average family size has shrunk from five children per woman in the early 1960s to a bit more than 2.5 children per woman today. In more than two-fifths of the world's countries, couples are only just replacing themselves in the population, or they are having fewer than the *c*.2.1 children needed for net replacement (UNPD [United Nations Population Division] 2002). Contrary to public impressions, this does not mean that these populations are now stable or declining. The large proportion of people of childbearing age in populations that up until recently were growing fairly rapidly guarantees that more births than deaths will occur for essentially an average human lifetime after replacement fertility is reached. Also, high levels of migration mean that most nations experiencing low fertility rates will continue to grow (International Organization for Migration 2000; UNPD 2002).

GLOBAL TRENDS: THE CASE OF GLOBAL MARINE FISHERIES

Given the means and opportunity, fishers like hunters before them ultimately deplete the resources that they target (e.g. Clovis of North America: Alroy 2001). Understanding this analogy is important, as it provides a framework for understanding the severe depletion of smaller and mid-size mammals in Africa ('bushmeat': Bowen-Jones 1998) and large fishes in the world's oceans. The recent marine fish biomass declines (e.g. Christensen *et al.* 2003 for the North Atlantic) are primarily attributable, not to scientific incompetence in monitoring (Malakoff 2002), nor shifts in distribution (Bigot 2002), nor regime shifts (Steele 1998), but to overfishing (Jackson *et al.* 2001).

When fishing starts in a new area, the large fishes go first, as they are relatively easier to catch than small fishes (e.g. with harpoons or lines) and tend to provide a better return on investment (Pauly *et al.* 2002). Large fish, with their low natural mortalities and relatively high age at first maturity (Pauly 1980; Froese & Binohlan 2000), cannot sustain substantial fishing pressure and rapidly decline (Denney *et al.* 2002), forcing the fishers either to move on to smaller fishes, and/or to other, previously unexploited areas. As larger fish tend to have higher trophic levels than smaller fish – indeed, the latter are usually the prey of the former – this process, now called 'fishing down marine food webs', leads to declining trends in the mean trophic level of fisheries landings (Pauly *et al.* 1998; though see Caddy *et al.* 1998).

The 'fishing down' process has occurred around the world (Table 1.2) and the broad pattern is that fishing targets a succession of species until the residual species mix ceases to support a fishing economy. This implies a

Country/area ^a	Years	Decline	Source and remarks
Iceland	1900–1999	1918–1999	Valtsson and Pauly (2003), based on comprehensive catch database of Valtsson (2001)
Celtic Sea	1945–1998	1946–2000	Pinnegar <i>et al.</i> (2002), based on trophic levels estimated from stable isotopes of nitrogen
Gulf of Thailand	1963-1982;	1965–1982;	Christensen (1998); Pauly and Chuenpagdee (2003)
	1963-1997	1965-1997	
Eastern Canada	1950–1997	1957–1997	Pauly <i>et al.</i> (2001), based on data submitted to FAO by the Canadian Department of Fisheries and Oceans
Western Canada	1873–1996	1910–1996	Pauly <i>et al.</i> (2001), based on comprehensive dataset assembled by S. Wallace
Cuban EEZ	1960-1995	1960-1995	Pauly et al. (1998) and Baisre (2000)
East Coast, USA	1950–2000	all	R. Chuenpagdee <i>et al.</i> unpublished data; emphasis on Chesapeake Bay
Chinese EEZ	1950-1998	1970-1998	Pang and Pauly (2001)
West Central	1950-2000;	1950-2000;	D. Pauly and M. L. Palomares unpublished data, based
Atlantic	1950–2000	1950-2000	on FAO data (Area 41), disaggregated into USA (North) and other countries (South)
World, tuna and billfishes	1950–2000	1950–2000	D. Pauly and M. L. Palomares unpublished data, based on FAO data (ISSCAAP [International Standard Statistical Classification of Aquatic Animals and Plants] group 36 only)
World, all fishes	1950–2000	1950–2000	Pauly and Watson (2003), based on spatially disaggregated data

Table 1.2. Occurrence of 'fishing down' using local/detailed data sets, following the original presentation of this phenomenon by Pauly et al. (1998), based on the global FAO catch dataset

^a EEZ, Exclusive Economic Zone.

transition from fisheries targeting large fish (e.g. northern cod) to those targeting smaller fishes (e.g. capelin) or invertebrates (e.g. northern shrimp and snow crab). In the north-east Atlantic, fish species that mature later, grow more slowly and have lower rates of potential population increase, have exhibited greater long-term declines in abundance than closely related species which are smaller, grow faster and are more fecund (Jennings et al. 1998, 1999; Dulvy et al. 2000). In sharks, rays and skates, large size combined with low fecundity makes them particularly vulnerable to overexploitation; many populations have been rapidly declining (Dulvy et al. 2000) and are likely to be rendered extinct in the coming decades (Dulvy et al. 2003). In some cases, depletion of keystone species may have important indirect effects on community and habitat structure (e.g. Dulvy et al. 2004).

Exports have become a major issue in fisheries, with marine products being amongst the most heavily traded commodities (Pauly et al. 2002). The general trend is for developed countries to increasingly to compensate for the shortfall of products from traditional fishing grounds in their exclusive economic zones (EEZs) by fishing in developing countries (Pauly et al. 2005). Given the debts that most developing countries have run up with respect to international lenders, this implies that marine resources, and their underlying ecosystems, suffer from increased pressure to make up shortfalls. Examples include the countries of West Africa, whose dependence on financial support from the European Union forces them to sign fisheries agreements providing access for European fishing fleets under terms that appear unfair to these countries (Kaczynski & Fluharty 2002). In Argentina, demersal

Major features	Representative groups
Large- to moderate-sized, predaceous, territorial reef fishes and rockfishes with late age at maturity, very low natural mortality rates and low recruitment rates versus adult stock size	Snappers, sea basses, emperors, rockfishes, sea breams
Large- to moderate-sized shelf dwelling, soft bottom predators susceptible to bottom trawling	Cods, flounders, soles, rockfishes, croakers, skates
Large- to moderate-sized schooling midwater fishes susceptible to midwater trawling	Hakes, rockfishes, armorheads, rougheyes
Large- to moderate-sized shelf dwelling, schooling, pelagic fishes	Bonitos, sierras, capelin, eulachon, salmon, sharks
Any species with exceptionally high monetary value	Bluefin tuna, red snappers, halibuts, medicinal fishes, aquarium fishes, groupers, salmon, red mullets, billfish

Table 1.3. Marine fish that are unlikely to survive, given continuation of present fisheries trends

Source: Adapted from Parrish (1995, 1998) and Pauly (2000).

resources which in the early 1980s were among the few that were both large and underexploited have now collapsed under the pressure of both national and international fleets, licensed for their ability to generate foreign exchange (Sánchez 2002).

Given present trends and pressures, present target species are not expected to survive and small non-palatable non-schooling species will be those that will fare best (Table 1.3). In extreme cases, finfish may be progressively replaced by consumer species such as jellyfish, which have few predators and may impede any finfish recovery (e.g. northern Benguela, Namibia: Lynam *et al.* 2006).

The fishing industry on its own is incapable of reversing the fishing down of food webs. There are other social forces, which can and will play an increasing role in the international debates on fisheries. Foremost is the community of non-governmental organizations devoted to maintaining or re-establishing 'healthy' marine ecosystems, and striving for ecosystem-based fisheries management. Public debate about fisheries was unheard of 20 years ago, and many conservation biologists feel they need to debunk those who deny the need for action (e.g. Lomborg 2001). Global catches are declining (Watson & Pauly 2001; Pauly *et al.* 2002) but catches can and usually do remain high when stocks collapse. Thus the cod off eastern Canada yielded good catches until the fishery had to be closed because there were literally no fish left (Myers *et al.* 1997).

Some suggest that aquaculture could help compensate for overfishing, and a quarter of human fish consumption

now derives from aquaculture. However, as currently practised, aquaculture also causes environmental damage, raising questions about how to meet food demands and preserve environmental quality (WRI 1998). Aquaculture in fact exists in two fundamentally different forms. One is devoted to the farming of bivalves (e.g. oysters, mussels) and/or freshwater fish (e.g. carp, tilapia) and relies mainly on plant matter to generate a net addition to the fish food supply available to consumers. This is based predominantly in developing countries (mainly in China, but also in countries such as the Philippines and Bangladesh), and supplies cheap animal protein where it is needed (New 2002). By contrast, the other form of aquaculture involves the farming of carnivorous fish such as salmon or sea bass, and increasingly, the fattening of wild-caught bluefin tuna. In nature, salmon, sea bass and bluefin tuna have high trophic levels, hence it is impossible to feed them only on vegetable matter. This implies that as this form of aquaculture increases, there will be fewer cheap fish (e.g. sardine, herring, mackerel and anchovies) available for humans to buy and eat. It is thus not adding to global fish supply, and instead increases the pressure on wild fish stocks (Naylor et al. 2000).

This second type of aquaculture predominates, and it has led to massive imports by developed countries of meal from fishes caught and ground up in developing countries, exacerbating fishing pressures in these regions. Coastal pollution and diseases emanating from the uneaten food and facces of these marine feed-lot operations are also seen as a major constraint to the development of the industry (New 2002).

Fisheries need to be reinvented as providers of a healthy complement to grain-based diets. Nor can they remain subject to a free-for-all among distant-water fleets; they can however become a regular source of income for communities whose members act within natural constraints (Pitcher 2001). Such reinvented fisheries will be smaller in size, and they will hopefully rely more on fish biomass being exported from marine protected areas closed to fishing.

Humans consume around 86 million tonnes of fish per year, nearly 15.7 kg per person, which is more than twice the 1950 level. Inhabitants of the economically less developed countries (including China) will likely increase their total consumption of food fish to the 2025 time horizon by as much as 34 million tonnes, whereas consumption will remain relatively static elsewhere (Delgado *et al.* 2003). Given current population trends and the global market for fishery products, the question remains as to whether fishing pressure can ever be reduced and aquatic ecosystems restored.

GLOBAL TRENDS IN AQUATIC BIODIVERSITY AND EXTINCTION

With the exception of seven countries, all nations of the world are now signatories to the UN Convention on Biological Diversity, introduced in 1992 at the Rio 'Earth Summit'. The Convention committed nations 'to achieve by 2010 a significant reduction of the current rate of biodiversity loss at the global, regional and national level as a contribution to poverty alleviation and to the benefit of all life on earth'. However, both marine and freshwater ecosystems continue to be degraded and species lost all around the world.

Freshwater systems occupy only 0.8% of Earth's surface (McAllister *et al.* 1997), but they are rich in species and vital as habitat. Perhaps 12% of all animal species live in fresh water (Abramovitz 1996). Due to their limited area, freshwater ecosystems contain only about 2.4% of all Earth's plant and animal species (Reaka-Kudla 1997); however, on a hectare-for-hectare basis, they are richer in species than the more extensive terrestrial and marine ecosystems (Table 1.4).

The alteration and damming of river systems for industrial and domestic use, irrigation and hydroelectric power have fragmented more than half of the world's large river systems. Some 83% of their total annual flow is affected

Ecosystem	Per cent of Earth's total habitat	Per cent of known species ^a	Relative species richness ^b
Freshwater	0.8	2.4	3
Terrestrial	28.4	77.5	2.7
Marine	70.8	14.7	0.2

Table 1.4. Species richness by ecosystem

^{*a*} Species do not sum to 100% because 5.3% of known symbiotic species are excluded.

^bRelative species richness is the ratio of the per cent of known species and the per cent of area occupied by the ecosystem.

Source: McAllister et al. (1997).

(52% moderately, 31% severely), with Europe's river flow being the most regulated and Australasia's the least (WWF [World Wide Fund for Nature] 2006). While many factors can simultaneously contribute to freshwater fish extinctions, habitat alteration and introduction of non-native species have been the major causes of species losses (Harrison & Stiassny 1999). Habitat alteration has contributed to 71% of extinctions, non-native species (which can compete with or feed on native species) to 54%, overfishing to 29% and pollution to 26% (Harrison & Stiassny 1999).

Dramatic declines in amphibian populations, including population crashes and mass localized extinction, have been noted since the 1980s from locations all over the world, and amphibian declines are thus perceived as one of the most critical threats to global biodiversity. A number of causes are believed to be involved, including habitat destruction and modification, overexploitation, pollution, pesticide use, introduced species, climate change, increased ultraviolet-B radiation (UV-B) and diseases. However, many of the causes of amphibian declines are still poorly understood, and amphibian declines are currently a topic of much ongoing research.

In an attempt to provide a quantitative overview of the long-term changing health of the planet, in 1998 the WWF produced a 'Living Planet Report', which has since been updated annually. The stated aim of this report was to answer the question: how fast is nature disappearing from the Earth? It introduced several useful concepts, including the Living Planet Index (LPI), the aggregate of the Forest Ecosystems Index, Freshwater Ecosystems Index (FEI) and Marine Ecosystems Index (MEI) (Fig. 1.4).




Fig. 1.4. Freshwater and marine Living Planet Indices (FEI [Freshwater Ecosystems Index] and MEI [Marine Ecosystems Index] respectively) from 1970 to 2003 (WWF 2006).

The FEI is based on the population trends of 344 freshwater vertebrate species, of which 387 are temperate and 51 tropical. These species include every mammal, bird, reptile, amphibian and fish for which time series population data are available, and they now encompass 11 mammals, 153 birds, 17 reptiles, 69 amphibians and 94 fishes (WWF 2006). The FEI points to freshwater species having on average declined by about 28% since 1970 (Fig. 1.4), the decline being more marked in temperate than tropical systems. Studies around the world have pointed to declines in freshwater animal populations, notable among which are amphibians (e.g. Wyman 1990; Wake 1991; Pounds & Crump 1994; Houlahan et al. 2000). However, large declines have also occurred in many freshwater fish (e.g. Moyle & Leidy 1992; Stiassny 1996; WCMC [World Conservation Monitoring Centre] 1998) and bivalve populations (e.g. Bogan 1993; Ricciardi et al. 1998). Some 81 fish species are recorded to have become extinct during the past century, and a further 11 are extinct in the wild but remain as captive populations (Fig. 1.5).

A major proportion of known extinctions (notably, 50 species of cichlid) have resulted from the ecological effects of the deliberate introduction of the Nile perch *Lates niloticus* into Lake Victoria in the mid twentieth century (WCMC 1998). In North America alone, some 123 freshwater animal species have been recorded as extinct since 1900, and hundreds of additional species of fishes, molluscs, crayfishes and amphibians are imperilled (Ricciardi & Rasmussen 1999). In total, 734 species of fish are classified as threatened, of which 84% are freshwater species (IUCN [World Conservation Union] 1996).




Fig. 1.5. Cumulative sum of known fish species extinctions by decade (WCMC 1998).

The MEI is based on population trends of 274 marine species, including 48 marine mammals, 112 seabirds, 7 reptiles (turtles) and 107 fishes. The MEI has declined by about 27% since 1970, a slightly slower decline than is evident in the FEI (WWF 2006) but nonetheless substantial (Fig. 1.4). Relatively stable trends are evident in the Pacific and in the Arctic/Atlantic Oceans, in comparison with dramatic declines in the Indian/Southeast Asian and Southern Oceans (Table 1.5). However, overall increases in the populations of seabirds and some mammal species in the Atlantic and Pacific Oceans since 1970 mask declines in many fish species, especially those of economic importance such as cod and tuna, as well as turtles and other species that are caught as by-catch.

There is a public misconception that there is little likelihood of a marine species ever becoming extinct (Roberts & Hawkins 1999). However, this has happened in the past and is set to continue and possibly increase in frequency in the future, as habitats are changed and exploitation continues. Notable among large vulnerable species are the sea mammals, populations of which were decimated by whaling and hunting up until the second half of the twentieth century. Steller's sea cow Hydrodamalis gigas was hunted to extinction in 1767 and the Caribbean monk seal Monachus tropicalis in 1952 (Day 1990), while species such as the Mediterranean monk seal Monachus monachus and Gulf porpoise Phocoena sinus remain critically endangered. Several species of fish and molluscs have also been lost, and many remain critically endangered (Roberts & Hawkins 1999).

Among fisheries globally, 75% have been judged to be fully fished, overfished or depleted (FAO [Food and Agriculture Organization of the United Nations] 2004). In addition to the effects of exploitation, the biodiversity of many coral reef and coastal marine species may also be influenced by habitat loss (Friedlander & Parrish 1998; Wilson et al. 2006). Of 58 marine fish population extirpations, exploitation was the primary cause of decline in 40 cases, the remainder being attributable to habitat loss, although the effects of habitat loss and exploitation on abundance are not always readily separable (Dulvy et al. 2003). Given ongoing increases in fishing effort and globally declining catches (Watson & Pauly 2001; Pauly 2002), and in the absence of a global network of relatively large marine protected areas (Russ & Zeller 2003), it is expected that large marine fishes, both pelagic and demersal, could be fished to extinction in the next few decades (e.g. see Sadovy & Cheung 2003; Dulvy et al. 2003). The fisheries targeting these species will obviously go bankrupt in the process, although they may continue to last for a while if propped up by sufficient subsidies.

HOW MANY ECOSYSTEMS?

The number of distinct ecosystems recognized differs markedly and this is particularly so for the aquatic realm. It has often been a matter of personal taste as to how many are deemed to exist and on what criteria they are separated. Ecosystems blend into one another and sharp boundaries are a convenience for cartographers rather than a reality of nature. There are many potential ways of dividing up the world's aquatic systems including by depth, geography, climatic or hydrological conditions, and primary productivity, or the nature of the biological communities. Which of these criteria are the most useful will depend on the purpose of the classification. At one end of the spectrum authors have recognized only two aquatic ecosystems, namely marine and freshwater systems, whilst other texts describe as many as 57 distinct systems in the marine environment alone (Longhurst et al. 1995).

Marine systems

The oceans cover about 71% of the Earth's surface and reach depths of more than 10000 m. They extend from regions where precipitation exceeds evaporation to where the opposite is true. Perhaps the most general possible division consists of five marine 'biomes' separated on the basis of depth, temperature and nutrient status. These include: warm continental shelves, cold continental shelves,

warm open oceans, cold open oceans and upwelling waters (Bradshaw 1977). More detailed treatments generally take their lead from work in terrestrial biomes and divide aquatic systems into distinct types of biological community such as coral reefs, saltmarshes or mangroves. Thus, based on other work (Whittaker 1975; Friday & Ingram 1985; Barnes & Hughes 1988; Archibold 1995), Chapman and Reiss (1998) suggested 10 distinct marine systems and five freshwater systems. Other schemes reflect less the types of assemblage present and more the geographic location or climatological/ hydrological conditions.

An approach adopted by IUCN, several UN agencies and the US National Oceanographic and Atmospheric Administration (NOAA), is that of Large Marine Ecosystems (LMEs). The 64 recognized LMEs are relatively large geographical units, characterized by distinct bathymetry, hydrography, productivity and trophically dependent populations (Sherman & Alexander 1986; Sherman et al. 2005). Another possible scheme for dividing up the world's aquatic systems is that used by the FAO. Twenty-seven major fishing areas are recognized for statistical purposes, and consist of eight major inland areas, each covering the waters of one of the eight continents, and 19 major marine areas covering the waters of the Atlantic, Indian, Pacific and Southern Oceans, and adjacent seas. The latter are generally divided along lines of latitude and longitude and range in size from 3 million km² (Mediterranean and Black Sea) to 49 million km² (Pacific, Eastern Central) (FAO 1994).

A further possible scheme for dividing the world's marine systems is that of the UNEP Regional Seas Programme initiated in 1974 as an approach to the control of marine pollution and resource management. The Programme has action plans for 13 regions and involves the participation of more than 140 coastal states and territories. Thus far, however, there are large parts of the world's oceans that are not covered by this scheme, and the open ocean is a major source for concern (Pauly *et al.* 2003; Chapters 20–22).

Freshwater environment

In nearly all existing schemes to describe the world's aquatic systems, there are fewer types of freshwater than marine systems. At one extreme, the scheme of Baily (1998) included no freshwater system *per se*, but instead all rivers, streams, lakes and ponds were included as part of the particular ecoregion in which they lay, as defined by climatic

conditions and altitude. A similar approach was adopted by Bradshaw (1977). The most common distinction is that between running waters (e.g. streams and rivers) and still waters (e.g. lakes and ponds) (e.g. Chapman & Reiss 1998). Three other types of freshwater systems may be highlighted, namely cool temperate bogs, tropical freshwater swamp forests (e.g. those of the Amazon basin) and temperate freshwater swamp forests (e.g. the Everglades) (Chapman & Reiss 1998); these have also been recognized in the *Ecosystems of the World* book series (Goodall 1977–2000), albeit viewed as terrestrial rather than freshwater systems (Gore 1983; Lugo *et al.* 1990).

Discussion

All of the schemes mentioned have both advantages and disadvantages. For example, the FAO system is comprehensive in that all the world's oceans and freshwater systems are included, but the divisions used are not always biologically appropriate. Thus, many of the species found in the north-east Atlantic (e.g. cod and haddock) are also found in the north-west Atlantic and Arctic Sea, whilst inland boundaries such as those between Asia, the former USSR and Europe are not clearly defined in terms of their catchments, where certain rivers (e.g. Danube and Amur) start in one continent but flow into another.

For this book, the most appropriate scheme was one based on a categorization of ecosystems largely by particular biological communities (e.g. coral reefs, rivers), although this system has problems of its own. For example, is an apparently coherent community of organisms sufficiently similar throughout the world to be grouped? For instance, are coral reefs on the Great Barrier Reef adequately similar to those in the Caribbean to be labelled together? For the purposes of this book, the answer to this question is 'yes', since coral reefs around the world generally face similar types of threat including global warming, eutrophication and overfishing, and might thus be expected to respond in a broadly similar manner (Chapter 16). Likewise, rivers across the world face threats such as canalization, channelization, eutrophication, pollution and excessive water extraction for irrigation (Chapter 2). Scientists in general are more accustomed to writing global reviews relating to a single type of community, rather than reviewing issues relating to all types of community at a particular geographic location or region.

Starting with the list of 15 aquatic systems and their definitions in Chapman and Reiss (1998), a number of

factors were important in determining which of these groupings were to be retained, which further subdivided and which joined into a single category. Chapman and Reiss (1998) recognized many types of shallow community underlain by soft substrata (e.g. marine mudflats, sandy beaches, temperate saltmarshes, mangroves), but did not include seagrass beds, which are ecologically distinctive and comparatively well studied (e.g. McRoy & Helfferich 1977). In addition, 'marine mudflats' and 'sandy beaches' tend to encompass only shallow or intertidal communities and do not seem to include soft-sediment systems at greater depths on continental shelves, many of which support substantial trawl fisheries (Jennings & Kaiser 1998). A better solution for describing the myriad softbottom categories was to include 'sandy beaches', 'saltmarsh/mudflats' and 'subtidal soft-bottom' communities.

For marine systems underlain by hard substrata, Chapman and Reiss (1998) recognized 'marine rocky shore' 'continental shelf benthos' and 'coral reef'. Both the terms 'marine rocky shore' and 'continental shelf benthos' are somewhat vague, the former being used to encompass rocky intertidal communities, and the latter to include subtidal communities such as kelp beds which may reach 50 m depth (Chapman & Reiss 1998). Therefore, clearer terminology for these widely researched ecosystems was required ('rocky intertidal': Chapter 14).

With respect to freshwater systems, the large inland seas such as the North American and African Great Lakes, and the Aral and Caspian Seas, are here recognized as being distinct from smaller bodies of standing water, since many of these have their own characteristic biotas and are affected by somewhat different threats, where factors such as eutrophication may represent a more important challenge. One type of aquatic system which has been neglected from nearly all global syntheses, but which has been the subject of considerable scientific interest (e.g. Wilkens *et al.* 2000), is that of subterranean water courses. Finally, given their importance to the origins and modern sustenance of human societies, and in spite of overlaps with other ecosystems, it was felt useful to include flood plains as a separate ecosystem.

GOALS OF THIS BOOK

While much emphasis is being put on climate change, its anthropogenic drivers and its likely impacts on ecosystems and society, the direct impacts of human population growth and industrialization have long had, are having, and will continue to have substantial consequences for the extent, structure and functioning of aquatic ecosystems. It is broadly known that many salty and freshwater ecosystems on land have already been particularly affected by human agency, and those of the open oceans, especially the deep sea, little altered. However there have been no comprehensive attempts to compare expert opinions of the status and likely trends across all aquatic ecosystems at a global level. This book originated in a series of reviews published in the journal Environmental Conservation which were further progressed by expert groups at the 5th International Conference on Environmental Future at the Eidgenössische Technische Hochschule (ETH, Zürich, Switzerland, March 2003) and brought to fruition thereafter. The book will systematically explore current and likely future states of aquatic ecosystems, spanning from running waters and small water bodies to the massive marine systems around the poles, and in the open oceans. The overall questions to which answers were sought included the following: What are the key threats to the status of each of the aquatic ecosystems? To what extent might each of these ecosystems change between now and the 2025 time horizon? How do the patterns of threat and likely change vary among these ecosystems and what drives these variations? What information does this overview vield for the conservation science and management of the world's aquatic ecosystems? Because the 21 ecosystems selected for treatment are classifiable into seven categories (flowing waters, still waters, freshwater wetlands, coastal wetlands, rocky shores, soft shores, and vast marine systems), the opportunity was provided not only to make comparisons between particular ecosystems but also to compare among the different ecosystem types. The book accordingly has seven principal parts, each with its own preamble, and in the final Chapter 23 aims to synthesize the information on trends and prognoses, and draw broad conclusions about the threats and trajectories of the world's aquatic ecosystems, their overall structures and ecological functions. The IPCC process has helped to show how building scientific consensus and effectively addressing audiences outside that of science is a long iterative process. The global environmental assessment of aquatic ecosystems that this book represents is one scientifically based step towards sharing sound information and ensuring this ultimately contributes to the development of appropriate international policies and actions.

References

- Abbott, I. A. & Hollenberg, G. J. (1976) Marine Algae of California. Stanford, CA, USA: Stanford University Press.
- Åberg, E. (1992) Tree colonization of three mires in southern Sweden. In: *Peatland Ecosystems and Man: An Impact Assessment*, ed. O. M. Bragg, P. D. Hulme, H. A. P. Ingram & R. A. Robertson, pp. 268–270. Dundee, UK: Department of Biological Sciences, University of Dundee.
- Abernethy, R. K. & Turner, R. E. (1987) US forested wetlands: 1940–1980. *BioScience* 37: 721–727.
- Abramovitz, J. N. (1996) Imperiled Waters, Impoverished Future: The Decline of Freshwater Ecosystems, Worldwatch Paper No. 128. Washington, DC, USA: Worldwatch Institute.
- Acharya, G. (2000) Approaches to valuing the hidden hydrological services of wetland ecosystems. *Ecological Economics* 35: 63–74.
- ACIA (Arctic Climate Impact Assessment) (2004) *Impacts of a Warming Arctic*. Cambridge, UK: Cambridge University Press.
- Ackefors, H. & Enell, M. (1994) The release of nutrients and organic matter from aquaculture systems in Nordic countries. *Journal of Applied Ichthyology* 10: 225–241.
- Adam, P. (1978) Geographical variation in British saltmarsh vegetation. *Journal of Ecology* 66: 339–366.
- Adam, P. (1990) *Saltmarsh Ecology*. Cambridge, UK: Cambridge University Press.
- Adam, P. (2000) Morecambe Bay saltmarshes: 25 years of change. In: *British Saltmarshes*, ed. B. R. Sherwood,
 B. G. Gardiner & T. Harris, pp. 81–107. Tresaith, UK: Forrest Text.
- Adam, P. (2002) Saltmarshes in a time of change. Environmental Conservation 29: 39–61.
- Adams, N. L. (2001) UV radiation evokes negative phototaxis and covering behavior in the sea urchin *Strongylocentrotus droebachiensis. Marine Ecology Progress Series* 213: 87–95.
- Adams, W. M. (2001) Green Development: Environment and Sustainability in the Third World, 2nd edn. London, UK: Routledge.

Adey, W. H. & Steneck, R. S. (2001) Thermogeography over time creates biogeographic regions: a temperature/space/ time-integrated model and an abundance-weighted test for benthic marine algae. *Journal of Phycology* 37: 677–698.

Adey, W. H. (1964) The genus *Phymatolithon* in the Gulf of Maine. *Hydrobiologia* 24: 377–420.

- Adis, J. & Junk, W. J. (2002) Terrestrial invertebrates inhabiting lowland river floodplains of Central Amazonia and Central Europe: a review. *Freshmater Biology* 47: 711–731.
- Ærtebjerg, G., Andersen, J. H. & Hansen, O. S. (2003) Nutrients and eutrophication in Danish marine waters: a challenge for science and management. Unpublished report. Roskilde, Denmark: National Environmental Research Institute.
- Aerts, R., Wallén, B. & Malmer, N. (1992) Growth-limiting nutrients in *Sphagnum*-dominated bogs subject to low and high atmospheric nitrogen supply. *Journal of Ecology* 80: 131–140.
- Agawin, N. S. R. & Duarte, C. M. (2002) Evidence of direct particle trapping by a tropical seagrass meadow. *Estuaries* 25: 1205–1209.
- Agnew, D. J. & Nicol, S. (1996) Marine disturbances: commercial fishing. In: *Foundations for Research West of the Antarctic Peninsula*, Antarctic Research Series vol. 70, ed.
 R. M. Ross, E. E. Hofmann & L. B. Quetin, pp. 417–35. Washington, DC, USA: American Geophysical Union.
- Aioi, K. & Nakaoka, M. (2003) The seagrasses of Japan. In: World Atlas of Seagrasses, ed. E. P. Green & F. T. Short, pp. 201–210. Berkeley, CA, USA: University of California Press.
- Airoldi, L. (1998) Roles of disturbance, sediment stress, and substratum retention on spatial dominance in algal turf. *Ecology* 79: 2759–2770.
- Aladin, N. V. & Potts, W. T. W. (1992) Changes in the Aral Sea ecosystems during the period 1960–1990. *Hydrobiologia* 237(2): 67–79.
- Aladin, N. V., Filippov, A. A., Plotnikov, I. S., Orlova, M. I. & Williams, W. D. (1998) Changes in the structure and function of biological communities in the northern part of

the Aral Sea (Small Aral Sea), 1984–1994. International Journal of Salt Lake Research 7: 301–343.

- Albers, P. H. (2002) Sources, fate, and effects of PAHs in shallow water environments: a review with special reference to small watercraft. *Journal of Coastal Research* Special Issue 37: 143–150.
- Alcamo, J., Döll, P., Kaspar, F. & Siebert, S. (1997) Global Change and Global Scenarios of Water Use and Availability: An Application of WaterGAP1.0. Kassel, Germany: Center for Environmental Systems Research (CESR), University of Kassel.
- Alcocer, J. & Escobar, E. (1990) The drying up of the Mexican Plateau axalapazcos. *Salinet* 4: 44–46.
- Alcocer, J. & Williams, W. D. (1996) Historical and recent changes in Lake Texcoco, a saline lake in Mexico. *International Journal of Salt Lake Research* 5: 45–61.
- Alcocer, J., Escobar, E. & Lugo, A. (2000) Water use (and abuse) and its effects on the crater-lakes of Valle de Santiago, Mexico. *Lakes and Reservoirs: Research and Management* 5: 145–149.
- Aldhous, P. (2003) The world's forgotten crisis. *Nature* **422**: 251–253.
- Alewell, C., Manderscheid, B., Meesenburg, H. & Bittersohl, J. (2000) Is acidification still an ecological threat? *Nature* 407: 856–857.
- Al Hamed, M. I. (1966) Limnological studies on the inland waters of Iraq. *Bulletin of the Iraq Natural History Museum* 3: 1–22.
- Alin, S., Cohen, A., Bills, R., et al. (1999) Effects of landscaped disturbance on animal communities in Lake Tanganyika, East Africa. *Conservation Biology* 13: 1017–1033.
- Allan, J. D. & Flecker, A. S. (1993) Biodiversity conservation in running waters. *BioScience* 43: 32–43.
- Allan, J. D., Abell, R., Hogan, Z., et al. (2005) Overfishing of inland waters. BioScience 55: 1041–1051.
- Allen, J. R., Slinn, D. J., Shammon, T. M., Hartnoll, R. G. & Hawkins, S. J. (1998) Evidence for eutrophication of the Irish Sea over four decades. *Limnology and Oceanography* 43: 1970–1974.
- Allen, J. R. L. (2000) Morphodynamics of Holocene salt marshes: a review sketch from the Atlantic and Southern North Sea coasts of Europe. *Quaternary Science Reviews* 19: 1155–1231.
- Allen, M., Raper, S. & Mitchell, J. (2001) Uncertainty in the IPCC's *Third Assessment Report. Science* 293: 430–433.
- Allendorf, F. W. & Lundquist, L. L. (2003) Introduction: population biology, evolution, and control of invasive species. *Conservation Biology* 17: 24–30.

- Allison, G. W., Lubchenco, J. & Carr, M. H. (1998) Marine reserves are necessary but not sufficient for marine conservation. *Ecological Applications* 8: 79–92.
- Alm, J., Sculman, L., Walden, J., et al. (1999) Carbon balance of a boreal bog during a year with an exceptionally dry summer. *Ecology* 80: 161–174.
- Almendinger, J. E. & Leete, J. H. (1998a) Peat characteristics and groundwater chemistry of calcareous fens in the Minnesota River Basin, USA. *Biogeochemistry* 43: 17–41.
- Almendinger, J. E. & Leete, J. H. (1998b) Regional and local hydrogeology of calcareous fens in the Minnesota River Basin, USA. *Wetlands* 18: 184–202.
- Alongi, D. M. (1998) Coastal Ecosystem Processes. Boca Raton, FL, USA: CRC Press.
- Alongi, D. M. (2002) Present state and future of the world's mangrove forests. *Environmental Conservation* 29: 331–349.
- Alroy, J. (2001) A multispecies overkill simulation of the end-Pleistocene megafaunal mass extinction. *Science* 292: 1893–1896.
- AMAP (1998) *AMAP Assessment Report: Arctic Pollution Issues.* Oslo, Norway: Arctic Monitoring and Assessment Programme.
- Ambroggi, R. P. (1977) Underground reservoirs to control the water cycle. *Scientific American* 236: 21–28.
- Ambrose, R. F. (2000) Wetland mitigation in the United States: assessing the success of mitigation policies. *Wetlands* 19: 1–27.
- Amon, J. P., Thompson, C. A., Carpenter, Q. J. & Miner, J. (2002) Temperate zone fens of the glaciated midwestern USA. *Wetlands* 22: 301–317.
- Amoros, C. & Petts, G. E., eds. (1993) Hydrosystèmes Fluviaux. Paris, France: Masson.
- Amsberry, L., Baker, M. A., Ewanchuk, P. J. & Bertness, M. D. (2000) Clonal integration and the expansion of *Phragmites australis*. *Ecological Applications* 10: 1110–1118.
- Amthor, J. S. (1995) Terrestrial higher-plant response to increasing atmospheric (CO₂) in relation to the global carbon cycle. *Global Change Biology* 1: 243–274.
- Andersen, U. V. (1995a) Resistance of Danish coastal vegetation types to human trampling. *Biological Conservation* 71: 223–230.
- Andersen, U. V. (1995b) Invasive aliens: a threat to the Danish coastal vegetation? In: *Directions in European Coastal Management*, ed. M. G. Healy & P. J. Doody, pp. 335–344. Cardigan, UK: Samara Publishing Ltd.

Anderson, D. M., Glibert, P. M. & Burkholder, J. M. (2002a) Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25: 704–726.

Anderson, D. M., Overpeck, J. T. & Gupta, A. K. (2002b) Increase in the Asian Southwest monsoon during the past four centuries. *Science* 297: 596–599.

Anderson, J. A. R. (1983) The tropical peatswamps of western Malesia. In: *Mires: Swamp, Bog, Fen and Moor, Regional Studies, Ecosystems of the World 4B*, ed.
A. J. P. Gore, pp. 181–199. Amsterdam, the Netherlands: Elsevier.

Anderson, O. F. & Clark, M. R. (2003) Analysis of bycatch in the fishery for orange roughy, *Hoplostethus* atlanticus, on the South Tasman Rise. *Marine Freshwater Research* 54: 643–652.

Anderson, R. J., Carick, P., Levitt, G. J. & Share, A. (1997) Holdfasts of adult kelp *Ecklonia maxima* provide refuges from grazing for recruitment of juvenile kelps. *Marine Ecology Progress Series* 159: 265–273.

Anderson, S. & Moss, B. (1993) How wetland habitats are perceived by children: consequences for children's education and wetland conservation. *International Journal* for Science Education 15: 473–485.

Anderson, T. W. (1994) Role of macroalgal structure in the distribution and abundance of a temperate reef fish. *Marine Ecology Progress Series* 113: 279–290.

Andersson, F. & Olsson, B., eds. (1985) Lake Gårdsjön: an acid forest lake and its catchment. *Ecological Bulletins* (*Stockholm*) Special Issue 37.

Andersson, G., Berggren, H., Cronberg, G. & Gelin, C. (1978) Effects of planktivorous and benthivorous fish on organisms and water chemistry in eutrophic lakes. *Hydrobiologia* 59: 9–15.

Andréasson-Gren, I.-M. & Groth, K.-H. (1995) Economic evaluation of Danube floodplains. Unpublished report. Gland, Switzerland: WWF International.

Andrew, N. L. (1993) Spatial heterogeneity, sea urchin grazing, and habitat structure on reefs in temperate Australia. *Ecology* 74: 292–302.

Andrew, N. L. (1994) Survival of kelp adjacent to areas grazed by sea urchins in New South Wales, Australia. *Australian Journal of Ecology* 19: 466–472.

Andrew, N. L. & Choat, J. H. (1982) The influence of predation and non-specific adults on the abundance of juvenile *Evechinus chloroticus* (Echinoidea: Echinodermata). *Oecologia* 54: 80–87. Andrew, N. L. & Jones, G. P. (1990) Patch formation by herbivorous fish in a temperate kelp forest. *Oecologia* 85: 57–68.

Andrew, N. L. & O'Neill, A. L. (2000) Large-scale patterns in habitat structure on subtidal rocky reefs in New South Wales. *Marine and Freshwater Research* 51: 255–263.

Andrew, N. L. & Underwood, A. J. (1993) Density-dependent foraging in the sea urchin *Centrostephanus rodgersii* on shallow subtidal reefs in New South Wales, Australia. *Marine Ecology Progress Series* **99**: 89–98.

Angel, A. & Ojeda, F. P. (2001) Structure and trophic organization of subtidal fish assemblages on the northern Chilean coast: the effect of habitat complexity. *Marine Ecology Progress Series* 217: 81–91.

Angus, S. & Elliott, M. M. (1992) Erosion in Scottish machair with particular reference to the Outer Hebrides. In: Coastal Dunes: Geomorphology, Ecology and Management for Conservation, ed. R. W. G. Carter, T. G. F. Curtis & M. J. Sheehy-Skeffington, pp. 93–112. Rotterdam, the Netherlands: A. A. Balkema.

Annadotter, H., Cronberg, G., Aagren, R., et al. (1999) Multiple techniques for lake restoration. *Hydrobiologia* 395/396: 77–85.

Anon. (1999) Wetlands: A Source of Life, Conclusions of the 2nd International Conference on Wetlands and Development, 10–14 November 1998, Dakar, Senegal. Wageningen, the Netherlands: Wetlands International; Gland, Switzerland: IUCN; and Dakar, Senegal: Ministry of Environment and Nature Protection.

Anon. (2003) Indien: Ende aller Sorgen. *Der Spiegel* **40**: 116–117 (in German).

Anon. (2004) A remedy for financial turbulence? *The Economist* **371**: 72.

Antila, C. K., Daehler, C. C., Rank, N. E. & Strong, D. R. (1998) Greater male fitness of a rare invader (*Spartina alterniflora* Poaceae) threatens a common native (*Spartina foliosa*) with hybridization. *American Journal of Botany* 85: 1597–1601.

Antila, C. K., King, R. A., Ferris, C., Ayres, D. R. & Strong, D. R. (2000) Reciprocal hybrid formation of

Spartina in San Francisco Bay. Molecular Ecology 9: 765–770.

Applegate, V. & Moffett, J. (1955) The sea lamprey. Scientific American 192: 36–41.

Archibold, O. W. (1995) Ecology of World Vegetation. London, UK: Chapman & Hall.

Armentano, T. V. & Menges, E. S. (1986) Patterns of change in the carbon balance of organic soil-wet lands of the temperate zone. *Journal of Ecology* 74: 755–774. Arntz, W. E., Valdivia, E. & Zeballos, J. (1988) Impact of El Niño 1982–83 on the commercially exploited invertebrates (Mariscos) of the Peruvian shore. *Meeresforschung* 32: 3–22.

Aronson, R. B. & Precht, W. F. (2001) Evolutionary paleoecology of Caribbean coral reefs. In: *Evolutionary Paleoecology: The Ecological Context of Macroevolutionary Change*, ed. W. D. Allmon & D. J. Bottjer, pp. 171–233. New York, USA: Columbia University Press.

Arp, W. J. & Drake, B. G. (1991) Increased photosynthetic capacity of *Scirpus olneyi* after 4 years of exposure to elevated CO₂. *Plant, Cell and Environment* 14: 1003–1006.

Arp, W. J., Drake, B. G., Pockman, W. T., Curtis, P. S. & Whigham, D. F. (1993) Interactions between C₃ and C₄ salt marsh plant species during four years of exposure to elevated atmospheric CO₂. *Vegetatio* 104/105: 133–143.

Arrigo, K. R., Worthen, D., Schnell, A. & Lizotte, M. P. (1998) Primary production in Southern Ocean waters. *Journal of Geophysical Research – Oceans* 103: 15587–15600.

Arrow, K., Bolin, B., Costanza, R., et al. (1995) Economic growth, carrying capacity and the environment. Science 268: 520–521.

Arthington, A. H. & Pusey, P. J. (2003) Flow restoration and protection in Australian rivers. *River Research and Applications* 19: 377–395.

Aselmann, I. & Crutzen, P. J. (1989) Global distribution of natural wetlands and rice paddies, their net primary productivity, seasonality and possible methane emissions. *Journal of Atmospheric Chemistry* 8: 307–358.

Ashjian, C. J., Smith, S. L. & Lane, P. V. Z. (1995) The northeast water polynya during summer 1992: distribution and aspects of secondary production of copepods. *Journal of Geophysical Research – Oceans* 100: 4371–4388.

Ashton, E. D., Macintosh, J. & Hogarth. P. (2003) A baseline study of the diversity and community ecology of crab and molluscan macrofauna in the Sematan mangrove forest, Sarawak, Malaysia. *Journal of Tropical Ecology* 19: 127–142.

Ashworth, W. (1987) *The Late, Great Lakes: An Environmental History.* Detroit, MI, USA: Wayne State University Press.

Asiaweek (2000) The bottom line. *Asiameek* 24(45). Available online at www.asiaweek.com/asiaweek/magazine/2000/ 1117/bottomline.html

Aubrey, D. G. (1993) Coastal erosion's influencing factors include development, dams, wells, and climate change. *Oceanus* 36: 5–9. Auster, P. J. & Langton, R. W. (1999) The effects of fishing on fish habitat. *American Fisheries Society Symposium* 22: 150–187.

Awosika, L. F., Ibe, A. C. & Ibe, C. E. (1993) Anthropogenic activities affecting sediment load balance along the west African coastline. In: *Coastlines of Western Africa*, ed. L. F. Awosika, A. C. Ibe & P. Shroader, pp. 26–39. New York, USA: American Society of Civil Engineers.

Ayres, D. R., Garcia-Rossi, D., Davis, H. G. & Strong, D. R. (1999) Extent and degree of hybridization between exotic (*Spartina alterniflora*) and native (*S. foliosa*) cordgrass (Poaceae) in California, USA determined by random amplified polymorphic DNA (RAPDs). *Molecular Ecology* 8: 1179–1186.

Ayres, D. R., Strong, D. R. & Baye, P. (2003) *Spartina foliosa* (Poaceae): a common species on the road to rarity? *Madroño* **50**: 209–213.

Baan, P. J. A. & van Buuren, J. T. (2003) Testing of Indicators for the Marine and Coastal Environment in Europe. Copenhagen, Denmark: European Environment Agency.

Babcock, R.C, Kelly, S., Shears, N. T., Walker, J. W. & Willis, T. J. (1999) Changes in community structure in temperate marine reserves. *Marine Ecology Progress Series* 189: 125–134.

Bach, S. S., Borum, J., Fortes, M. D. & Duarte, C. M. (1998) Species composition and plant performance of mixed seagrass beds along a siltation gradient at Cape Bolinao, the Philippines. *Marine Ecology Progress Series* 174: 247–256.

Baden, S. P. & Pihl, L. (1990) Effects of oxygen depletion on the ecology, blood physiology and fishery of the Norway lobster, *Nephrops norvegicus. Marine Ecology Progress Series* 67: 151–155.

Baer, K. E. & Pringle C. M. (2000) Special problems of urban river conservation: the encroaching megapolis. In: *Global Perspectives in River Conservation*, ed. P. J. Boon,
B. R. Davies & G. E. Petts, pp. 385–402. Chichester, UK: John Wiley.

Baeteman, C. (1994) Subsidence in coastal lowlands due to groundwater withdrawal: the geological approach. *Journal of Coastal Research* Special Issue 37: 61–75.

Bailey, R. G. (1998) Ecoregions: The Ecosystem Geography of the Oceans and Continents. New York, USA: Springer-Verlag.

Baird, R. C., Thompson, N. P., Hopkins, T. L. & Weiss, W. R. (1975) Chlorinated hydrocarbons in mesopelagic fishes of the eastern Gulf of Mexico. *Bulletin of Marine Science* 25: 473–481. Baisre, J. (2000) Chronicles of Cuban Marine Fisheries (1935–1995): Trend Analysis and Fisheries Potential, FAO Fisheries Technical Paper 394. Rome, Italy: Food and Agriculture Organization of the United Nations.

Baker, A. C., Starger, C. J., McClanahan, T. R. & Glynn, P. W. (2004) Corals' adaptive response to climate change. *Nature* 430: 741.

Bakker, J. B. (1989) Nature Management by Grazing and Cutting: On the Ecological Significance of Grazing and Cutting – Regimes Applied to Restore Former Species-Rich Grassland Communities in the Netherlands. Dordrecht, the Netherlands: Kluwer.

Bakker, S. A., Van den Berg, N. J. & Speleers, B. P. (1994) Vegetation transitions of floating wetlands in a complex of turbaries between 1937 and 1989 as determined from aerial photographs with GIS. *Vegetatio* 114: 161–167.

Baldwin, N., Saalfeld, R., Ross, M. & Buettner, H. (1979) Commercial Fish Production in the Great Lakes 1867–1977, Technical Report No. 3. Ann Arbor, MI, USA: Great Lakes Fishery Commission.

Balgos, M. C., Ricci, N., Walker, L., et al. (2005) Draft: Compilation of summaries of national and regional ocean policies. The Nippon Foundation Research Task Force on National Ocean Policies, Gerard J. Mangone Center for Marine Policy, University of Delaware [www document]. URL http://www.globaloceans.org/tops2005/pdf/ OceanPolicySummaries. pdf#search=%22%22national% 20ocean%20policy%22%22

Ball, M. C., Cochrane, M. J. & Rawson, H. M. (1997) Growth and water use of the mangroves *Rhizophora apiculata* and *R. stylosa* in response to salinity and humidity under ambient and elevated concentrations of atmospheric CO₂. *Plant Cell and Environment* 20: 1158–1166.

Balls, P. W., Macdonald, A., Pugh, K. & Edwards, A. C. (1995) Long-term nutrient enrichment of an estuarine system: Ythan, Scotland (1958–1993). *Environmental Pollution* **90**: 311–321.

Baltuck, M., Dickey, J., Dixon, T. & Harrison, G. A. (1996) New approaches raise questions about future sea-level change. *Eos* 77: 385–388.

Bandeira, S. O. & Gell, F. (2003) The seagrasses of Mozambique and Southeastern Africa. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 105–112.
Berkeley, CA, USA: University of California Press.

Barber, K. E. (1981) *Peat Stratigraphy and Climate Change*. Rotterdam, the Netherlands: A.A. Balkema.

Barber, R. T., Vijayakumar, A. & Cross, F. A. (1972) Mercury concentrations in recent and ninety-year-old benthopelagic fish. *Science* 178: 636–639. Barbier, E. B. (1997) The economic determinants of land degradation in developing countries. *Philosophical Transactions of the Royal Society of London B* 352: 891–899.

Barbier, E. B. (2000) The economic linkages between rural poverty and land degradation: some evidence from Africa. *Agriculture, Ecosystems and Environment* 82: 355–370.

Barbier, E. B. & Thompson, J. R. (1998) The value of water: floodplain versus large-scale irrigation benefits in Northern Nigeria. *Ambio* 27: 434–440.

Barbraud, C. & Weimeskirch, H. (2001) Emperor penguins and climate change. *Nature* 411: 183–186.

Barkai, A. & McQuaid, C. (1988) Predator-prey role reversal in a marine benthic ecosystem. *Science* 242: 4875.

Barlow, M. & Clarke, T. (2002) *Blue Gold: The Battle Against Corporate Theft of the World's Water*. Toronto, Ontario Canada: Stoddart.

Barnes, R. S. & Hughes, R. N. (1988) An Introduction to Marine Ecology, 2nd edn. Oxford, UK: Blackwell, Scientific Publications.

Barnett, P. R. O. (1971) Some changes in intertidal sand communities due to thermal pollution. *Proceedings of the Royal Society of London B* 177: 353–364.

Barnett, P. R. O. & Hardy, B. L. S. (1969) The effects of temperature on the benthos near the Hunterston generating station, Scotland. *Chesapeake Science* 16: 255–256.

Barnett, T. P., Pierce, D. W. & Schnur, R. (2001) Detection of anthropogenic climate change in the world's oceans. *Science* 292: 270–274.

Barraclough, S. & Finger-Stich, A. (1996) Social Implications of Commercial Shrimp Farming in Asia. Geneva, Switzerland: United Nations Research Institute for Social Development.

Barrales, H. L. & Lobban, C. (1975) The comparative ecology of *Macrocystis pyrifera* with emphasis on the forests of Chubut, Argentina. *Ecology* 63: 657–677.

Bart, D. & Hartman, J. M. (2003) The role of large rhizome dispersal and low salinity windows in the establishment of common reed, *Phragmites australis*, in salt marshes: new links to human activities. *Estuaries* 26: 433–436.

Bartley, D. M. (2000) International mechanisms for the conservation and sustainable use of wetlands. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 331–345. Leiden, the Netherlands: Backhuvs.

Bascand, L. D. (1970) The roles of *Spartina* species in New Zealand. *Proceedings of the New Zealand Ecological Society* 17: 33–40. Baumgartner, T. R., Soutar, A. & Ferreira-Bartrina, V. (1992) Reconstruction of the history of Pacific sardine and northern anchovy populations over the past two millennia from sediments of the Santa Barbara basin, California. *CalCOFI Report* 33: 24–40.

Bax, N. J. (2000) Stock Assessment Report 2000: Orange Roughy (Hoplostethus atlanticus). Hobart, Tasmania, Australia: South East Fishery Stock Assessment Group.

Baye, P. (1990) Ecological history of an artificial foredune ridge on a northeastern barrier spit. In: *Proceedings of the Symposium on Coastal Sand Dunes*, ed. R. G. D. Davidson-Arnott, pp. 389–403. Ottawa, Ontario, Canada: National Research Council Canada.

Bayley, P.B. (1995) Understanding large river-floodplain ecosystems. *BioScience* **45**: 143–158.

Bayliss, B., Brenman, K., Elliot, I., et al. (1997) Vulnerability Assessment of Predicted Climate Change and Sea Level Rise in the Alligator Rivers Region, Northern Territory Australia, Supervising Scientist Report No. 123. Canberra, ACT, Australia: Supervising Scientist.

Bayly, I. A. E. (1967) The general biological classification of aquatic environments with special reference to those in Australia. In: *Australian Inland Waters and their Fauna: Eleven Studies*, ed. A. H. Weatherley, pp. 78–104. Canberra, ACT, Australia: Australian National University Press.

Beamish, R. J. (1993) Climate and exceptional fish production off the west coast of North America. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 2270–2291.

Beardall, J., Beer, S. & Raven, J. A. (1998) Biodiversity of marine plants in an era of climate change: some predictions based on physiological performance. *Botanica Marina* 41: 113–123.

Bebianno, M. J. & Langston, W. J. (1991) Metallothionein induction in *Mytilus edulis* exposed to cadmium. *Marine Biology* 108: 91–96.

Bebianno, M. J. & Langston, W. J. (1992) Cadmium induction of metallothionein synthesis in *Mytilus* galloprovincialis. Comparative Biochemistry and Physiology C 103: 79–85.

Bebianno, M. J. & Langston, W. J. (1995) Induction of metallothionein synthesis in the gill and kidney of *Littorina littorea* exposed to cadmium. *Journal of the Marine Biological Association of the United Kingdom* 75: 173–186.

Bebianno, M. J., Nott, J. A. & Langston, W. J. (1993) Cadmium metabolism in the clam *Ruditapes decussata*: the role of metallothioneins. *Aquatic Toxicology* 27: 315–333.

Beck, B. F. & Herring J. G. (2002) Geotechnical and environmental applications of karst geology and hydrology. In: Proceedings of the 8th Interdisciplinary Conference on Sinkholes and the Engineering and Environmental Impacts of Karst, ed. B. F. Beck & J. G. Herring, pp. 83–88. Rotterdam, the Netherlands: A. A. Balkema.

Beck, M. W., Heck Jr, K. L., Able, K. W., *et al.* (2003) The role of nearshore ecosystems as fish and shellfish nurseries. *Issues in Ecology* **11**: 1–12.

Beck, W. (1998) Warmer and wetter 6000 years ago? *Science* **279**: 1003–1004.

Becker, A. & Grünewald, U. (2003) Flood risk in Central Europe. *Science* **300**: 1099.

Becker, B. (1997) Sustainability assessment: a review of values, concepts and methodological approaches. *The World Bank – Issues in Agriculture* 10: 1–63.

Bedford, B. L. (1996) The need to define hydrologic equivalence at the landscape scale for freshwater wetland mitigation. *Ecological Applications* 6: 57–68.

Bedford, B. L. (1999) Cumulative effects on wetland landscapes: links to wetland restoration in the United States and southern Canada. *Wetlands* 19: 775–788.

Bedford, B. L. & Godwin, K. S. (2003) Fens of the United States: distribution, characteristics, and scientific connection versus legal isolation. *Wetlands* 23: 608–629.

Bedford, B. L. & Preston, E. M. (1988) Developing the scientific basis for assessing cumulative effects of wetland loss and degradation on landscape functions: status, perspectives, and prospects. *Environmental Management* 12: 751–771.

Bedford, B. L., Leopold, D. J. & Gibbs, J. (2001) Wetland ecosystems. In: *Encyclopedia of Biodiversity*, vol. 5, ed. S. A. Levin, pp. 781–804. New York, USA: Academic Press.

Beeftink, W. G. (1966) Vegetation and habitat of the salt marshes and beach plains in the south-western part of the Netherlands. *Wentia* 15: 83–108.

Beer, S. & Koch, E. (1996) Photosynthesis of marine macroalgae and seagrass in globally changing CO₂ environments. *Marine Ecology Progress Series* 141: 199–204.

Beeton, A. (1969) Changes in the environment and biota of the Great Lakes. In: *Eutrophication: Causes, Consequences, Correctives Symposium Proceedings*, pp. 150–187. Washington, DC, USA: National Academy of Sciences.

Beeton, A. (1984) The world's Great Lakes. *Journal of Great Lakes Research* 10: 106–13.

Beeton, A. (2002) Large freshwater lakes: present state, trends, and future. *Environmental Conservation* 29: 21–38.

Beeton, A. (1965) Eutrophication of the St Lawrence Great Lakes. *Limnology and Oceanography* **10**: 240–254.

- Beeton, A. & Edmondson, W. (1972) The eutrophication problem. *Journal of the Fisheries Research Board of Canada* 29: 673–682.
- Beeton, A. & Hageman, J. (1994) Impact of Dreissena polymorpha on the zooplankton community of western Lake Erie. Verhandlungen Internationale Vereinigung für Limnologie 25: 2349 (abstract).

Beeton, A., Sellinger, C. & Reid, D. (1999) An introduction to the Laurentian Great Lakes ecosystem. In: *Great Lakes Fisheries Policy and Management*, ed. W. W. Taylor & C. P. Ferreri, pp. 3–54. East Lansing, MI, USA: Michigan State University Press.

Bellamy, D. J. & Rieley, J. (1967) Ecological statistics of a miniature bog. *Oikos* 18: 33–40.

Bellan, G. & Bellan-Santini, D. (2001) A review of littoral tourism, sport and leisure activities: consequences on marine flora and fauna. *Aquatic Conservation: Marine and Freshwater Ecosystems* 11: 325–333.

Beltman, B., Van den Broek, T. & Bloemen, S. (1995) Restoration of acidified rich-fen ecosystems in the Vechtplassen area: successes and failures. In: *Restoration of Temperate Wetlands*, ed. B. D. Wheeler, S. C. Shaw, W. J. Fojt & R. A. Robertson, pp. 273–286. Chichester, UK: John Wiley.

Beltman, B., Van den Broek, T., Van Maanen, K. & Vaneveld, K. (1996) Measures to develop a rich-fen wetland landscape with a full range of successional stages. *Ecological Engineering* 7: 299–314.

Belyea, L. R. & Clymo, R. S. (2001) Feedback control of the rate of peat formation. *Proceedings of the Royal Society* of London B 268: 1315–1321.

Belyea, L. R. & Malmer, N. (2004) Carbon sequestration in peatland: patterns and mechanisms of response to climate change. *Global Change Biology* **10**: 1043–1052.

Benayas, J. M. R., Colomer, M. G. S. & Levassor, C. (1999) Effects of area, environmental status and environmental variation on species richness per unit area in Mediterranean wetlands. *Journal of Vegetation Science* 10: 275–80.

Benda, L. E., Poff, N. L., Tague, C., *et al.* (2002) How to avoid train wrecks when using science in environmental problem solving. *BioScience* 52: 1127–1136.

Benke, A. C., Chaubey, I., Ward, M. & Dunn, E. L. (2000) Flood pulse dynamics of an unregulated river floodplain in the Southeastern US coastal plain. *Ecology* 81: 2730–2741.

Bennett, E. M., Carpenter, S. R., Peterson, G. D., et al. (2003) Why global scenarios need ecology. Frontiers in Ecology and Environment 1: 322–329.

Benson, L., Kashgarian, M., Rye, R., et al. (2002) Holocene multidecadal and multicentennial droughts affecting

Northern California and Nevada. *Quaternary Science Reviews* 21: 659–682.

Berdnikov, S. V., Selyutin, V. V., Vasilchenko, V. V. & Caddy, J. F. (1999) Trophodynamic model of the Black and Azov Sea pelagic ecosystem: consequences of the comb jelly, *Mnemiopsis leidyi*, invasion. *Fisheries Research* 42: 261–289.

Berendonk, T. U. & Bonsall, M. B. (2002) The phantom midge and a comparison of metapopulation structures. *Ecology* 83: 116–128.

Berendse, F., van Breemen, N., Rydin, H., et al. (2001) Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production in *Sphagnum* bogs. *Global Change Biology* 7: 591–598.

Bergkamp, G., Pirot, J.-Y. & Hostetter, S., eds. (2000) Integrated Management of Wetlands and Water Resources.
Proceedings of the 2nd International Wetlands Conference, Dakar, 1998. Gland, Switzerland: Ramsar.

Berglund, B. E., Malmer, N. & Persson, T. (1991) Landscape-ecological aspects of long-term changes in the Ystad area. *Ecological Bulletins* 41: 405–424.

Bergstad, O. A. (1990) Distribution, population structure, growth and reproduction of the roundnose grenadier (*Coryphaenoides rupestris*) (Pisces: Macrouridae) in the deep waters of the Skagerrak. *Marine Biology* 107: 25–39.

Berkes, F. (1987) The common property resource problem and the fisheries of Barbados and Jamaica. *Environmental Management* 11: 225–235.

Berkes, F., Hughes, T. P., Steneck, R. S., *et al.* (2006) Globalization, roving bandits, and marine resources. *Science* **311**: 1557–1558.

Berkes, F., Mahon, R., McConney, P., Pollnac, R. & Pomeroy, R. (2001) Managing Small-Scale Fisheries: Alternative Directions and Methods. Ottawa, Ontario, Canada: International Development Research Centre.

Berkmüller, K., Mukherjee, S. & Mishra. B. (1990) Grazing and cutting pressures on Ranthambhore National Park, Rajasthan, India. *Environmental Conservation* 17: 135–140.

Bernstein, B. B. & Jung, N. (1980) Selective pressures and coevolution in a kelp canopy community in Southern California. *Ecological Monographs* 49: 335–355.

Bernstein, B. B. & Welsford, R. W. (1982) An assessment of feasibility of using high-calcium quicklime as an experimental tool for research into kelp bed/sea urchin ecosystems in Nova Scotia. *Canadian Technical Report of Fisheries and Aquatic Sciences* **968**: 1–51.

Bertness, M. D. (1984) Habitat and community modification by an introduced herbivorous snail. *Ecology* **65**: 370–381. Bertness, M. D. & Ellison, A. M. (1987) Determinants of pattern in a New England marsh plant community. *Ecological Monographs* 57: 129–147.

Bertness, M. D., Ewanchuk, P. & Silliman, B. R. (2002) Anthropogenic modification of New England salt marsh landscapes. *Proceedings of the National Academy of Sciences* of the USA 99: 1395–1398.

Bethoux, J. P., Morin, P., Chaumery, C., et al. (1998) Nutrients in the Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to environmental change. *Marine Chemistry* 63: 155–169.

Bett, B. J. (2001) UK Atlantic Margin Environmental Survey: introduction and overview of bathyal benthic ecology. *Continental Shelf Research* 21: 917–956.

Betts, R. A. (2000) Offset of the potential carbon sink from boreal forestation by decreases in surface albedo. *Nature* 408: 187–190.

Beutel, M. W., Horne, A. J., Roth, J. C. & Barratt, N. J. (2001) Limnological effects of anthropogenic desiccation of a large, saline lake, Walker Lake, Nevada. *Hydrobiologia* 466(1–3): 91–105.

Beyth, M. (2002) The Red Sea and the Mediterranean Dead Sea canals project [www document]. URL http://www. mfa.gov.il/mfa/go.asp?MFAH0mb90.

Biebl, R. & McRoy, C. P. (1971) Plasmatic resistance and rate of respiration and photosynthesis of *Zostera marina* at different salinities and temperatures. *Marine Biology* 8: 48–56.

Bigot, J. (2002) Les stocks ne diminuent pas, ils se déplacent. L'Industrie Nouvelle 2851: 10.

Bijlsma, L., Ehler, C. N., Klein, R. J. L., et al. (1996) Coastal zones and small islands. In: Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change, ed. Intergovernmental Panel on Climate Change, pp. 289–324. Cambridge, UK: Cambridge University Press.

Billett, D. S. M., Bett, B. J., Rice, A. L., et al. (2001) Longterm change in the megabenthos of the Porcupine Abyssal Plain (NE Atlantic). Progress in Oceanography 50: 325–348.

Bird, E. C. F. (1985) Coastline Changes: A Global Review. Chichester, UK: John Wiley.

Bird, E. C. F. (1990) Artificial beach nourishment on the shores of Port Phillip Bay, Australia. *Journal of Coastal Research* Special Issue 6: 55–68.

Bird, E. C. F. (1993) Submerging Coasts: The Effects of a Rising Sea Level on Coastal Environments. Chichester, UK: John Wiley.

Bird, E. C. F. & Cullen, P. W. (1996) Recreational uses and problems of Port Phillip Bay, Australia. In: *Recreational Uses of Coastal Areas*, ed. P. Fabbri, pp. 39–51. Dordrecht, the Netherlands: Kluwer. Birkeland, C. (1982) Terrestrial runoff as a cause of outbreaks of *Acanthaster planci* (Echinodermata: Asteroidea). *Marine Biology* 69: 175–185.

Birkeland, C., Dayton, P. K. & Engstrom, N. A. (1982) A stable system of predation on a holothurian by four asteroids and their top predator. *Australian Museum Memoirs* 16: 175–189.

Biswas, A. K. (1998) Deafness to global water crisis: causes and risks. *Ambio* 27: 492–493.

Bjorgo, E., Johannessen, O. M. & Miles, M. W. (1997) Analysis of merged SSMR–SSMI time series of Arctic and Antarctic sea ice parameters 1978–1995. *Geophysical Research Letters* 24: 413–416.

Bjorndal, K. A. (1980) Nutrition and grazing behaviour of the green turtle *Chelonia mydas*. *Marine Biology* 56: 147–154.

Blasco, F. (1983) Mangroves of Gambia and Senegal. Toulouse, France: University of Toulouse and UN Sahelian Office.

Blasco, F. (1984) Taxonomic considerations of the mangrove species. In: *The Mangrove Ecosystem*, ed. S. C. Snedaker & J. G. Snedaker, pp. 81–90. Paris, France: UNESCO.

Blasco, F., Aizpuru, M. & Gers, C. (2001) Depletion of the mangroves of Continental Asia. Wetlands Ecology and Management 9: 255–266.

Blasco, F., Gauquelin, T., Rasolofoharinoro, M., Aizpuri, M. & Caldairou, V. (1998) Recent advances in mangrove studies using remote sensing data. *Marine and Freshmater Research* 49: 287–296.

Blossey, B. (2003) A framework for evaluating potential ecological effects of implementing biological control of *Phragmites australis. Estuaries* 26: 607–617.

Blowers, A. & Smith, S. (2003) Introducing environmental issues: the environment of an estuary. In: Understanding Environmental Issues, ed. S. Hinchliffe, A. Blowers & J. Freeland, pp. 3–48. Chichester, UK: John Wiley.

Bobbink, R., Ashmore, M., Braun, S., Flückiger, W. & Van den Wyngaert, I. J. J. (2003) Empirical nitrogen critical loads for natural and semi-natural ecosystems: 2002 update. In: *Empirical Critical Loads for Nitrogen*, Environmental Documentation No. 124, ed. B. Acherman & R. Bobbink, pp. 43–170. Berne, Switzerland: Swiss Agency for the Environment, Forests and Landscape.

Bodkin, J. L. (1988) Effects of kelp forest removal on associated fish assemblages in central California. *Journal of Experimental Marine Biology and Ecology* 117: 227–238.

Boehlert, G. W. (1986) Productivity and population maintenance of seamount resources and future research directions. In: *Environment and Resources of Seamounts* in the North Pacific, NOAA Technical Report,

ed. R. Uchida, S. Hayasi & G. W. Boehlert, pp. 95–101. Seattle, WA, USA: National Marine Fisheries Service.

Boehlert, G. W., Yoklavich, M. M. & Chelton, D. B. (1989) Time series growth in the genus *Sebastes* from the northeast Pacific Ocean. *Fishery Bulletin* 87: 791–806.

Boesch, D. F., Brinsfield, R. B. & Magnien, R. E. (2001) Chesapeake Bay eutrophication: scientific understanding of ecosystem restoration and challenges for agriculture. *Journal of Environmental Quality* **30**: 303–320.

Boesch, D. F., Josselyn, M. N., Mehta, A. J., et al. (1994) Scientific assessment of coastal wetland loss, restoration, and management in Louisiana. *Journal of Coastal Research* Special Issue 20: 1–103.

Bogan, A. E. (1993) Freshwater bivalve extinctions (Mollusca, Unionoida): a search for causes. *American Zoologist* 33: 599–609.

Bohrer, B., Heidenreich, H., Schimmele, M. & Schultze, M. (1998) Numerical prognosis for salinity profiles of future lakes in the opencast mine of Merseburg-Ost. *International Journal of Salt Lake Research* 7: 235–260.

Boje, R. & Tomczak, M. (1978) Upwelling Ecosystems. New York, USA: Springer-Verlag.

Bokn, T., Berge, J. A., Green, N. & Rygg, B. (1990) Invasion of the planktonic algae *Chrysochromulina polylepis* along south Norway in May–June 1988: acute effects on biological communities along the coast. In: *Eutrophication and Algal Blooms in North Sea Coastal Zones, the Baltic and Adjacent Areas: Prediction and Assessment of Preventative Actions,* Water Pollution Research Report No. 12, ed. C. Lancelot, G. Billen & H. Barth, pp. 183–193. Brussels, Belgium: Commission of the European Communities.

Bokuniewicz, H. C. (1990) Tailoring local responses to rising sea level: a suggestion for Long Island. *Shore and Beach* 58(3): 22–25.

Bolen, E. G. & Guthery, F. S. (1982) Playas, irrigation, and wildlife in West Texas. North American Wildlife Conference 47: 528–541.

Bologna, P. A. X. & Steneck, R. S. (1993) Kelp beds as habitat for American lobster *Homarus americanus*. *Marine Ecology Progress Series* 100: 127–134.

Bolton, J. J. (1996) Patterns of species diversity and endemism in comparable temperate brown algal floras. *Hydrobiologia* 326/327: 173–178.

Bolton, J. J. & Anderson, R. J. (1987) Temperature tolerances of two southern African *Ecklonia* species (Alariaceae: Laminariales) and of hybrids between them. *Marine Biology* 96: 293–297. Bonacci, O., Zdenko, T. & Trninic, D. (1992) Effects of dams and reservoirs on the hydrological characteristics of the Lower Drava River. *Regulated Rivers: Research and Management* 7: 349–357.

Bondesan, M., Castiglioni, G. B., Elmi, C., et al. (1995) Coastal areas at risk from storm surges and sea-level rise in northeastern Italy. *Journal of Coastal Research* 11: 1354–1379.

Bongaarts, J., Mauldin, W. P. & Phillips, J. F. (1990). The demographic impact of family planning programs. *Studies in Family Planning* 21: 299–310.

Bonner, W. N. (1982) *Seals and Man.* Seattle, WA, USA: University of Washington Press.

Bonsdorff, E., Blomqvist, E. M., Mattila, J. & Norkko, A. (1997) Coastal eutrophication: causes, consequences and perspectives in the archipelago areas of the northern Baltic Sea. *Estuarine, Coastal and Shelf Science* 44: 63–72.

Boorman, L. A. (1989) The grazing of British sand dune vegetation. *Proceedings of the Royal Society of Edinburgh B* 96: 75–88.

Boorman, L. A. (2003) Saltmarsh Review: An Overview of Coastal Saltmarshes, their Dynamic and Sensitivity Characteristics for Conservation and Management, JNCC Report No. 334. Peterborough, UK: Joint Nature Conservation Council.

Bootsma, H. & Hecky, R. (1993) Conservation of the African Great Lakes: a limnological perspective. *Conservation Biology* 7: 644–656.

Bootsma, H. & Hecky, R. (2003) A comparative introduction to the biology and limnology of the African Great Lakes. *Journal of Great Lakes Research* 29(2): 3–19.

Borum, J. (1996) Shallow waters and land/sea boundaries. In: *Eutrophication in Coastal Marine Ecosystems*, ed.
K. Richardson & B. B. Jorgensen, pp. 179–203.
Washington, DC, USA: American Geophysical Union.

Boström, C., Baden, S. & Krause-Jensen, D. (2003) The seagrasses of Scandinavia and the Baltic Sea. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 35–54. Berkeley, CA, USA: University of California Press.

Botch, M. S. & Masing, V. V. (1983) Mire ecosystems in the USSR. In: *Mires: Swamp, Bog, Fen and Moor, Regional Studies, Ecosystems of the World 4B*, ed. A. J. P. Gore, pp. 95–152. Amsterdam, the Netherlands: Elsevier.

Bothner, M. H., Buchholtz ten Brink, M. & Manheim, F. T. (1998) Metal concentrations in surface sediments of Boston Harbor: changes with time. *Marine Environmental Research* 45: 127–155. Bothner, M. H., Takada, H., Knight, I. T., et al. (1994) Sewage contamination in sediments beneath a deep-ocean dump site off New York. *Marine Environmental Research* 38: 43–59.

Botsford, L. W., Castilla, J. C. & Peterson, C. H. (1997) The management of fisheries and marine ecosystems. *Science* 277: 509–515.

Botton, M. L., Loveland, R. E. & Jacobsen, T. R. (1988) Beach erosion and geochemical factors: influence on spawning success of horseshoe crabs (*Limulus polyphemus*) in Delaware Bay. *Marine Biology* 99: 325–332.

Botton, M. L., Loveland, R. E. & Jacobsen, T. R. (1994) Site selection by migratory shorebirds in Delaware Bay, and its relationship to beach characteristics and abundance of Horseshoe Crab (*Limulus polyphemus*) eggs. *Auk* 111: 605–616.

Boudouresque, C. F., Charbonel, E., Meinesz, A., *et al.* (2000) A monitoring network based on the seagrass *Posidonia oceanica* in the northwestern Mediterranean Sea. *Biologia Marina Mediterranea* 7: 328–331.

Boulé, M. E. (1994) An early history of wetland ecology. In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 57–74. Amsterdam, the Netherlands: Elsevier.

Boulton A. J., Sheldon, F., Thoms, M. C. & Stanley, E. H. (2000) Problems and constraints in managing rivers with variable flow regimes. In: *Global Perspectives on River Conservation: Science, Policy and Practice*, ed. P. J. Boon, B. R. Davies & G. E. Petts, pp. 411–426. Chichester, UK: John Wiley.

Bourke, R. H. & Garrett, R. P. (1987) Sea ice thickness distribution in the Arctic Ocean. *Cold Regions Science and Technology* 13: 259–280.

Bourke, R. H. & McLaren, A. S. (1992) Contour mapping of Arctic Basin ice draft and roughness parameters. *Journal of Geophysical Research* 97: 17715–17728.

Bourque, B. J. (1995) Diversity and Complexity in Prehistoric Maritime Societies: A Gulf of Maine Perspective. New York, USA: Plenum Press.

Bowen-Jones, E. (1998) A Review of the Commercial Bushmeat Trade, with Emphasis on Central/West Africa and the Great Apes. Cambridge, UK: Ape Alliance/Flora International.

Bowes, G. (1993) Facing the inevitable: plants and increasing atmospheric CO₂. *Annual Review of Plant Physiology and Plant Molecular Biology* 44: 309–332.

Bowles, I. (2001) Environmental meltdown in the White house. *Boston Globe* 22 April.

Bowman, M. J. (1995) The Ramsar Convention comes of age. Netherlands International Law Review 42: 1–52. Bowman, M. (2002) The Ramsar Convention on Wetlands: has it made a difference? In: *Yearbook of International Cooperation on Environment and Development 2002–03*, eds.
O. S. Stokke & B. Øystein, pp. 61–68. London, UK: Earthscan Publications.

Bowman, R. S. & Lewis, J. R. (1977) Annual fluctuations in the recruitment of *Patella vulgata* L. *Journal of the Marine Biological Association of the United Kingdom* 57: 793–815.

Boyd, P. W. & Newton, P. P. (1999) Does pelagic community structure determine downward particulate organic carbon flux in different oceanic provinces? *Deep-Sea Research I* 46: 63–91.

Boyer, K. E. & Zedler, J. B. (1996) Damage to cordgrass by scale insects in a constructed salt marsh: effects of nitrogen additions. *Estuaries* 19: 1–12.

Boyer, K. E. & Zedler, J. B. (1998) Effects of nitrogen additions on the vertical structure of a constructed cordgrass marsh. *Ecological Applications* 8: 692–705.

Boyer, K. E. & Zedler, J. B. (1999) Nitrogen addition could shift species composition in a restored California salt marsh. *Restoration Ecology* 7: 74–85.

Boyer, K. E., Callaway, J. C. & Zedler, J. B. (2000) Evaluating the progress of restored cordgrass (*Spartina foliosa*) marshes: belowground biomass and tissue N. *Estuaries* 23: 711–721.

Boynton, W. R., Hagy, J. D., Murray, L., Stokes, C. & Kemp, W. M. (1996) A comparative analysis of eutrophication patterns in a temperate coastal lagoon. *Estuaries* 19: 408–421.

Bradbury, R. & Seymour, R. (1997) Waiting for COTS. Proceedings of the 8th International Coral Reef Symposium, Panama, June 24–29 1996, vol. 1, ed. H. A. Lessios & I. E. Macintyre, pp. 1357–1362. Balboa, Panama: Smithsonian Tropical Research Institute.

Bradbury, R. H., Hammond, L. S., Moran, P. J. & Reichelt, R. E. (1985) Coral reef communities and the crown-of-thorns starfish: evidence for qualitatively stable cycles. *Journal of Theoretical Biology* 113: 69–80.

Bradley, D. C. & Ormerod, S. J. (2001) Community persistence among stream invertebrates tracks the North Atlantic Oscillation. *Journal of Animal Ecology* 70: 987–996.

Bradley, D. C. & Ormerod, S. J. (2002) Long-term effects of catchment liming on invertebrates in upland streams. *Freshmater Biology* 47: 161–171.

Bradley, P. M. & Morris, J. T. (1991) The influence of salinity on the kinetics of NH₄+ uptake in *Spartina alterniflora*. *Oecologia* 85: 375–380. Bradley, P. M. & Morris, J. T. (1992) Effect of salinity on the critical nitrogen concentration of *Spartina alterniflora* Loisel. *Aquatic Botany* 43: 149–161.

Bradshaw, C. J. A., Thompson, C. M., Davis, L. S. & Lalas, C. (1999) Pup density related to terrestrial habitat use by New Zealand fur seals. *Canadian Journal of Zoology* 77: 1579–1586.

Bradshaw, M. J. (1977) *Earth, the Living Planet*. Sevenoaks, UK: Hodder & Stoughton.

Branch, G. M. & Moreno, C. A. (1994) Intertidal and subtidal grazers. In *Rocky Shores: Exploitation in Chile and South Africa*, ed. W. Siegfried, pp. 75–100. New York, USA: Springer-Verlag.

Branch, G. M. & Odendaal, F. (2003) Marine protected areas and wave action: impacts on a South African limpet, *Cymbula oculus. Biological Conservation* 114: 255–269.

Branch, G. M. & Steffani, C. N. (2004) Can we predict the effects of alien species? A case-history of the invasion of South Africa by *Mytilus galloprovincialis* (Lamarck). *Journal* of Experimental Marine Biology and Ecology 300: 189–215.

Branch, G. M., Eekhout, S. & Bosman, A. L. (1990) Shortterm effects of the Orange River floods on the intertidal rocky-shore communities of the open coast. *Transactions of* the Royal Society of South Africa 47: 331–354.

Bravard, J. P. (1987) *Le Rhône du Léman à Lyon*. Lyon, France: La Manufacture.

Bray, R. N. (1981) Influence of water currents and zooplankton densities on daily foraging movements of blacksmith, *Chromis punctipinnis*, a planktivorous reef fish. *Fishery Bulletin of the US* 78: 829–841.

Breen, P. A. & Mann, K. H. (1976) Changing lobster abundance and the destruction of kelp beds by sea urchins. *Marine Biology* 34: 137–142.

Breitburg, D. L., Baxter, J. W., Hatfield, C. A., et al. (1998) Understanding effects of multiple stressors: ideas and challenges. In: Successes, Limitations, and Frontiers in Ecosystem Science, ed. M. L. Pace & P. M. Groffman, pp. 416–431. New York, USA: Springer-Verlag.

Breitburg, D. L., Sanders, J. G., Gilmour, C. C., *et al.* (1999) Variability in responses to nutrients and trace elements, and transmission of stressor effects through an estuarine food web. *Limnology and Oceanography* **44**: 837–863.

Breitfuss, M. J., Connolly, R. M. & Dale, P. E. R. (2003) Mangrove distribution and mosquito control: transport of *Avicennia marina* propagules by mosquito-control runnels in southeast Queensland saltmarshes. *Estuarine, Coastal and Shelf Science* 56: 573–579. Bren, L. J. (1988) Effects of river regulation on flooding of a riparian red gum forest on the River Murray, Australia. *Regulated Rivers: Research and Management* 2: 65–77.

Bricker, S. B. (1993) The history of Cu, Pb, and Zn inputs to Narragansett Bay, Rhode Island as recorded by saltmarsh sediments. *Estuaries* **16**: 589–607.

Bricker, S. B., Clement, C. G., Pirhalla, D. E., Orlando, S. P. & Farrow, D. R. G. (1999) National Estuarine Eutrophication Assessment: Effects of Nutrient Enrichment in the Nation's Estuaries. Silver Spring, MD, USA: National Oceanic and Atmospheric Administration, National Ocean Service, Special Projects Office and the National Centers for Coastal Ocean Science.

Bricker-Urso, S., Nixon, S. W., Cochran, J. K., Hirschberg, D. J. & Hunt, C. (1989) Accretion rates and sediment accumulation in Rhode Island salt marshes. *Estuaries* 12: 300–317.

Brierley, A. S. & Thomas, D. N. (2002) Ecology of Southern Ocean pack ice. Advances in Marine Biology 43: 171–276.

Brierley, A. S., Fernandes, P. G., Brandon, M. A., et al. (2002) Antarctic krill under sea ice: elevated abundance in a narrow band just south of ice edge. *Science* 295: 1890–1892.

Briers, R. A., Cariss, H. M. & Gee, J. H. R. (2002) Dispersal of adult stoneflies (Plecoptera) from upland streams draining catchments with contrasting land-use. *Archiv für Hydrobiologie* 155: 627–644.

Briers, R. A., Gee, J. H. R., Cariss, H. H. & Geoghan, R. (2004) Interpopulation dispersal by adult stoneflies detected by stable isotope enrichment. *Freshmater Biology* 49: 425–431.

Brinson, M. M. (1990) Riverine forests. In: *Forested Wetlands*, ed. A. E. Lugo, M. M. Brinson & S. Brown, pp. 87–141. Amsterdam, the Netherlands: Elsevier.

Brinson, M. M. (1993) A Hydrogeomorphic Classification for Wetlands, Technical Report No. WRP-DE-4. Washington, DC, USA: US Army Corps of Engineers. Available online at http://el.erdc.usace.army.mil/wetlands/pdfs/ wrpde4.pdf

Brinson, M. M. & Malvárez, A. I. (2002) Temperate freshwater wetlands: types, status, and threats. *Environmental Conservation* 29: 115–133.

Brito, M. A., Sobrevila, C., Dalponte, J. C. & Borges, G. A. (undated) Setting Conservation Priorities in the State of Mato Grosso, Brazil, Relatório ao Centro de Dados para Conservação. Cuiabá, Brasil: Fundação Estadual de Meio Ambiente. Brock, J. H. (1994) *Tamarix* spp. (salt cedar), and invasive exotic woody plant in arid and semi-arid riparian habitats of western USA. In: *Ecology and Management of Invasive Riverside Plants*, ed. L.C. de Waal, L. E. Child, P. M. Wade & J. H. Brock, pp. 27–44. New York, USA: John Wiley.

Brönmark, C. & Hansson, L.-A. (2002) Environmental issues in lakes and ponds: current state and perspectives. *Environmental Conservation* 29: 290–306.

Brooks, T. M., Mittermeier, R. A., Mittermeier, C. G., et al. (2002) Habitat loss and extinction in the hotspots of biodiversity. *Conservation Biology* 16: 909–923.

Brough, C. N. & White, K. N. (1990) Localization of metals in the gastropod *Littorina saxatilis* (Prosobranchia, Littorinoidea) from a polluted site. *Acta Zoologica* 71: 77–88.

Brouns, J. J. (1994) Seagrasses and climate change. In: Impact of Climate Change on Ecosystems and Species: Marine and Coastal Ecosystems, ed. J. C. Pernetta, R. Leemnas, D. Elder & S. Humphrey, pp. 59–71, Gland, Switzerland: IUCN.

Broussard, J. (1983) National Audubon Society v. Superior Court, 33 Cal. 3d 419, California Supreme Court, CA, USA.

Browman, H. I., Rodriguez, C. A., Beland, F., et al. (2000) Impact of ultraviolet radiation on marine crustacean zooplankton and ichthyoplankton: a synthesis of results from the estuary and Gulf of St Lawrence, Canada. Marine Ecology Progress Series 199: 281–311.

Brown, A. C. (1983) The ecophysiology of sandy-beach animals: a partial review. In: *Sandy Beaches as Ecosystems*, ed. A. McLachlan & T. Erasmus, pp. 575–605. The Hague, the Netherlands: Dr W. Junk.

Brown, A. C. (1985) The Effects of Crude Oil Pollution on Marine Organisms, Report No. 99. Pretoria, South Africa: South African National Scientific Programmes.

Brown, A. C. (1994) Plutonium and marine life. *Transactions of the Royal Society of South Africa* 49: 213–224.

Brown, A. C. (1996) Behavioural plasticity as a key factor in the survival and evolution of the macrofauna on exposed sandy beaches. *Revista Chilena de Historia Natural* **69**: 469–474.

Brown, A. C. (2001) Biology of sandy beaches. In: *Encyclopedia of Ocean Sciences*, vol. 5, ed. J. H. Steele,
S. A. Thorpe & K. K. Turekian, pp. 2496–2504. London,
UK: Academic Press.

Brown, A. C. & McLachlan, A. (1990) *Ecology of Sandy Shores.* Amsterdam, the Netherlands: Elsevier.

Brown, A. C. & McLachlan, A. (2002) Sandy shores ecosystems and the threats facing them: some predictions for the year 2025. *Environmental Conservation* 29: 62–77. Brown, A. C., Davies, B. R., Day, J. A. & Gardiner, A. J. C. (1991) Chemical pollution loading of False Bay. *Transactions* of the Royal Society of South Africa 47: 703–716.

Brown, A. C., McLachlan, A., Kerley, G. I. H. & Lubke, R. A. (2000) Functional ecosystems: sandy beaches and dunes. In: *Summary Marine Biodiversity Status Report for South Africa*, ed. B. D. Durham & J. C. Pauw, pp. 4–5. Pretoria, South Africa: National Research Foundation.

Brown Jr, E., Busiahn, T., Jones, M. & Argyle, R. (1999) Allocating Great Lakes forage bases in response to multiple demand. In: *Great Lakes Fisheries Policy and Management*, ed. W. W. Taylor & C. P. Ferreri, pp. 385–394. East Lansing, MI, USA: Michigan State University Press.

Brown, K. (2002) Water scarcity: forecasting the future with spotty data. *Science* 297: 926–927.

Brown, L. R. (2003) *Plan B: Rescuing a Planet under Stress and a Civilization in Trouble*. London, UK: W.W. Norton.

Brown, R., Ebener, M. & Gorenflo, T. (1999) Great Lakes commercial fisheries: historical overview and prognosis for the future. In: *Great Lakes Fisheries Policy and Management*, ed. W. W. Taylor & C. Ferreri, pp. 307–354. East Lansing, MI, USA: Michigan State University Press.

Brünig, E. F. (1990) Oligotropic forested wetlands in Borneo. In: *Ecosystems of the World*, vol.15, *Forested Wetlands*, ed. A. E. Lugo, M. Brinson & S. Brown, pp. 299–334. Amsterdam, the Netherlands: Elsevier.

Brunke, M. & Gonser, T. (1997) The ecological significance of exchange processes between rivers and groundwater. *Freshwater Biology* 37: 1–33.

Brunsden, D. & Moore, R. (1999) Engineering geomorphology on the coast: lessons from west Dorset. *Geomorphology* 31: 391–409.

Bryan, G. W. & Langston, W. J. (1992) Bioavailability, accumulation, and effects of heavy metals in sediments with special reference to United Kingdom estuaries: a review. *Environmental Pollution* 76: 89–131.

Bryan, G. W., Gibbs, P. E., Hummerstone, L. G. & Burt, G. R. (1986) The decline of the gastropod *Nucella lapillus* around southwest England: evidence for the effect of tributyl tin from antifouling paints. *Journal of the Marine Biological Association of the United Kingdom* 66: 611–640.

Bryant, D., Burke, L., McManus, J. & Spalding, M. (1998) *Reefs at Risk.* Washington, DC, USA: World Resources Institute.

Bryant, J. C. & Chabreck, R. H. (1998) Effects of impoundment on vertical accretion of coastal marsh. *Estuaries* 21: 416–422. Budd, A. F. (2000) Diversity and extinction in the Cenozoic history of Caribbean reefs. *Coral Reefs* 19: 25–35.

Budd, J., Beeton, A., Stumpf, R., Culver, D. & Kerfoot, C. (2001) Satellite observations of *Microcystis* blooms in western Lake Erie. *Verhandlungen Internationale Vereinigung für Limnologie* 27: 3787–3793.

Buddemeier, R. W. & Smith, S. V. (1988) Coral reef growth in an era of rapidly rising sea level: predictions and suggestions for long-term research. *Coral Reefs* 7: 51–56.

Buijse, A. D., Coops, H., Staras, M., et al. (2002) Restoration strategies for river floodplains along large lowland rivers in Europe. Freshwater Biology 47: 889–907.

Bujang, J. S. & Zakeia, M. H. (2003) The seagrasses of Malaysia. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 166–176. Berkeley, CA, USA: University of California Press.

Bulgarian Ministry of Agriculture and Forests (2001) Strategy for the protection and restoration of floodplain forests on the Bulgarian Danube islands. Unpublished report. Vienna, Austria: WWF International Danube Carpathian Programme.

Bulkley, J., Wright, S. & Wright, D. (1984) Preliminary study of the diversion of 283 m³ s⁻¹ (10000 cfs) from Lake Superior to the Missouri River basin. *Journal of Hydrology* **68**: 461–472.

Bull, D., Kemp, A. E. S. & Weedon, G. P. (2000) A 160-ky-old record of El Niño–Southern Oscillation in marine production and coastal run off from Santa Barbara Basin, California, USA. *Geology* 28: 1007–1010.

Bumb, B. L. & Bannante, C. A. (1996) The Role of Fertilizer in Sustaining Food Security and Protecting the Environment to 2020, IFPRI Discussion Paper No. 17. Washington, DC, USA: World Resources Institute.

Bunn, S. E. & Arthington, A. H. (2002) Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management* 30: 455–467.

Bunn, S. E., Boon, P. I., Brock, M. A. & Schofield, N. J., eds. (1997) National Wetlands R&D Program Scoping Review, Special Publication No. 1/97. Canberra, ACT, Australia: Land and Water Resources R&D Corporation.

Bunt, J. S. (1992) Introduction. In: *Tropical Mangrove Ecosystems*, ed. A. I. Robertson & D. M. Alongi, pp. 1–6. Washington, DC, USA: American Geophysical Union.

Burby, R. J. (2002) Flood insurance and floodplain management: the US experience. *Environmental Hazards* 3: 111–119. Burd, F. (1992) Erosion and Vegetation Change on the Salt Marshes of Essex and North Kent Between 1973 and 1978, Research and Survey in Nature Conservation No. 42. Peterborough, UK: Nature Conservancy Council.

Burdick, D. M. & Konisky, R. A. (2003) Determinants of expansion of *Phragmites australis*, common reed, in natural and impacted coastal marshes. *Estuaries* 26: 407–416.

Burdick, D. M. & Short, F. T. (1999) The effects of boat docks on eelgrass beds in coastal waters of Massachusetts. *Environmental Management* 23: 231–240.

Burdick, D. M., Short, F. T. & Wolf, J. (1993) An index to assess and monitor the progression of the wasting disease in eelgrass, *Zostera marina*. *Marine Ecology Progress Series* 94: 83–90.

Burger, J. (1991) Foraging behaviour and effects of human disturbance on the Piping Plover (*Charadrius melodus*). *Journal of Coastal Research* 7: 39–52.

Burger, J. & Gochfeld, M. (1991) Human activity influence and diurnal and nocturnal foraging of Sanderlings (*Calidris alba*). Condor 93: 259–265.

Burger, J., Gochfeld, M. & Niles, L. J. (1995) Ecotourism and birds in coastal New Jersey: contrasting responses of birds, tourists and managers. *Environmental Conservation* 22: 56–65.

Burke, L., Kura, Y., Kassem, K., et al. (2001) Pilot Analysis of Global Ecosystems (PAGE): Coastal Ecosystems.
Washington, DC, USA: World Resources Institute.

Burkett, V. & Kusler, J. (2000) Climate change: potential impacts and interactions in wetlands of the United States. *Journal of the American Water Resources* Association 36: 313–320.

Burkholder, J. M. (1998) Implications of harmful microalgae and heterotrophic dinoflagellates in management of sustainable marine fisheries. *Ecological Applications* 8 (Suppl.): S37–S62.

Burns, K. A. & Codi, S. (1998) Contrasting impacts of localized versus catastrophic oil spills in mangrove sediments. *Mangroves and Salt Marshes* 2: 63–74.

Burns, K. A. & Yelle-Simmons, L. (1994) The Galeta oil spill. IV. Relationship between sediment and organism hydrocarbon loads. *Estuarine, Coastal and Shelf Science* 38: 397–412.

Burns, K. A., Levings, S. & Garrity, S. (1993) How many years before mangrove ecosystems recover from a catastrophic oil spill? *Marine Pollution Bulletin* 26: 239–248. Burrows, M. T., Hawkins, S. J. & Southward, A. J. (1992) A comparison of reproduction in co-occurring chthamalid barnacles, *Chthamalus stellatus* (Poli) and *Chthamalus montagui* (Southward). *Journal of Experimental Marine Biology and Ecology* 160: 229–249.

Burrows, M., Kawai, K. & Hughes, R. (1999) Foraging by mobile predators on a rocky shore: underwater TV observations of movements of blennies *Lipophrys pholis* and crabs *Carcinus maenas*. *Marine Ecology Progress Series* 187: 237–250.

Busnita, T. (1967) Die Ichthyofauna der Donau. In: Limnologie der Donau, ed. R. Liepolt, pp. 198–224. Stuttgart, Germany: Schweizerbartsche Verlagsbuchhandlung.

Bussau, C., Schriever, G. & Thiel, H. (1995) Evaluation of abyssal metazoan meiofauna from a manganese nodule area of the eastern South Pacific. *Vie Milieu* **45**: 39–48.

Bustamante, R. H. & Branch, G. M. (1996a) The dependence of intertidal consumers on kelp-derived organic matter on the west coast of South Africa. *Journal of Experimental Marine Biology Ecology* 196: 1–28.

Bustamante, R. H. & Branch, G. M. (1996b) Large scale patterns and trophic structure of southern African rocky shores: the roles of geographic variation and wave exposure. *Journal of Biogeography* 23: 339–351.

Bustamante, R. H., Branch, G. M. & Eekhout, S. (1995*a*) Maintenance of an exceptional intertidal grazer biomass in South Africa: subsidy by subtidal kelps. *Ecology* **76**: 2314–2329.

Bustamante, R. H., Branch, G. M., Eekhout, S., et al. (1995b) Gradients of intertidal productivity around the coast of South Africa and their relationship with consumer biomass. *Oecologia* 102: 189–201.

Bustnes, J. O., Lonne, O. J., Skjoldal, H. R., et al. (1995)
Sea ducks as predators on sea urchins in a northern kelp forest. In: *Ecology of Fjords and Coastal Waters*, ed.
H. R. Skjoldal, C. Hopkins & K. E. Erikstad, pp. 599–608.
Amsterdam, the Netherlands: Elsevier.

Busuioc, A., Chen, D. L. & Hellström, C. (2001) Performance of statistical downscaling models in GCM validation and regional climate change estimates: application for Swedish precipitation. *International Journal of Climatology* 21: 557–578.

Cabello-Pasini, A., Lara-Turrent, C. & Zimmerman, R. C. (2002) Effects of storms on photosynthesis, carbohydrate content and survival of eelgrass populations from a coastal lagoon and the adjacent open ocean. *Aquatic Botany* 74: 149–164.

Caddy, J., Csirke, J., Garcia, S. & Grainger, R. (1998) How pervasive is 'Fishing down marine food webs'? *Science* 282: 183.

CAFF (Arctic Council Working Group on Conservation of Arctic Flora and Fauna). (2001) *Arctic Flora and Fauna: Status and Conservation*. Helsinki, Finland: Edita.

Cahill, T. A., Gill, T. E., Reid, J. S., Gearhart, E. A. & Gillette, D. A. (1996) Saltating particles, playa crusts and dust aerosols at Owens (dry) Lake, California. *Earth Surface Processes and Landforms* 21: 621–639.

Caillet, G. M., Andrews, A. H., Burton, E. J., et al. (2001) Age determination and validation studies of marine fishes: do deep-dwellers live longer? *Experimental Gerontology* 36: 739–764.

Cairns Jr, J. & Heckman, J. R. (1996) Restoration ecology: the state of an emerging field. *Annual Review of Energy and the Environment* **21**: 167–189.

Caldeira, K. & Wickett, M. E. (2003) Anthropogenic carbon and ocean pH. *Nature* **425**: 365.

Callaway, J. C. & Josselyn, M. N. (1992) The introduction and spread of smooth cordgrass *(Spartina alterniflora)* in South San Francisco Bay. *Estuaries* 15: 218–226.

Cambell, G. W. & Lee, D. S. (1996) Atmospheric deposition of sulphur and nitrogen species in the UK. *Freshwater Biology* 36: 151–167.

Cambridge, M. L., Chiffings, A. W., Brittan, C., Moore, L. & McComb, A. J. (1986) The loss of seagrass in Cockburn Sound, Western Australia. II. Possible causes of seagrass decline. *Aquatic Botany* 24: 269–285.

Campana, S. E., Zwanenburg, K. C. T. & Smith, J. N. (1990) ²¹⁰Pb/²²⁶Ra determination of longevity in red fish. *Canadian Journal of Fisheries and Aquatic Sciences* 47: 163–165.

Campbell, I. D., Campbell, C., Yu, Z. C., Vitt, D. H. & Apps, M. J. (2000) Millennial-scale rhythms in peatlands in the western interior of Canada and in the global carbon cycle. *Quaternary Research* 54: 155–158.

Campbell, L., Hecky, R. & Dixon, D. (2003) Review of mercury in Lake Victoria (East Africa): implications for human and ecosystem health. *Journal of Toxicology and Environmental Health B* 6: 325–356.

Carbiener, R. & Schnitzler, A. (1990) Evolution of major pattern models and processes of alluvial forests of the Rhine in the rift valley (France/Germany). *Vegetatio* 88: 115–129.

Carbonell, M., Nathai-Gyan, N. & Finlayson, C. M., eds. (2001) Science and Local Communities: Strengthening Partnerships for Effective Wetland Management. Memphis, TN, USA: Ducks Unlimited. Cardinale, B. J., Palmer, M. A. & Collins, S. L. (2002) Species diversity enhances ecosystem functioning through interspecific facilitation. *Nature* 415: 426–429.

Cardoso, P. G., Pardal, M. A., Raffaelli, D., Baeta, A. & Marques, J. C. (2004) Macroinvertebrate response to different species of macroalgal mats and the role of disturbance history. *Journal of Experimental Marine Biology* and Ecology 308: 207–220.

Cargill, S. M. & Jefferies, R. L. (1984) Nutrient limitation of primary production in a sub-arctic salt marsh. *Journal of Applied Ecology* 21: 657–668.

Carlson, D. B., O'Bryan, P. D. & Rey, J. R. (1994)
The management of Florida's (USA) salt marsh impoundments for mosquito control and natural resource enhancement. In: *Global Wetlands: Old World and New*, ed.
W. J. Mitsch, pp. 805–814. Amsterdam, the Netherlands: Elsevier.

Carlton, J. T. (1989) Man's role in changing the face of the oceans: biological conservation of near-shore environments. *Conservation Biology* 3: 265–273.

Carlton, J. T. (1999) Molluscan invasions in marine and estuarine communities. *Malacologia* 41: 439–454.

Carlton, J. T. (2000) Global change and biological invasions in the oceans. In: *Invasive Species in a Changing World*, ed. H. A. Mooney & R. J. Hobbs, pp. 31–53. Washington, DC, USA: Island Press.

Carlton, J. T. & Geller, J. B. (1993) Ecological roulette: the global transport of nonindigenous marine organisms. *Science* 261: 78–82.

Carpenter, R. C. (1988) Mass-mortality of a Caribbean sea urchin: immediate effects on community metabolism and other herbivores. *Proceedings of the National Academy of Sciences of the USA* 85: 511–514.

Carpenter, S. R., Caraco, N. F., Correll, D. L., et al. (1998) Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications* 8: 559–568.

Carr, M. H. (1989) Effects of macroalgal assemblages on the recruitment of temperate zone reef fishes. *Journal of Experimental Marine Biology and Ecology* **126**: 59–76.

Carreiro-Silva, M., McClanahan, T. R. & Kiene, W. E. (2005) The role of inorganic nutrients and herbivory in controlling microbioerosion of carbonate substrate. *Coral Reefs* 24: 214–221.

Carricart-Ganivet, J. P. & Beltran-Torres, A. U. (2000) Skeletal extension, density and calcification rate of the reef building coral *Montastraea annularis* (Ellis and Solander) in the Mexican Caribbean. *Bulletin of Marine Science* **66**: 215–224. Carroll, P. & Crill, P. M. (1997) Carbon balance of a temperate poor fen. *Global Biogeochemical Cycles* 11: 349–356.

Casagrande, R. A., Balme, G. & Blossey, B. (2003) *Rhizedra lutosa*, a natural enemy of *Phragmites australis* in North America. *Estuaries* 26: 602–606.

Castellanos, E. M., Figueroa, M. E. & Davy, A. J. (1994) Nucleation and facilitation in saltmarsh succession: interactions between *Spartina maritima* and *Arthrocnemum perenne. Journal of Ecology* 82: 239–248.

Castilla, J. C. (1996) Copper mine tailing disposal in northern Chile rocky shores: *Enteromorpha compressa* (Chlorophyta) as a sentinel species. *Environmental Monitoring and Assessment* **40**: 171–184.

Castilla, J. C. (1999) Coastal marine communities: trends and perspectives from human-exclusion experiments. *Trends in Ecology and Evolution* 14: 280–283.

Castilla, J. C. (2000) Roles of experimental marine ecology in coastal management and conservation. *Journal of Experimental Marine Biology and Ecology* **250**: 3–21.

Castilla, J. C. (2001) Marine ecosystems, human impacts on. Encyclopaedia of Biodiversity 4: 27–35.

Castilla, J.C & Camus, P. (1992) The Humboldt–El Niño scenario: coastal benthic resources and anthropogenic influences, with particular reference to the 1982/83 ENSO. *South African Fournal of Marine Science* **12**: 703–712.

Castilla, J. C. & Duran, L. R. (1985) Human exclusion from the rocky intertidal zone of Central Chile: the effects on *Concholepas concholepas* (Gastropoda). *Oikos* 45: 391–399.

Castilla, J. C. & Moreno, C. A. (1982) Sea urchins and *Macrocystis pyrifera*: experimental test of their ecological relations in southern Chile. In: *Echinoderms: Proceedings* of the International Conference, Tampa Bay, FL, USA, ed. J. M. Lawrence, pp. 257–263. Rotterdam, the Netherlands: A. A. Balkema.

Castillo, J. M., Fernández-Baco, L., Castellanos, E. M., et al. (2000) Lower limits of Spartina densiflora and S. maritima in a Mediterranean salt marsh determined by different ecophysiological tolerances. Journal of Ecology 88: 801–812.

Castro, M. S., Driscoll, C. T., Jordan, T. E., Reay, W. G. & Boynton, W. R. (2003) Sources of nitrogen to estuaries in the United States. *Estuaries* 26: 803–814.

Causey, B., Delany, J., Diaz, E., et al. (2002) Status of coral reefs in the US Caribbean and Gulf of Mexico: Florida, Texas, Puerto Rico, US Virgin Islands, Navassa. In: Status of Coral Reefs of the World: 2002, ed. C. Wilkinson, pp. 251–276. Townsville, Queensland, Australia: Australian Institute of Marine Science. Cavalieri, D. J., Gloersen, P., Parkinson, C. L., Comiso, J. C. & Zwally, H. J. (1997) Observed hemispheric asymmetry in global sea ice changes. *Science* 278: 1104–1106.

Cayan, D. R., Redmond, K. T. & Riddle, L. G. (1999) ENSO and hydrologic extremes in the western United States. *Journal of Climatology* **12**: 2881–2893.

Cearreta, A., Irabien, M. J., Leorri, E., *et al.* (2000) Recent anthropogenic impacts on the Bilbao estuary, northern Spain: geochemical and macrofaunal evidence. *Estuarine*, *Coastal and Shelf Science* **50**: 571–592.

Cerda, M. & Castilla, J. C. (2001) Diversity and biomass of macro-invertebrates in intertidal matrices of the tunicate *Pyura praeputialis* (Heller, 1878) in the Bay of Antofagasta, Chile. *Revista Chilena de Historia Natural* 74: 841–853.

Ceulemans, R. & Mousseau, M. (1994) Effects of elevated atmospheric CO₂ on woody plants. *New Phytologist* 127: 425–446.

Chalker-Scott, L. (1995) Survival and sex ratios of the intertidal copepod, *Tigriopus californicus*, following ultraviolet-B (290–320 nm) radiation exposure. *Marine Biology* 123: 799–804.

Chambers, F. M. (1997) Bogs as treeless wastes: the myth and the implications for conservation. In: *Conserving Peatlands*, ed. L. Parkyn, R. E. Stoneman & H. A. P. Ingram, pp. 168–175. Wallingford, UK: CAB International.

Chambers, F. M., Mauquoy, D. & Todd, P. A. (1999*a*) Recent rise to dominance of *Molinia caerulea* in environmentally sensitive areas: new perspectives from palaeoecological data. *Journal of Applied Ecology* **36**: 719–733.

Chambers, R. M., Meyerson, L. A. & Saltonstall, K. (1999b) Expansion of *Phragmites australis* into tidal wetlands of North America. *Aquatic Botany* 64: 261–273.

Chapelle, F. H. (2001) Ground-Water Microbiology and Geochemistry. New York, USA: John Wiley.

Chapman, A. R. O. (1986) Age versus stage: an analysis of age- and size-specific mortality and reproduction in a population of *Laminaria longicruris* Pyl. *Journal of Experimental Marine Biology and Ecology* 97: 113–122.

Chapman, A. R. O. & Johnson, C. R. (1990) Disturbance and organization of macroalgal assemblages in the Northwestern Atlantic. *Hydrobiologia* 192: 77–121.

Chapman, D. M. (1989) Coastal Dunes of New South Wales: Status and Management. Technical Report No. 89/3.
Sydney, New South Wales, Australia: University of Sydney Coastal Studies Unit.

Chapman, J. L. & Reiss, M. (1998) *Ecology: Principles and Applications*, 2nd edn. Cambridge, UK: Cambridge University Press. Chapman, L. J., Balirwa, J., Bugenyi, F. W. B., Chapman, C. & Crisman, T. L. (2001) Wetlands of East Africa: biodiversity, exploitation and policy perspectives. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, vol. 2, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 101–131. Leiden, the Netherlands: Backhuvs.

Chapman, M. G. & Bulleri, F. (2003) Intertidal seawalls: new features of landscape in intertidal environments. *Landscape and Urban Planning* **62**: 159–172.

Chapman, P. M., Paine, M. D., Arthur, A. D. & Taylor, L. A. (1996) A triad study of sediment quality associated with a major, relatively untreated marine sewage discharge. *Marine Pollution Bulletin* 32: 47–64.

Chapman, V. J. (1974) Salt Marshes and Salt Deserts of the World, 2nd edn. Lehre, Germany: Cramer.

Chapman, W. L. & Walsh, J. E. (1993) Recent variations of sea ice and air temperature in high latitudes. *Bulletin of the American Meteorological Society* 74: 33–47.

Charles, C. D., Hunter, D. E. & Fairbanks, R. D. (1997) Interaction between the ENSO and the Asian Monsoon in a coral record of tropical climate. *Science* 277: 925–928.

Charman, D. (2002) *Peatlands and Environmental Change*. Chichester, UK: John Wiley.

Chauhan, M. & Gopal, B. (2001) Biodiversity and management of Keoladeo National Park (India): a wetland of international importance. In: *Biodiversity in Wetlands:* Assessment, Function and Conservation, vol. 2, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 217–256. Leiden, the Netherlands: Backhuys.

Chavez, F. P., Ryan, J., Lluch-Cota, S. E. & Miguel Niquen, C. (2003) From anchovies to sardines and back: multidecadal change in the Pacific Ocean. *Science* 299: 217–221.

Chazottes, V., Le Campion-Alsumard, T. & Peyrot-Clausade, M. (1995) Bioerosion rates on coral reefs: interactions between macroborers, microborers and grazers (Moorea, French Polynesia). *Palaeo* 113: 189–198.

Chen, J., Carlson, B. E. & Del Genio, A. D. (2002) Evidence for strengthening of the tropical general circulation in the 1990s. *Science* 295: 838–841.

Chesney, E. J., Baltz, D. M. & Thomas, R. G. (2000) Louisiana estuarine and coastal fisheries and habitats: perspectives from a fish's eye view. *Ecological Applications* 10: 350–366.

Chesterfield, E. A. (1986) Changes in the vegetation of the river red gum forest at Barmah, Victoria. *Australian Forestry* 49: 4–15. Choat, J. H. & Ayling, A. M. (1987) The relationship between habitat and fish faunas of New Zealand reefs. *Journal of Experimental Marine Biology and Ecology* **110**: 228–284.

Choat, J. H. & Schiel, D. R. (1982) Patterns of distribution and abundance of large brown algae and invertebrate herbivores in subtidal regions of northern New Zealand. *Journal of Experimental Marine Biology and Ecology* 60: 129–162.

Christensen, J. H. & Christensen, O. B. (2002) Severe summertime flooding in Europe. *Nature* 421: 805.

Christensen, T. R. & Keller, M. (2004) Element interactions and trace gas exchange. In: *Interactions of the Major Biogeochemical Cycles: Global Change and Human Impacts*, ed. J. M. Melillo, C. B. Field & B. Moldan, pp. 247–258. Washington, DC, USA: Island Press.

Christensen, T. R., Lloyd, D., Svensson, B., et al. (2002) Biogenic controls on trace gas fluxes in northern wetlands. *Global Change News Letter* 51: 9–15.

Christensen, V. (1998) Fishery-induced changes in the marine ecosystem: insights from models of the Gulf of Thailand. *Journal of Fish Biology* 53 (Suppl. A): 128–142.

Christensen, V. & Pauly, D., eds. (1993) Trophic Models of Aquatic Ecosystems: ICLARM Conference Proceedings 26. Manila, Philippines: International Center for Living Aquatic Resources.

Christensen, V., Guénette, S., Heymans, J. J., *et al.* (2003) Hundred-year decline of North Atlantic predatory fishes. *Fish and Fisheries* 4: 1–24.

Christie, J., Becker, M., Cowden, J. & Vallentyne, J. (1986) Managing the Great Lakes Basin as a home. *Journal of Great Lakes Research* 12: 2–17.

Chung, C.-H. (1985) The effects of introduced *Spartina* grass on coastal morphology in China. *Zeitschrift für Geomorphologie* (Suppl.) 57: 169–174.

Chung, C.-H. (1994) Creation of *Spartina* plantations as an effective measure for reducing coastal erosion in China.
In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 443–452. Amsterdam, the Netherlands: Elsevier.

Chung, C.-H., Zhuo, R. Z. & Xu, G. W. (2004) Creation of *Spartina* plantations for reclaiming Dongtai, China, tidal flats and offshore sands. *Ecological Engineering* **23**: 135–150.

Cincotta, R. P. & Engelman, R. (2000) Nature's Place: Human Population and the Future of Biological Diversity. Washington, DC, USA: Population Action International.

Cisneros-Mata, M. A., Botsford, L. & Quinn, J. F. (1997) Projecting the viability of *Totoba macdonaldi*, a population with unknown age dependent variability. *Ecological Applications* 7: 968–980. Ciupek, R. B. (1986) Protecting wetlands under Clean Water Act 404: EPA's conservation policy on mitigation. *Natural Wetlands Newsletter* 8: 12–15.

Clark, J. R. (1983) *Coastal Ecosystem Management*. Malabar, FL, USA: R.E. Krieger.

Clark, J. R. (1995) Coastal Zone Management Handbook. Boca Raton, FL, USA: Lewis Publishers.

Clark, J. R. (1998) Endangered and Threatened Species: Threatened Status for Johnson's Seagrass, Department of Commerce, National Oceanic and Atmospheric Administration, 50 CFR Part 227. Federal Register, Vol. 63, No. 177. Available online at www.nmfs.noaa.gov/pr/pdfs/fr/fr63-49035.pdf

Clark, J. S., Carpenter, S. R., Barber, M., *et al.* (2001) Ecological forecasts: an emerging imperative. *Science* 293: 657–660.

Clark, M. (1999) Fisheries for orange roughy (*Hoplostethus atlanticus*) on seamounts in New Zealand. Oceanologica Acta 22: 593–602.

Clark, M. R. (1995) Experience with the management of orange roughy (*Hoplostethus atlanticus*) in New Zealand, and the effects of commercial fishing on stocks over the period 1980–1993. In: *Deep-Water Fisheries of the North Atlantic Oceanic Slope*, ed. A. G. Hopper, pp. 251–266. Dordrecht, the Netherlands: Kluwer.

Clark, M. R. & Tracey, D. M. (1994) Changes in a population of orange roughy, *Hoplostethus atlanticus*, with commercial exploitation on the Challenger Plateau, New Zealand. *Fishery Bulletin* 92: 236–253.

Clark, M. R., Finncham, D. J. & Tracey, D. M. (1994) Fecundity of orange roughy (*Hoplostethus atlanticus*) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 28: 193–200.

Clark, M. W. (1998) Management implications of metal transfer pathways from a refuse tip to mangrove sediments. *Science of the Total Environment* 222: 17–34.

Clark, R., ed. (1999) *Global Environment Outlook 2000.* London, UK: Earthscan Publications for United Nations Environment Programme.

Clark, R. B. (1992) *Marine Pollution*, Oxford, UK: Clarendon Press.

Clark, R. B., Frid, C. L. J. & Atrill, M. (1997) Marine Pollution. Oxford, UK: Clarendon Press.

Clark, R. L. & Guppy, J. C. (1988) A transition from mangrove forest to freshwater wetland in the monsoon tropics of Australia. *Journal of Biogeography* 15: 665–684.

Clark, W. C., Mitchell, R. B. & Cash, D. W. (2006) Evaluating the influence of global environmental assessments. In: *Global Environmental Assessments: Information and Influence*, ed. R. B. Mitchell, W. C. Clark, D. W. Cash, and N. M. Dickson, pp. 1–28. Cambridge, MA, USA: MIT Press.

Clarke, A. (1996) The influence of climate change on the distribution and evolution of organisms. In: *Animals and Temperature: Phenotypic and Evolutionary Adaptation*, ed. I. A. Johnston & A. F. Bennett, pp. 377–407. Cambridge, UK: Cambridge University Press.

Clarke, A. & Harris, C. M. (2003) Polar marine ecosystems: major threats and future change. *Environmental Conservation* 30: 1–25.

Clarke, F. W. (1920) *The Data of Geochemistry*, US Geological Survey Bulletin No. 695. Menlo Park, CA, USA: US Geological Survey.

Clarke, T. (2003) Lightning satellite survey. Nature 421: 228.

CLIMAP Project Members (1976) The surface of the ice-age Earth. *Science* 191: 1131–1137.

Cloern, J. E. (2001) Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology Progress Series* 210: 223–253.

Close, A. (1990) The impact of man on the natural flow regime. In: *The Murray*, ed. N. Mackay & D. Eastburn, pp. 61–74. Canberra, ACT, Australia: Murray–Darling Basin Commission.

Clough, B. F., Boto, K. G. & Attiwill, P. M. (1983) Mangroves and sewage : a re-evaluation. In: *Biology and Ecology of Mangroves*: vol. 8, *Tasks for Vegetation Science*, ed.
H. J. Teas, pp. 151–161. The Hague, the Netherlands: Dr W. Junk.

Cluis, D. & Laberge, C. (2002) Analysis of El Niño effect on the discharge of selected rivers in the Asia Pacific region. *Water International* 27: 279–293.

Clymo, R. S. (1965) Experiments on breakdown of Sphagnum in two bogs. Journal of Ecology 53: 747–758.

Clymo, R. S. (1973) The growth of *Sphagnum*: some effects of environment. *Journal of Ecology* 61: 849–869.

Clymo, R. S. (1984) The limits to peat bog growth. *Philosophical Transactions of the Royal Society B* 303: 605–654.

Clymo, R. S. & Hayward, P. M. (1982) The ecology of *Sphagnum*. In: *Bryophyte Ecology*, ed. A. J. E. Smith, pp. 229–289. London, UK: Chapman & Hall.

Clymo, R. S. & Pearce, D. M. E. (1995) Methane and carbon dioxide production in, transport through, and efflux from a peatland. *Philosophical Transactions of the Royal Society of London B* 350: 249–259.

Clymo, R. S., Turunen, J. & Tolonen, K. (1998) Carbon accumulation in peatland. *Oikos* 81: 368–388.

Coale, K. H., Johnson, K. S., Fitzwater, S. E., *et al.* (1996) A massive phytoplankton bloom induced by an

ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. *Nature* 383: 495–501.

- Cobb, K. M., Charles, C. D., Cheng, H. & Edwards, R. L. (2003) El Niño/Southern Oscillation and tropical Pacific climate during the last millennium. *Nature* 424: 271–276.
- Cognetti, G. (2001) Marine eutrophication: the need for a new indicator system. *Marine Pollution Bulletin* 42: 163–164.

Cohen, J. E. (1995) *How Many People Can the Earth Support?* New York, USA: W.W. Norton.

Cohen, J. E. (1997) Estimates of coastal population. *Science* **278**: 1209.

Cohen, J. E. (2003) Human population @ the next half century. *Science* **302**: 1172–1176.

Cohen, J. E., Small, C., Mellinger, A., Gallup, J. & Sachs, J. (1997) Estimates of coastal populations. *Science* 287: 1211–1212.

COHMAP Members (1988) Climatic changes over the last 18 000 years: observations and model simulations. *Science* 241: 1043–1052.

Colborn, T., Davidson, A., Green, S., et al. (1990) Great Lakes, Great Legacy? Washington, DC, USA and Ottawa, Ontario, Canada: The Conservation Foundation and the Institute for Research on Public Policy.

Cole, J., Dunbar, R., McClanahan, T. & Muthiga, N. (2000) Tropical Pacific forcing of decadal variability in SST in the western Indian Ocean. *Science* 287: 617–619.

Cole, J. J., Peierls, B. L., Caraco, N. F. & Pace, M. L. (1993) Nitrogen loading of rivers as a human-driven process. In: *Humans as Components of Ecosystems*, ed. M. J. McDonnell & S. T. A. Pickett, pp. 141–157. New York, USA: Springer-Verlag.

Cole, R. G. & Babcock, R. C. (1996) Mass mortality of a dominant kelp (Laminariales) at Goat Island, North-eastern New Zealand. *Marine Freshwater Resource* 47: 907–911.

Cole, R. G. & Syms, C. (1999) Using spatial pattern analysis to distinguish causes of mortality: an example from kelp in north-eastern New Zealand. *Journal of Ecology* 87: 963–972.

Coleman, R. A., Goss-Custard, J. D., LeV. dit Durell, S. E. A. & Hawkins, S. J. (1999) Limpet *Patella* spp. consumption by oystercatchers *Haematopus ostralegus*: a preference for solitary prey items. *Marine Ecology Progress Series* 183: 253–261.

Coles, R. G. (1996) Coastal Management and Community Coastal Resource Planning in the Asia Pacific Region.
Canberra, ACT, Australia: Winston Churchill Memorial Trust.

Cohen, A. N. & Carlton, J. T. (1998) Accelerating invasion rate in a highly invaded estuary. *Science* 279: 555–562.

Coles, R. G. & Fortes, M. (2001) Protecting seagrass: approach and methods. In: *Global Seagrass Research Methods*, ed. F. T. Short & R. G. Coles, pp. 445–464. Amsterdam, the Netherlands: Elsevier.

Coles, R. G., McKenzie, L. & Campbell, S. (2003*a*) The seagrasses of Eastern Australia. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 131–147. Berkeley, CA, USA: University of California Press.

Coles, R. G., McKenzie, L., Campbell, S., Fortes, M. & Short, F. (2003b) The seagrasses of the Western Pacific Islands. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 177–186. Berkeley, CA, USA: University of California Press.

Coles, S. L. & Al-Rayami, K. A. (1996) Beach tar concentrations on the Muscat coastline, Gulf of Oman, Indian Ocean, 1993–1995. *Marine Pollution Bulletin* 32: 609–614.

Coles, S. L. & Brown, B. E. (2003) Coral bleaching: capacity for acclimatization and adaptation. *Advances in Marine Biology* 46: 183–223.

Coles, S. M. (1979) Benthic microalgal populations on intertidal sediments and their role as precursors to salt marsh development. In: *Ecological Processes in Coastal Environments*, ed. R. L. Jefferies & A. J. Davy, pp. 25–42. Oxford, UK: Blackwell Scientific Publications.

Collie, J. S. & DeLong, A. K. (1999) Multispecies interactions in the Georges Bank fish community. In: *Alaska Sea Grant College Program, Ecosystem Approaches for Fisheries Management*, Publication No. AK-SG-99-01, pp. 187–210. Alaska, USA: Alaska Sea Grant.

Collie, J. S., Escanero, G. A. & Valentine, P. C. (1997) Effects of bottom fishing on the benthic megafauna of Georges Bank. *Marine Ecology Progress Series* 155: 159–172.

Collie, J. S., Hall, S. J., Kaiser, M. J. & Poiner, I. R. (2000) A quantitative analysis of fishing impacts on shelf-sea benthos. *Journal of Animal Ecology* **69**: 785–798.

Collier, L. M. & Pinn, E. H. (1998) An assessment of the acute impact of sea lice treatment with ivermectin on a benthic community. *Journal of Experimental Marine Biology and Ecology* 23: 131–147.

Collingham, Y. C. & Huntley, B. (2000) Impacts of habitat fragmentation and patch size upon migration rates. *Ecological Applications* **10**: 131–144.

Collins, M. A., Belchier, M. & Everson, I. (2003) Why the fuss about toothfish? *Biologist* 50: 116–119.

Congressional Natural Hazards Caucus Work Group (2001) Discussion paper for the Congressional Natural Hazards Caucus [www document]. URL http://www.agiweb.org/ workgroup/discussion_paper0101.html Conlan, K., White, K. N. & Hawkins, S. J. (1992) The hydrography and ecology of a redeveloped brackish-water dock. *Estuarine, Coastal and Shelf Science* 35: 435–452.

Conley, D. J., Stalnacke, P., Pitkanen, H. & Wilander, A. (2000) The transport and retention of dissolved silicate by rivers in Sweden and Finland. *Limnology and Oceanography* 45: 1850–1853.

Connell, J. H. (1961) The influence of interspecific competition and other factors on the distribution of the barnacle *Chthamalus stellatus*. *Ecology* **42**: 710–723.

Connell, J. H. (1972) Community interactions on marine rocky intertidal shores. *Annual Review of Ecology and Systematics* 3: 169–192.

Connell, J. H. (1997) Disturbance and recovery of coral assemblages. *Coral Reefs* 16: S101–S113.

Connolly, R. M. (1999) Saltmarshes as habitat for fish and nektonic crustaceans: challenges in sampling designs and wetlands. *Australian Journal of Ecology* 24: 422–430.

Connolly, R. M., Dalton, A. & Bass, D. A. (1997) Fish use of an inundated flat in a temperate Australian estuary. *Australian Journal of Ecology* 22: 222–226.

Connolly, S. R. & Roughgarden, J. (1998) A latitudinal gradient in Northeast Pacific intertidal community structure: evidence for an oceanographically based synthesis of marine community theory. *American Naturalist* 151: 311–326.

Conover, R. J. & Huntley, M. (1991) Copepods in ice covered seas: distribution, adaptations to seasonally limited food, metabolism, growth patterns and life cycle strategies in polar seas. *Journal of Marine Systems* 2: 1–41.

Conserve Africa Foundation (2005) Poverty and environment in Africa: an overview [www document]. URL http:// www.conserveafrica.org.uk/poverty.html

Constable, A. J., de La Mare, W. K., Agnew, D. J., Everson, I.
& Miller, D. (2000) Managing fisheries to conserve the Antarctic marine ecosystem: practical implementation of the Convention on the Conservation of Antarctic Marine Living Resources. *ICES Journal of Marine Science* 57: 778–791.

Contreras-Balderas, S. & Lozano-Vilano, M. L. (1994) Water, endangered fishes, and development perspectives in arid lands of Mexico. *Conservation Biology* 8: 379–387.

Convey, P., Scott, D. & Fraser, W. R. (2003) Biophysical and habitat changes in response to climate alteration in the Arctic and Antarctic. *Advances in Applied Biodiversity Science* 4: 79–84. Cook, P., Robbins, J., Endicott, D., et al. (2003) Effects of aryl hydrocarbon receptor mediated early life stage toxicity on lake trout populations in Lake Ontario during the 20th century. Environmental Science and Technology 37: 3864–3877.

Correa, J. A., Castilla, J. C., Ramirez, M., et al. (1999) Copper, copper mine tailings and their effect on marine algae in northern Chile. *Journal of Applied Phycology* 11: 57–67.

Corredor, J. E., Howarth, R. W., Twilley, R. R. & Morell, J. M. (1999) Nitrogen cycling and anthropogenic impact in the tropical interamerican seas. *Biogeochemistry* 46: 163–178.

Corredor, J. E., Morrell, J. M. & Del Castillo, C. E. (1990) Persistence of spilled crude oil in a tropical marine environment. *Marine Pollution Bulletin* **21**: 385–388.

Cortes, J. (1993) Comparison between Caribbean and eastern Pacific coral reefs. *Revistade Biologica Tropicale* **41** (Suppl. 1): 19–21.

Cortes, J., Macintyre, I. G. & Glynn, P. W. (1994) Holocene growth history of an eastern Pacific fringing reef, Punta Islotes, Costa Rica. *Coral Reefs* 13: 65–73.

Cosgrove, W.J. & Rijsberman, F.R. (2000) *World Water Vision: Making Water Everybody's Business.* London, UK: Earthscan Publications.

Costa, L. T., Farinha, J. C., Hecker, N. & Tomàs Vives, P., eds. (1996) Mediterranean Wetland Inventory, vol. 1, A Reference Manual [www document]. URL http://www. wetlands.org/RSDB/Default.htm

Costanza, R., d'Arge, R., de Groot, R., *et al.* (1997) The value of the world's ecosystem services and natural capital. *Nature* **387**: 253–260.

Costello, M. J. & Read, P. (1994) Toxicity of sewage sludge to marine organisms: a review. *Marine Environmental Research* 37: 23–42.

Coulson, J. C. & Butterfield, J. (1978) An investigation of the biotic factors determining the rates of plant decomposition on blanket bog. *Journal of Ecology* 66: 631–650.

Coulter, G. & Mubamba, R. (1993) Conservation in Lake Tanganyika, with special reference to underwater parks. *Conservation Biology* 7: 678–685.

Coutant, C. C. (1981) Foreseeable effects of CO₂-induced climate change. Fresh-water concerns. *Environmental Conservation* 8: 285–297.

Covich, A. P. (1996) Stream biodiversity and ecosystem processes. *Bulletin of the North American Benthological Society* 13: 294–303.

Covich, A. P., Austen, M., Bärlocher, F., et al. (2004) The role of biodiversity in the functioning of freshwater and marine benthic ecosystems. *BioScience* 54: 767–775. Covich, A. P., Fritz, S. C., Lamb, P. J., et al. (1997) Potential effects of climate change on aquatic ecosystems of the great plains of North America. *Hydrological Processes* 11: 993–1021.

Covich, A. P., Palmer, M. A. & Crowl, T. A. (1999) Role of benthic invertebrate species in freshwater ecosystems. *BioScience* 49: 119–127.

Cowardin, L. M., Carter, V., Golet, F.C & LaRoe, E. T. (1979) Classification of Wetlands and Deepwater Habitats of the United States, Biological Services Program No. FWS/ OBS-79/31. Washington, DC, USA: US Fish and Wildlife Service, US Department of the Interior.

Cowen, R. K. (1983) The effect of sheephead (*Semicossyphus pulcher*) predation on red sea urchin populations: an experimental analysis. *Oecologia* **58**: 249–255.

Cowen, R. K., Lwiza, K. M. M., Sponaugle, S., Paris, C. B. & Olson, D. B. (2000) Connectivity of marine populations: open or closed? *Science* 287: 857–859.

Craft, C. B. & Casey, W. P. (2000) Sediment and nutrient accumulation in floodplain and depressional freshwater wetlands of Georgia, USA. *Wetlands* **20**: 323–332.

Crain, A. D., Bolten, A. B. & Bjorndal, K. A. (1995) Effects of beach nourishment on sea turtles: review and research initiatives. *Restoration Ecology* 3: 95–104.

Crane, R. G. & Barry, R. G. (1984) The influence of clouds on climate with a focus on high latitude interactions. *Journal of Climatology* 4: 71–93.

Crawford, R. E. (2002) Secondary wake turbidity from small boat operation in a shallow sandy bay. *Journal of Coastal Research* Special Issue **37**: 49–64.

Creasey, S. S. & Rogers, A. D. (1999) Population genetics of bathyal and abyssal organisms. *Advances in Marine Biology* 35: 1–151.

Creed, J. C. (2003) Seagrasses of South America: Brazil, Argentina and Chile. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 263–270. Berkeley, CA, USA: University of California Press.

Creed, J. C., Phillips, R. C. & Van Tussenbroek, B. I. (2003) The seagrasses of the Caribbean. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 254–262. Berkeley, CA, USA: University of California Press.

Cresser, M. S. (2000) The critical loads concept: milestone or millstone for the new millennium? *Science of the Total Environment* 249: 51–62.

Cronin, M., Davies, I. M., Newton, A., et al. (1998) Trace metal concentrations in deep sea fish from the North Atlantic. Marine Environmental Research 45: 225–238. Crooks, S., Schutten, J., Sheern, G. D., Pye, K. & Davy, A. J. (2002) Drainage and elevation as factors in the restoration of salt marsh in Britain. *Restoration Ecology* **10**: 591–602.

Crow, G.E. (1993) Species diversity in aquatic angiosperms: latitudinal patterns. *Aquatic Botany* 44: 229–258.

Crowe, T. P., Thompson, R. C., Bray, S. & Hawkins, S. J. (2000) Impacts of anthropogenic stress on rocky intertidal communities. *Journal of Aquatic Ecosystem Stress and Recovery* 7: 273–297.

Croxall, J. P. & Nicol, S. (2004) Management of Southern Ocean fisheries global forces and future sustainability. *Antarctic Science* 16: 568–584.

Croxall, J. P., Prince, P. A., Rothery, P. & Wood, A. G. (1997)
Population changes in albatrosses at South Georgia. In: *Albatross Biology and Conservation*, ed. G. Robertson &
R. Gales, pp. 69–83. Chipping Norton, NSW, Australia: Surrey Beatty.

Croxall, J. P., Trathan, P. N. & Murphy, E. J. (2002) Environmental change and Antarctic seabird populations. *Science* **297**: 1510–1514.

Crunkilton, R. (1985) Subterranean contamination of Meramec Spring by ammonium nitrate and urea fertilizer and its implication on rare biota. In: *Proceedings of the* 1984 National Cave Management Symposium, 25, pp. 151–158. Columbia, MO, USA: Missouri Speleological Survey.

Crutzen, P. (2006) Albedo enhancement by stratospheric sulphur injections: a contribution to resolve a policy dilemma? *Climatic Change* 77: 211–220.

Cryer, M., Hartill, B. & Oshea, S. (2002) Modification of marine benthos by trawling: toward a generalization for the deep ocean. *Ecological Applications* 12: 1824–1839.

Cullen, P. (2003) Salinity. In: *Ecology: An Australian Perspective*, ed. P. Attiwill & B. Wilson, pp. 474–488. Melbourne, Victoria, Australia: Oxford University Press.

Culver, D. C. & Holsinger, J. R. (1992) How many species of troglobites are there? *National Speleological Society Bulletin* 54: 79–80.

Culver, D. C., Jones, W. K. & Holsinger, J. R. (1992) Biological and hydrological investigation of the Cedars, Lee County, Virginia: an ecologically significant and threatened karst area. In: *Proceedings of the 1st International Conference* on Groundwater Ecology, ed. J. A. Stanford & J. J. Simons, pp. 281–290. Bethesda, MD, USA: American Water Research Association.

Curran, M. A. J., van Ommen, T. D., Morgan, V. I., Phillips, K. L. & Palmer, A. S. (2003) Ice core evidence for Antarctic sea ice decline since the 1950s. *Science* **302**: 1203–1206. Curry, J. A. & Webster, P. J. (1999) *Thermodynamics of Atmospheres and Oceans.* San Diego, CA, USA: Academic Press.

Curry, J. A., Schramm, J. L., Serreze, M. C. & Ebert, E. E. (1995) Water vapor feedback over the Arctic Ocean. *Journal* of *Geophysical Research* **100**: 14223–14229.

Curtis, P. S. & Wang, X. (1998) A meta-analysis of elevated CO₂ effects on woody plant mass, form, and physiology. *Oecologia* 113: 299–313.

Curtis, P. S., Balduman, L. M., Drake, B. G. & Whigham, D. F. (1990) Elevated atmospheric CO₂ effects on belowground processes in C₃ and C₄ estuarine marsh communities. *Ecology* 71: 2001–2006.

Curtis, P. S., Drake, B. G., Leadley, P. W., Arp, W. J. & Whigham, D. F. (1989*a*) Growth and senescence in plant communities exposed to elevated CO₂ concentrations on an estuarine marsh. *Oecologia* 78: 20–26.

Curtis, P. S., Drake, B. G. & Whigham, D. F. (1989*b*) Nitrogen and carbon dynamics in C₃ and C₄ estuarine plants grown under elevated CO₂ *in situ. Oecologia* 78: 297–301.

Cutshall, N. H., Naidu, J. R. & Pearcy, W. G. (1978) Mercury concentrations in Pacific hake, *Merluccius productus* (Ayres), as a function of length and latitude. *Science* 200: 1489–1491.

Cwynar, L. C. & Spear, R. W. (1991) Reversion of forest to tundra in the central Yukon. *Ecology* 72: 202–212.

Czaya, E. (1981) *Rivers of the World*. New York, USA: Van Nostrand Reinhold.

Daan, R. & Mulder, M. (1996) On the short-term and long-term impact of drilling activities in the Dutch sector of the North Sea. *ICES Journal of Marine Science* 53: 1036–1044.

Dacey, J. W. H., Drake, B. G. & Klug, M. J. (1994) Stimulation of methane emission by carbon dioxide enrichment of marsh vegetation. *Nature* 370: 47–49.

Dadnadji, K. K. & van Wetten, J. C. J. (1993) Traditional management systems and irrigation of small scale interventions in the Lagone floodplains of Chad. In: *Towards the Wise Use of Wetlands*, ed. T. J. Davis,

pp. 74–81. Gland, Switzerland: Ramsar Convention Bureau. Dadswell, M. (1974) *Distribution, Ecology, and Postglacial*

Dispersal of Certain Crustaceans and Fishes in Eastern North America, Publications in Zoology No. 11. Ottawa, Ontario, Canada: National Museums of Canada.

Dahdouh-Guebas, F. (2002) The use of remote sensing and GIS in the sustainable management of tropical coastal ecosystems. *Environment, Development and Sustainability* 4: 93–112. Dahl, T. E. (1990) Wetlands: Losses in the United States 1780s to 1980s. Washington, DC, USA: US Fish and Wildlife Service.

Dahl, T. E. & Zoltai, S. C. (1997) Forested northern wetlands of North America. In: Northern Forested Wetlands: Ecology and Management, ed. C. C. Trettin, M. F. Jurgensen, D. F. Grigal, M. R. Gale & J. K. Jeglum, pp. 3–18. Boca Raton, FL, USA: CRC Press.

Dahm, C. N. & Molles Jr, M. C. (1991) Streams in semi-arid regions as sensitive indicators of global climate change. In: *Global Climate Change and Freshwater Ecosystems*, ed.
P. Firth & S. G. Fisher, pp. 250–260. New York, USA: Springer-Verlag.

Dahm, C. N., Cleverly, J. R., Allred Coonroyd, J. E., et al. (2002) Evapotranspiration at the land/water interface in a semi-arid drainage basin. *Freshmater Biology* 47: 831–843.

Daily, G. C. & Ehrlich, P. (1996) Socioeconomic equity, sustainability, and Earth's carrying capacity. *Ecological Applications* 6: 991–1001.

Daka, E. R. & Hawkins, S. J. (2004) Tolerance to heavy metals in *Littorina saxatilis* from a metal-contaminated estuary in the Isle of Man. *Journal of the Marine Biological Association* of the United Kingdom 84: 393–400.

Daka, E. R., Allen, J. R. & Hawkins, S. J. (2003) Heavy metal contamination in sediment and biomonitors from sites around the Isle of Man. *Marine Pollution Bulletin* 46: 784–791.

Dale, P. E. R. (1994) An Australian perspective on coastal wetland management and vector control. In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 771–780. Amsterdam, the Netherlands: Elsevier.

Daly, H. E. (1991) Elements of environmental macroeconomics. In: *Ecological Economics: The Science and Management of Sustainability*, ed. R. Constanza, pp. 32–46. New York, USA: Columbia University Press.

Daly, M. A. & Mathieson, A. C. (1977) The effects of sand movement on intertidal seaweeds and selected invertebrates at Bound Rock, New Hampshire, USA. *Marine Biology* 43: 45–55.

Dando, P. R., Southward, A. J., Southward, E. C., *et al.* (1992) Shipwrecked tube worms. *Nature* **356**: 667.

Dangles, O. & Malmqvist, B. (2004) Species richness–decomposition relationships depend on species dominance. *Ecology Letters* 7: 395–402.

Danielopol, D. L., Griebler, C., Gunatilaka, A. & Notenboom, J. (2003) Present state and future prospects for groundwater ecosystems. *Environmental Conservation* **30**: 104–130. Danielopol, D. L., Pospisil, P., Dreher, J., et al. (2000)
A groundwater ecosystem in the wetlands of the Danube at Vienna (Austria). In: *Ecosystems of the World*, vol. 30, *Subterranean Ecosystems*, ed. H. Wilkens, D. C. Culver & W. F. Humphreys, pp. 581–602. Amsterdam, the Netherlands: Elsevier.

Darrigan, G. & Ezcurra de Drago, I. (2000) Limnosperma fortunei (Dunker, 1857) (Mytilidae) distribution in the Plata Basin. Medio Ambiente 13: 75–79.

Dasgupta, P. (2000) Population and resources: an exploration of reproductive and environmental externalities. *Population* and Development Review 26: 643–689.

Dasgupta, P. (2001) Human Well-Being and the Natural Environment. Oxford, UK: Oxford University Press.

da Silva, C. J., Wantzen, K. M., Nunes da Cunha, C. & Machado, F. de A. (2001) Biodiversity in the Pantanal Wetland, Brazil. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, vol. 2, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 187–215. Leiden, the Netherlands: Backhuys.

Daskalakis, K. D. & O'Connor, T. P. (1995) Distribution of chemical contaminants in US coastal and estuarine sediment. *Marine Environmental Research* 40: 381–398.

Dauer, D. M., Rodi Jr, A. J. & Ranasinghe, J. A. (1992) Effects of low dissolved oxygen events on the macrobenthos of the lower Chesapeake Bay. *Estuaries* 15: 384–391.

Daughton, C. G. & Ternes, T. A. (1999) Pharmaceuticals and personal care products in the environment: agents of subtle change? *Environmental Health Perspectives* 107: 907–938.

Dauterive, L. (2000) Rigs-to-Reefs Policy, Progress, and Perspective, Gulf of Mexico OCS Region Report. New Orleans, LA, USA: US Department of the Interior, Minerals Management Service.

D'Avanzo, C. D. & Kremer, J. N. (1994) Diel oxygen dynamics and anoxic events in an eutrophic estuary of Waquoit Bay, Massachusetts. *Estuaries* 17: 131–139.

Davidson, I., Vanderkam, R. & Padilla, M. (1999) Review of wetland inventory information in the Neotropics. In: *Global Review of Wetland Resources and Priorities for Wetland Inventory*, Supervising Scientist Report No. 144 / Wetlands International Publication No. 53, ed. C. M. Finlayson & A. G. Spiers, pp. 419–455. Canberra, ACT, Australia: Supervising Scientist.

Davis, J. (2000) Changes in a tidepool fish assemblage on two scales of environmental variation: seasonal and El Niño Southern Oscillation. *Limnology and Oceanography* 45: 1368–1379. Davis, R. & Short, F. T. (1997) Restoring eelgrass, *Zostera marina* L., habitat using a new transplanting technique: the horizontal rhizome method. *Aquatic Botany* 59: 1–15.

Day, D. (1990) Noah's Choice: True Stories of Extinction and Survival. London, UK: Penguin Books.

Day, E. & Branch, G. M. (2002) Effects of sea urchins (*Parechinus angulosus*) on recruits and juveniles of abalone (*Haliotis midae*). *Ecological Monographs* **72**: 133–149.

Day Jr, J. W., Hall, C. A. S., Kemp, W. M. & Yáñez-Arancibia, A., eds. (1989) *Estuarine Ecology*. New York, USA: John Wiley.

Dayton, P. K. (1975) Experimental studies of algal-canopy interactions in a sea otter dominated kelp community at Amchitka Island, Alaska. *Fishery Bulletin, United States* 73: 230–237.

Dayton, P. K. (1985a) Ecology of kelp communities. Annual Review of Ecology and Systematics 16: 215–245.

Dayton, P. K. (1985b) The structure and regulation of some South American kelp communities. *Ecological Monographs* 55: 447–468.

Dayton, P. K., Currie, V., Gerrodete, T., et al. (1984) Patch dynamics and stability of some California kelp communities. *Ecological Monographs* 54: 253–289.

Dayton, P. K., Tegner, M. J., Edwards, P. B. & Riser, K. L. (1998) Sliding baselines, ghosts, and reduced expectations in kelp forest communities. *Ecological Applications* 8: 309–322.

Dayton, P. K., Tegner, M. J., Edwards, P. B. & Riser, K. L. (1999) Temporal and spatial scales of kelp demography: the role of oceanography climate. *Ecological Monographs* 69: 219–250.

Dayton, P. K., Tegner, M. J., Parnell, P. E. & Edwards, P. B. (1992) Temporal and spatial patterns of disturbance and recovery in a kelp forest community. *Ecological Monographs* 62: 421–445.

Dayton, P. K., Thrush, S. F., Agardy, T. M. & Hofman, R. J. (1995) Environmental effects of fishing. *Aquatic Conservation: Marine and Freshwater Ecosystems* 5: 205–232.

Dean, J. M. (1996) A crisis and opportunity in coastal oceans: coastal fisheries as a case study. In: *Sustainable Development in the Southeastern Coastal Zone*, ed. F. J. Vernberg, W. B. Vernberg & T. Siewicki, pp. 81–88. Columbia, SC,

USA: University of South Carolina Press.

Dean, T. A. & Jewett, S. C. (2001) Habitat-specific recovery of shallow subtidal communities following the Exxon Valdez oil spill. *Ecological Applications* 11: 1456–1471. De Casabianca, M. L., Laugier, T. & Collart, D. (1997) Impact of shellfish farming eutrophication on benthic macrophyte communities in the Thau lagoon, France. *Aquaculture International* 5: 301–314.

Dederen, L. H. T. (1992) Marine eutrophication in Europe: similarities and regional differences in appearance. In: *Marine Coastal Eutrophication*, ed. R. A. Vollenweider, R. Marchetti & R. Viviani, pp. 673–682. Amsterdam, the Netherlands: Elsevier.

de Forges, R., Koslow, J. A. & Poore, G. C. B. (2000) Diversity and endemism of the benthic seamount fauna in the southwest Pacific. *Nature* 405: 944–947.

Degens, E. T. & Ross, D. A., eds. (1969) Hot Brines and Recent Heavy Metal Deposits in the Red Sea: A Geochemical and Geophysical Account. New York, USA: Springer-Verlag.

De Groot, A.J. (1995) Metals and sediments: a global perspective. In: *Metal Contaminated Aquatic Sediments*, ed.
H. E. Allen, pp. 1–20. Ann Arbor, MI, USA: Ann Arbor Press.

de Groot, R. S., Wilson, M. A. & Boumans, R. M. J. (2002) A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecological Economics* 41: 393–408.

Deichmann, U., Balk, D. & Yetman, G. (2001) Transforming population data for interdisciplinary usages: from census to grid [www document]. URL http://sedac.ciesin.columbia. edu/gpw/docs/gpw3_documentation_final.pdf

Deimling, E. A. & Liss, W. J. (1994) Fishery development in the eastern North Pacific: a natural–cultural system perspective, 1888–1976. *Fisheries Oceanography* 3: 60–77.

De Jonge, V. N. & De Jong, D. J. (1992) Role of tide, light and fisheries in the decline of *Zostera marina* L. in the Dutch Wadden Sea. *Netherlands Institute for Sea Research Publication Series* 20: 161–176.

De Jonge, V. N. (1990) Response of the Dutch Wadden Sea ecosystem to phosphorus discharges from the River Rhine. *Hydrobiologia* **195**: 49–62.

de la Mare, W. K. (1997) Abrupt mid-twentieth-century decline in Antarctic sea-ice extent from whaling records. *Nature* 389: 57–59.

DeLaune, R. D. & Pezeshki, S. R. (1994) The influence of subsidence and saltwater intrusion on coastal marsh stability: Louisiana Gulf coast, USA. *Journal of Coastal Research* Special Issue 12: 77–89.

Delgado, C. L., Wada, N., Rosegrant, M. W., Meijer, S. & Ahmed, M. (2003) *Fish to 2020: Supply and Demand in Changing Global Markets.* Manila, Philippines: International Food Policy Research Institute and World Fish Centre. Delgado, O. & Lapointe, B. E. (1994) Nutrient-limited productivity of calcareous versus fleshy macroalgae in a eutrophic, carbonate-rich tropical marine environment. *Coral Reefs* 13: 151–159.

Delgado, O., Ruiz, J., Perez, M., Romero, R. & Ballesteros, E. (1999) Effects of fish farming on seagrass in a Mediterranean Bay: seagrass decline after organic loading cessation. *Oceanologica Acta* 22: 109–117.

De Marsily, G. (1992) Creation of 'hydrological nature reserves', a plea for the defence of ground water. *Ground Water* **30**: 658.

Denevan, W. M., ed. (1966) *The Aboriginal Cultural Geography of the Llanos de Mojos of Bolivia*. Berkeley, CA, USA: University of California Press.

Denevan, W. M. (1976) The aboriginal population of Amazonia. In: *The Native Population of the Americas*, ed.
W. M. Denevan, pp. 205–234. Madison, WI, USA: University of Wisconsin Press.

den Hartog, C. (1970) *The Seagrasses of the World*. Amsterdam, the Netherlands: North Holland.

Denney, N. H., Jennings, S. & Reynolds, J. D. (2002) Life history correlates of maximum population growth rate in marine fishes. *Proceedings of the Royal Society of London B* 269: 2229–2237.

Dennis, K. C., Niang-Diop, I. & Nicholls, R. J. (1995) Sea-level rise and Senegal: potential impacts and consequences. *Journal of Coastal Research* Special Issue 14: 243–261.

Denny, M. & Paine, R. (1998) Celestial mechanics, sea-level changes, and intertidal ecology. *Biological Bulletin* 194: 108–115.

Denny, P. (1992) Africa. In: *Wetlands*, ed. M. Finlayson & M. Moser, pp. 115–148. Oxford, UK: Facts on File.

Denny, P. (1993) Wetlands of Africa. In: Wetlands of the World, vol. 1, Inventory, Ecology and Management,
ed. D. F. Whigham, D. Dykyjová & S. Hejný, pp. 1–128.
Dordrecht, the Netherlands: Kluwer.

Department of Land and Water Conservation (2002) *Compensatory Wetlands*, a discussion paper under the NSW Wetlands Management Policy. Sydney, NSW, Australia: Department of Land and Water Conservation.

Depledge, M. H. & Billinghurst, Z. (1998) Ecological significance of endocrine disruption in marine invertebrates. *Marine Pollution Bulletin* 39: 32–38.

Dermott, R. & Kerec, D. (1997) Changes in the deepwater benthos of eastern Lake Erie since the invasion of *Dreissena*: 1979–1993. *Canadian Journal of Fisheries and Aquatic Sciences* 54: 922–930. Derraik, J. G. B. (2002) The pollution of the marine environment by plastic debris: a review. *Marine Pollution Bulletin* 44: 842–852.

Desai, U., ed. (1998) Ecological Policy and Politics in Developing Countries: Economic Growth, Democracy, and Environment. Albany, NY, USA: State University of New York Press.

Deser, C., Walsh, J. E. & Timlin, M. S. (2000) Arctic sea ice variability in the context of recent atmospheric circulation trends. *Journal of Climate* 13: 617–633.

Desmond, J., Zedler, J. B. & Williams, G. D. (2000) Fish use of tidal creek habitats in two southern California salt marshes. *Ecological Engineering* 14: 233–252.

Detenbeck, N. E., Galatowitsch, S. M., Atkinson, J. & Ball, H. (1999) Evaluating perturbations and developing restoration strategies for inland wetlands in the Great Lakes Basin. *Wetlands* 19: 789–820.

Dette, H.-H. & Raudkivi, A. J. (1994) Beach nourishment and dune protection. In: *Coastal Engineering: Proceedings of the 24th Coastal Engineering Conference*, pp. 1934–1945. New York, USA: American Society of Civil Engineers.

Deuser, W. G. & Ross, E. H. (1980) Seasonal change in the flux of organic carbon to the deep Sargasso Sea. *Nature* 283: 364–365.

Devine, J. A., Baker, K. D. & Haedrich, R. L. (2006) Deep-sea fishes qualify as endangered. *Nature* 439: 29.

DFID/EC/UNDP/World Bank (2002) Linking Poverty Reduction and Environmental Management: Policy Challenges and Opportunities, World Bank Working Paper. Washington, DC, USA: The World Bank.

DHI Water and Environment (2001) Yellow River flood management sector project, China (2000).Unpublished report. Horsholm, Denmark: DHI Water and Environment.

Diaz, R. J. & Rosenberg, R. (1995) Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and Marine Biology Annual Review* 33: 245–303.

Dickhut, R. M. & Gustafson, K. E. (1995) Atmospheric inputs of selected polycyclic aromatic hydrocarbons and polychlorinated biphenyls to southern Chesapeake Bay. *Marine Pollution Bulletin* **30**: 385–396.

Dijkema, K. S. (1987) Geography of salt marshes in Europe. Zeitschrift für Geomorphologie 31: 489–499.

Dionne, J.-C. (1989) An estimate of shore ice action in a Spartina tidal marsh, St Lawrence estuary, Québec, Canada. Journal of Coastal Research 5: 281–293. Diop, E. S. (1990) La Côte Ouest-Africaine: du Saloum (Sénégal) à la Mellacorée (Rép. de Guinée), Collection Etudes et Thèses. Paris, France: ORSTOM.

Diop, E. S., Soumare, A., Diallo, N. & Guisse, A. (1997) Recent changes of the mangroves of the Saloum River estuary, Senegal. *Mangroves and Salt Marshes* 1: 163–172.

Dobson, A. P., Bradshaw, A. D. & Baker, A. J. M. (1997) Hopes for the future: restoration ecology and conservation biology. *Science* 277: 515–522.

Dobson, J. E., Bright, E. A., Ferguson, R. L., et al. (1995) NOAA Coastal Change Analysis Program (C-CAP): Guidance for Regional Implementation, NOAA Technical Report No. NFFS 123. Washington, DC, USA: US Department of Commerce.

Dodd, R. S. & Afzal-Rafii, Z. (2002) Evolutionary genetics of mangroves: continental drift to recent climate change. *Trees* 16: 80–86.

Dodd, R. S., Afzal-Rafii, Z., Kashani, N. & Budrick J. (2002) Land barriers and open oceans: effects on gene diversity and population structure in *Avicennia germinans* L. (Avicenniaceae). *Molecular Ecology* 11: 1327–1338.

Dodd, R. S., Blasco, F., Afzal-Rafii, Z. & Torquebiau, E. (1999) Mangroves of the United Arab Emirates: diversity at the bioclimatic extreme. *Aquatic Botany* 63: 291–304.

Donahue, D. L. (1999) The Western Range Revisited: Removing Livestock from Public Lands to Conserve Native Biodiversity. Norman, OK, USA: University of Oklahoma Press.

Done, T. (1992) Constancy and change in some Great Barrier Reef coral communities: 1980–1990. American Zoologist 32: 655–662.

Done, T. J. (1987) Simulation of the effects of *Acanthaster planci* on the population structure of massive corals in the genus *Porites*: evidence of population resilience? *Coral Reefs* 6: 75–90.

Donnelly, J. & Bertness, M. D. (2001) Rapid shoreward encroachment of salt marsh vegetation in response to sea-level rise. *Proceedings of the National Academy of Sciences* of the USA **98**: 14218–14223.

Doody, P. (1989) Conservation and development of the coastal dunes in Great Britain. In: *Perspectives in Coastal Dune Management*, ed. F. van der Meulen, P. D. Jungerius & J. H. Visser, pp. 53–67, The Hague, the Netherlands: SPB Academic Publishing.

Dosdat, A., Gaumet, F. & Chartois, H. (1995) Marine aquaculture effluent monitoring: methodological approach to the evaluation of nitrogen and phosphorus excretion by fish. *Aquacultural Engineering* 14: 59–84. Douglas, B. C. (1991) Global sea level rise. *Journal of Geophysical Research* 96: 6981–6992.

Douglas, B. C. & Peltier, W. C. (2002) The puzzle of global sea level rise. *Physics Today* March 2002: 35–40.

Douglas, M., Finlayson, C. M. & Storrs, M. J. (1998) Weed management in tropical wetlands of the Northern Territory, Australia. In: *Wetlands in a Dry Land: Understanding for Management*, ed. W. D. Williams, pp. 239–251. Canberra, ACT, Australia: Environment Australia.

Douglas-Westwood (2002) Into the deep [www document]. URL http://www.dw-1.com

Dove, S. (2004) Scleractinian coral with photoprotective host pigments are hypersensitive to thermal bleaching. *Marine Ecology Progress Series* 272: 99–116.

Dove, S., Hoegh-Guldberg, O. & Ranganathan, S. (2001) Major colour patterns in reef-building corals are due to a family of GFP-like proteins. *Coral Reefs* 19: 197–204.

Downing, J. A., McClain, M., Twilley, R., et al. (1999) The impact of accelerating land-use change on the N-cycle of tropical aquatic ecosystems: current conditions and projected changes. *Biogeochemistry* 46: 109–148.

Drake, B. G. (1992) A field study of the effects of elevated CO₂ on ecosystem processes in a Chesapeake Bay wetland. *Australian Journal of Botany* **40**: 579–595.

Drake, B. G., Muehe, G., Peresta, M. A., González-Meler & Matamala, R. (1996*a*) Acclimation of photosynthesis, respiration and ecosystem carbon flux of a wetland on Chesapeake Bay, Maryland to elevated atmospheric CO₂ concentration. *Plant and Soil* 187: 111–118.

Drake, B. G., Peresta, G., Beugeling, F. & Matamala, R. (1996b) Long-term elevated CO₂ exposure in a Chesapeake Bay wetland: ecosystem gas exchange, primary production, and tissue nitrogen. In: *Carbon Dioxide and Terrestrial Ecosystems*, ed. G. W. Koch & H. A. Mooney, pp. 197–214. New York, USA: Academic Press.

Drake, P., Baldó, F., Sáenz, V. & Arias, A. M. (1999) Macrobenthic community structure in estuarine pollution assessment on the Gulf of Cádiz (SW Spain): is the phylum-level meta-analysis approach applicable? *Marine Pollution Bulletin* 38: 1038–1047.

Dreher, J. & Gunatilaka, A. (2001) Management of urban groundwater. I. Quantitative aspects. In: Groundwater Ecology: A Tool For Management of Water Resources, ed. C. Griebler, D. L. Danielopol, J. Gibert, H. P. Nachtnebel & J. Notenboom, pp. 197–212. Luxembourg: Office for Official Publications of the European Communities. Dreyer, G. D. & Niering, W. A. (1995) Tidal marshes of Long Island Sound: ecology, history and restoration. *Connecticut College Arboretum Bulletin* 34: 1–72.

Driscoll, C. T., Lawrence, G. B., Bulger, A. J., *et al.* (2001) Acidic deposition in the Northeastern United States: sources and inputs, ecosystem effects, and management strategies. *BioScience* 51: 180–198.

Du Rietz, G. E. & Nannfeldt, J. A. (1925) Ryggmossen und Stigsbo Rödmosse: Die letzten lebenden Hochmoore der Gegend von Uppsala, Svenska Växtsociologiska Sällskapets Handlingar No. 3. Uppsala, Sweden: Amqvist & Wiksell.

Duarte, C. M. (1995) Submerged aquatic vegetation in relation to different nutrient regimes. *Ophelia* 41: 87–112.

Duarte, C. M. (2001) Seagrass ecosystems. In: *Encyclopedia* of *Biodiversity*, vol. 5, ed. S. L. Levin, pp. 255–268. San Diego, CA, USA: Academic Press.

Duarte, C. M. (2002) The future of seagrass meadows. *Environmental Conservation* **29**: 192–206.

Duarte, C. M. & Chiscano, C. L. (1999) Seagrass biomass and production: a reassessment. *Aquatic Botany* 65: 159–174.

Duarte, C. M., Martínez, R. & Barrón, C. (2002) Biomass, production and rhizome growth near the northern limit of seagrass (*Zostera marina*) distribution. *Aquatic Botany* 72: 183–189.

Duarte, C. M., Terrados, J., Agawin, N. S. W., et al. (1997) Response of a mixed Philippine seagrass meadow to experimental burial. *Marine Ecology Progress Series* 147: 285–294.

Dubinsky, Z. & Stambler, N. (1996) Marine pollution and coral reefs. *Global Change Biology* 2: 511–526.

Duda, A. M. & El-Ashry, M. T. (2000) Addressing the global water and environment crises through integrated approaches to the management of land, water and ecological resources. *Water International* 25: 115–126.

Dudgeon, D. (1999) Tropical Asian Streams: Zoobenthos, Ecology and Conservation. Hong Kong: Hong Kong University Press.

Dudgeon, D. (2000*a*) The ecology of tropical Asian rivers and streams in relation to biodiversity conservation. *Annual Review of Ecology and Systematics* **31**: 239–263.

Dudgeon, D. (2000b) Large-scale hydrological changes in tropical Asia: prospects for riverine biodiversity. *BioScience* 50: 793–806.

Dudgeon, D. (2002) An inventory of riverine biodiversity in monsoonal Asia: present status and conservation challenges. *Water Science and Technology* **11**: 11–19. Dudgeon, D., Arthington, A. H., Gessner, M. O., et al. (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews* 81: 163–182.

Dudley, N. (1990) *Nitrates: The Threat to Food and Water.* London, UK: Green Print.

Dugan, P. J., ed. (1990) Wetland Conservation: A Review of Current Issues and Required Action. Gland, Switzerland: IUCN.

Dugan, P. J. (1993) Wetlands in Danger. New York, USA: Oxford University Press.

Duggins, D. O. (1980) Kelp beds and sea otters: an experimental approach. *Ecology* **61**: 447–453.

Duggins, D. O. (1983) Starfish predation and the creation of mosaic patterns in a kelp dominated community. *Ecology* 64: 1610–1619.

Duggins, D. O., Eckman, J. E. & Sewell, A. T. (1990) Ecology of understory kelp environments. II. Effects of kelps on recruitment of benthic invertebrates. *Journal of Experimental Marine Biology and Ecology* 143: 27–45.

Duggins, D. O., Simenstad, C. A. & Estes, J. A. (1989) Magnification of secondary production by kelp detritus in coastal marine ecosystems. *Science* 245: 170–173.

Duke, N. C. (1995) Genetic diversity, distributional barriers and rafting continents: more thoughts on the evolution of mangroves. *Hydrobiologia* 295: 167–181.

Duke, N. C., Benzie, J. A. H., Goodall, J. A. & Ballment, E. R. (1998) Genetic structure and evolution of species in the mangrove genus *Avicennia* (Avicenniaceae) in the Indo-West Pacific. *Evolution* 52: 1612–1626.

Duke, N. C., Pinzón, Z. S. & Prada, M. C. (1997) Large-scale damage to mangrove forests following two large oil spills in Panama. *Biotropica* 29: 2–14.

Dulvy, N. K., Freckleton, R. P. & Polunin N. V. C. (2004) Coral reef cascades and the indirect effects of predator removal by exploitation. *Ecology Letters* 7: 410–416.

Dulvy, N. K., Metcalfe, J. D., Glanville, J., Pawson, M. G. & Reynolds, J. D. (2000) Fishery stability, local extinctions, and shifts in community structure in skates. *Conservation Biology* 14: 283–293.

Dulvy, N. K., Sadovy, Y. & Reynolds, J. D. (2003) Extinction vulnerability in marine populations. *Fish and Fisheries* 4: 25–64.

Dumont, H. (1995) Ecocide in the Caspian Sea. *Nature* 377: 673–674.

Dumortier, M., Verlinden, A., Beeckman, H. & van der Mijnsbrugge, K. (1996) Effects of harvesting dates and frequencies on above and below-ground dynamics in Belgian wet grasslands. *Ecoscience* 3: 190–198. Dunton, K. H. (1990) Growth and production in Laminaria solidungula: relation to continuous underwater light levels in the Alaskan High Arctic. Marine Biology 106: 297–304.

Dunton, K. H. & Dayton, P. K. (1995) The biology of high latitude kelp. In: *Ecology of Fjords and Coastal Waters: Proceedings of the Mare Nor Symposium on the Ecology of Fjords and Coastal Waters*, Tromsø, Norway, 5–9 December 1994, ed. H. R. Skjoldal, C. Hopkins, K. E. Erikstad & H. P. Leinass, pp. 499–507. Amsterdam, the Netherlands: Elsevier.

Dunton, K. H. & Schell, D. M. (1987) Dependence of consumers on macroalgal (*Laminaria solidungula*) carbon in an Arctic kelp community: delta-carbon-13 evidence. *Marine Biology* 93: 615–626.

Dunton, K. H. & Schonberg, S. V. (2002) Assessment of propeller scarring in seagrass beds of the South Texas Coast. *Journal of Coastal Research* Special Issue 37: 99–109.

Duplisea, D. E., Jennings, S., Malcolm, S. J., Parker, R. & Sivyer, D. (2001) Modelling the potential impacts of bottom trawl fisheries on soft sediment biochemistry of the North Sea. *Geochemical Transactions* 14: 1–6.

Dupont, L. M., Jahns, S., Marret, F. & Ning, S. (2000) Vegetation change in equatorial West Africa: time slices for the last 150 ka. *Paleogeography Paleoclimatology Paleoecology* 155: 95–122.

Durako, M. J. & Moffler, M. D. (1987) Factors affecting the reproductive ecology of *Thalassia testudinum* Hydrocharitacea. *Aquatic Botany* 27: 79–96.

Duran, L. R. & Castilla, J. C. (1989) Variation and persistence of the middle rocky intertidal community of central Chile, with and without human harvesting. *Marine Biology* **103**: 555–562.

Durgaprasad, M. K. & Anjaneyulu, Y., eds. (2003) Lake Kolleru: Environmental Status (Past and Present).
Hyderabad, Andhra Pradesh, India: B.S. Publications.

DWR (2003) Construction of the West Desert Pumping Project. Division of Water Resources, State of Utah [www document]. URL http://www.water.utah.gov/ construction/gsl/construction.htm.

Dynesius, M. & Nilsson, C. (1994) Fragmentation and flow regulation of river systems in the northern third of the world. *Science* 266: 753–762.

Eakin, C. M. (1996) Where have all the carbonates gone? A model comparison of calcium carbonate budgets before and after the 1982–1983 EI Niño at Uva Island in the eastern Pacific. *Coral Reefs* 15: 109–119. Ebeling, A. W., Laur, D. R. & Rowley, R. J. (1985) Severe storm disturbance and reversal of community structure in a southern California kelp forest. *Marine Biology* 84: 287–294.

Ebling, F. S., Hawkins, A. D., Kitching, J. A., Muntz, P. & Pratt, V. M. (1966) The ecology of Lough Ine. XVI. Predation and diurnal migrations in *Paracentrotus* community. *Journal of Animal Ecology* 35: 559–566.

EC (2002) Towards a Strategy to Protect and Conserve the Marine Environment, Communication from the Commission to the Council and the European Parliament. Brussels, Belgium: European Commission.

EC (2005) Directive of the European Parliament and of the Council Establishing a Framework for Community Action in the field of Marine Environmental Policy (Marine Strategy Directive), SEC (2005) 1290. Brussels, Belgium: European Commission.

Edelstein, T., Craigie, J. S. & McLachlan, J. (1969) Preliminary survey of the sublittoral flora of Halifax county. *Journal* of the Fisheries Research Board of Canada 26: 2703–2713.

Edgar, G. J. & Barrett, N. S. (2002) Effects of catchment activities on macrofaunal assemblages in Tasmanian estuaries. *Estuarine, Coastal and Shelf Science* **50**: 639–654.

Edinger, E. N. (1998) Reef degradation and coral biodiversity in Indonesia: effects of land-based pollution, destructive fishing practices and changes over time. *Marine Pollution Bulletin* **36**: 617–630.

Edmondson, W. T. (1991) *The Uses of Ecology: Lake Washington and Beyond.* Seattle, WA, USA: University of Washington Press.

Edmunds, P. J. & Carpenter, R. C. (2001) Recovery of Diadema antillarum reduces macroalgal cover and increases abundance of juvenile corals on a Caribbean reef. Proceedings of the National Academy of Sciences of the USA 98: 5067–5071.

Edwards, A. C. & Davis, D. E. (1974) Effects of herbicides on the *Spartina* salt marsh. In: *Ecology of Halophytes*, ed.
R. J. Reimold & W. H. Queen, pp. 531–545. New York, USA: Academic Press.

Edyvane, K. (2003) Conservation, Monitoring and Recovery of Threatened Giant Kelp (Macrocystis pyrifera) Beds in Tasmania, Report for Environment Australia. Hobart, Tasmania, Australia: Department of Primary Industries, Water and Environment. Available online at www.dpiwe. tas.gov.au/inter.nsf/Publications/HMUY-5TT2P6?open

Eganhouse, R. P. & Sherblom, P. M. (2001) Anthropogenic organic contaminants in the effluent of a combined sewer overflow: impact on Boston Harbor. *Marine Environmental Research* 51: 51–74. Ehrenfeld, J. G. (1990) Dynamics and processes of barrier island vegetation. *Aquatic Science* **2**: 437–480.

Eiane, K., Aksnes, D. L., Bagoien, E. & Kaartvedt, S. (1999) Fish or jellies: a question of visibility? *Limnology and Oceanography* 44: 1352–1357.

Eisenreich, S., Emmling, P. & Beeton, A. (1977) Atmospheric loading of phosphorus and other chemicals to Lake Michigan. *Journal of Great Lakes Research* 3: 291–304.

Eisler, R. (1987) Polycyclic Aromatic Hydrocarbon Hazards to Fish, Wildlife, and Invertebrates: A Synoptic Review, Biological Report No. 85(1.11). Washington, DC, USA: US Fish and Wildlife Service.

Eisler, R. (2000) Polycyclic aromatic hydrocarbons. In: *Handbook of Chemical Risk Assessment*, vol. 2, pp. 1343–1411. Boca Raton, FL, USA: Lewis Publishers.

Eisma, D. (1998) Intertidal Deposits: River Mouths, Tidal Flats, and Coastal Lagoons. Boca Raton, FL, USA: CRC Press.

Eliot, I., Finlayson, C. M. & Waterman, P. (1999) Predicted climate change, sea-level rise and wetland management in the Australian wet–dry tropics. *Wetlands Ecology and Management* 7: 63–81.

Ellenbroek, G. A., ed. (1987) *Ecology and Productivity of an African Wetland System: The Kafue Flats, Zambia.* The Hague, the Netherlands: Dr W. Junk.

Ellery, W. N. & McCarthy, T. S. (1998) Environmental change over two decades since dredging and excavation of the lower Boro River, Okavango Delta, Botswana. *Journal* of Biogeography 25: 361–378.

Ellingsen, K. E. (2002) Soft-sediment benthic biodiversity on the continental shelf in relation to environmental variability. *Marine Ecology Progress Series* 232: 15–27.

Elliott, W. R. (2000) Conservation of the North American cave and karst biota. In: *Ecosystems of the World*, vol. 30, Subterranean Ecosystems, ed. H. Wilkens, D. C. Culver & W. F. Humphreys, pp. 664–689. Amsterdam, the Netherlands: Elsevier.

Ellis, C. J. & Tallis, J. H. (2000) Climatic control of blanket mire development at Kentra Moss, north-west Scotland. *Journal of Ecology* 88: 869–889.

Ellis, D. V. & Pattisina, L. A. (1990) Widespread neogastropod imposex: a biological indicator of global TBT pollution. *Marine Pollution Bulletin* 21: 248–253.

Ellis, J., Cummings, V., Hewitt, J., Thrush, S. & Norkko, A. (2002) Determining effects of suspended sediment on condition of a suspension feeding bivalve (*Atrina zelandica*): results of a survey, a laboratory experiment and a field transplant experiment. *Journal of Experimental Marine Biology and Ecology* 267: 147–174. Ellis, L. M., Crawford, C. S. & Molles Jr, M. C. (2002) The role of the flood pulse in ecosystem-level process in southwestern riparian forests: a case study from the Middle Rio Grande. In: *Flood Pulsing and Wetlands: Restoring the Natural Balance*, ed. B. A. Middleton, pp. 51–108. New York, USA: John Wiley.

Ellis, R. (2003) *The Empty Ocean*. Washington, DC, USA: Island Press.

Ellison, A. M. & Farnsworth, E. J. (1996) Anthropogenic disturbance of Caribbean mangrove ecosystems: past impacts, present trends, and future predictions. *Biotropica* 28: 549–565.

Ellison, A. M. & Farnsworth, E. J. (2001) Mangrove communities. In: *Marine Community Ecology*, ed. M. D. Bertness, S. D. Gaines & M. E. Hay, pp. 423–442. Sunderland, MA, USA: Sinauer Associates.

Ellison, A. M., Farnsworth, E. J. & Merkt, R. E. (1999) Origins of mangrove ecosystems and the mangrove biodiversity anomaly. *Global Ecology and Biogeography Letters* 8: 95–115.

Ellison, J. C. & Stoddart, D. R. (1991) Mangrove ecosystem collapse during predicted sea-level rise: Holocene analogues and implications. *Journal of Coastal Research* 7: 151–165.

El-Raey, M., Nasr, S., Frihy, O., Desouki, S. & Dewidar, K. (1995) Potential impacts of accelerated sea-level rise on Alexandria Governate, Egypt. *Journal of Coastal Research* Special Issue 14: 190–204.

El-Sayed, S., Van Dijken, G. & Gonzalez-Rodas, G. (1996) Effects of ultraviolet radiation on marine ecosystems. *International Journal of Environmental Studies* 51: 199–216.

Elster, C., Perdomo, L. & Schnetter, M.-L. (1999) Impact of ecological factors on the regeneration of mangroves in the Cíenaga Grande de Santa Marta, Colombia. *Hydrobiologia* 413: 35–46.

EM-DAT (2004) The OFDA/CRED Emergency Disaster Database. Brussels, Belgium: Université Catholique de Louvain [www document]. URL http://www.em-dat.net/ disasters/list.php.

Emerson, S. (1985) Organic carbon preservation in marine sediments. In: *The Carbon Cycle and Atmospheric CO₂: Natural Variations Archean to Present*, ed. E. T. Sundquist & W. Broecker, pp. 711–717. Washington, DC, USA: American Geophysical Union.

Emmerson, M. C., Solan, M., Emes, C., Paterson, D. M. & Raffaelli, D. (2001) Consistent patterns and the idiosyncratic effects of biodiversity in marine ecosystems. *Nature* **411**: 73–77. Enfield, D. B. (1988) Is El Niño becoming more common? Oceanography 59: 123–127.

Engelman, R. & LeRoy, P. (1993) Sustaining Water: Population and the Future of Fresh Water Supplies. Washington, DC, USA: Population Action International.

Engelman, R., Cincotta, R. P., Dye, B., Gardner-Outlaw, T. & Wisnewski, J. (2000) *People in the Balance: Population and Natural Resources at the Turn of the Millennium*. Washington, DC, USA: Population Action International.

Environment Canada (2002) Risk management strategy for road salts [www document]. URL http://www.ec.gc.ca/ nopp/roadsalt/en/rms.htm

Epp, D. & Smoot, N. C. (1989) Distribution of seamounts in the North Atlantic. *Nature* 337: 254–257.

Erlandson, J. M. (2002) Anatomically modern humans, maritime voyaging, and the Pleistocene colonization of the Americas. In: *The First Americans: The Pleistocene Colonization of the New World*, ed. N. G. Jablonski, pp. 1–19. San Francisco, CA, USA: California Academy of Sciences.

Erlandson, J. M., Kennett, D. J., Ingram, B. L., *et al.* (1996) An archaeological and paleontological chronology for Daisy Cave (CA-SMI-261), San Miguel Island, California. *Radiocarbon* 38: 355–373.

Erlandson, J. M., Rick, T. C. & Vellanoweth, R. L. (2004) Human impacts on ancient environments: a case study from the Northern Channel Islands. In: *Voyages of Discovery: The Archaeology of Islands*, ed. S. M. Fitzpatrick, pp. 51–83. New York, USA: Praeger.

Esselink, J. (2000) Nature Management of Coastal Salt Marshes: Interactions between Anthropogenic Influences and Natural Dynamics. Haren, the Netherlands: Koeman en Bijkerk.

Esselink, P., Helder, G. J. F., Aerts, B. A. & Gerdes, K. (1997) The impact of grubbing by greylag geese (*Anser anser*) on the vegetation dynamics on a tidal marsh. *Aquatic Botany* 55: 261–279.

Esteban Morales, G. (2003) Methane hydrates in the Chilean continental margin. *Journal of Biotechnology* 6: 1–5.

Estes, J. A. & Duggins, D. O. (1995) Sea otters and kelp forests in Alaska: generality and variation in a community ecological paradigm. *Ecological Monographs* 65: 75–100.

Estes, J. A. & Steinberg, P. D. (1988) Predation, herbivory, and kelp evolution. *Paleobiology* 14: 19–36.

Estes, J. A., Duggins, D. O. & Rathbun, G. B. (1989) The ecology of extinctions in kelp forest communities. *Conservation Biology* 3: 252–264. Estes, J. A., Tinker, M. T., Williams, T. M. & Doak, D. F. (1998) Killer whale predation on sea otters linking oceanic and nearshore ecosystems. *Science* 282: 473–476.

EU (2001) Common strategy on the implementation of the Water Framework Directive [www document]. URL http://www.europa.eu.int/comm/environment/water/ water-framework

Evans, A. S., Webb, K. L. & Penhale, P. A. (1986) Photosynthetic temperature acclimation in two coexisting seagrasses *Zostera marina* and *Ruppia maritima*. *Aquatic Botany* 24: 185–198.

Evans, C. D., Cullen, J. M., Alewell, C., et al. (2001) Recovery from acidification in European surface waters. *Hydrology* and Earth System Sciences 5: 283–297.

Evans, N. T. & Short, F. T. (2005) Functional trajectory models for assessment of transplant development of seagrass, *Zostera marina* L., beds in the Great Bay Estuary, NH, USA. *Estuaries* 28: 936–947.

Evans, R. S. (1989) Saline water disposal options in the Murray Basin. BMR Journal of Australian Geology and Geophysics 11: 167–185.

Everard, M. (1997) Development of a British wetland strategy. *Aquatic Conservation: Marine and Freshwater Ecosystems* 7: 223–238.

Everson, I. (2002) Consideration of major issues in ecosystem monitoring and management. CCAMLR Science 9: 213–232.

Everson, I., ed. (2000) *Krill: Biology, Ecology and Fisheries.* Oxford, UK: Blackwell Science.

Ewel, K. C. & Odum, H. T., eds. (1984) *Cypress Swamps*. Gainesville, FL, USA: University of Florida Press.

Ewel, K. C., Twilley, R. R. & Ong, J. E. (1998) Different kinds of mangrove forests provide different goods and services. *Global Ecology and Biogeography Letters* 7: 83–94.

Fach, B. A., Hofmann, E. E. & Murphy, E. J. (2002) Modeling studies of Antarctic krill *Euphausia superba* survival during transport across the Scotia Sea. *Marine Ecology Progress Series* 231: 187–203.

Fagan, W. F., Lewis, M. A., Neubert, M. G. & van den Driessche, N. P. (2002) Invasion theory and biological control. *Ecology Letters* 5: 148–157.

Fagbami, A. A., Udo, E. J. & Odu, C. T. I. (1988) Vegetation damage in an oil field in the Niger delta of Nigeria. *Journal* of Tropical Ecology 4: 61–75.

Falkenmark, M. & Widstrand, C. (1992) Population and water resources: a delicate balance. *Population Bulletin* 47 (3): 1–36.

FAO (1990) *Fertilizer Yearbook*. New York, USA: United Nations.

FAO (1994) FAO Yearbook: Fishery Statistics, Catches and Landings 1992, vol. 74. Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (1996) *The State of World Fisheries and Aquaculture 1996*, Technical Report. Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (1998) Wetland Characterization and Classification for Sustainable Agricultural Development. Harare, Zimbabwe: Food and Agriculture Organization of the United Nations (FAO), Sub-Regional Office for East and Southern Africa (SAFR).

FAO (1999) Review of the State of World Fishery Resources: Inland Fisheries, FAO Fisheries Circular No. 942. Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (2001*a*) Asia-Pacific conference on early warning, prevention, preparedness, and management of disasters in food and agriculture. Unpublished report. Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (2001b) What Is the Code of Conduct for Responsible Fisheries? Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (2002) World Agriculture: Towards 2015/2030 – An FAO Perspective. Rome, Italy: Food and Agriculture Organization of the United Nations.

FAO (2003*a*) International Year of Rice, 2004. Rome, Italy: Food and Agriculture Organization of the United Nations. Available online at www.fao.org/rice2004/index_en.htm

FAO (2003b) Status and Trends in Mangrove Area Extent Worldwide, Forest Resources Assessment Working Paper No. 63, ed. M.L. Wilkie & S. Fortuna. Rome, Italy: Forest Resources Division, Food and Agriculture Organization of the United Nations.

 FAO (2004) The State of World Fisheries and Aquaculture 2004.
 Rome, Italy: Food and Agriculture Organization of the United Nations. Available online at www.fao.org

FAO (2006) The State of World Fisheries and Aquaculture 2006.Rome, Italy: Food and Agriculture Organization of the United Nations. Available online at www.fao.org

Farina, J. M. & Castilla, J. C. (2001) Temporal variation in the diversity and cover of sessile species in rocky intertidal communities affected by copper mine tailings in northern Chile. *Marine Pollution Bulletin* 42: 554–568.

Farman, J. C., Gardiner, B. G. & Shanklin, J. D. (1985) Large losses of total ozone in Antarctica reveal seasonal ClO_X/ NO_X interaction. *Nature* 315: 207–210.

Farnsworth, E. J., Ellison, A. M. & Gong, W. K. (1996) Elevated CO₂ alters anatomy, physiology, growth and reproduction of red mangrove (*Rhizophora mangle L.*). *Oecologia* 108: 599–609. Farrier, D. & Tucker, L. (2000) Wise use of wetlands under the Ramsar Convention: a challenge for meaningful implementation of international law. *Journal of Environmental Law* 12: 21–32.

Feldt, W. G., Kanisch, G., Kanisch, M. & Vobach, M. (1985) Radioecological studies of sites in the northeast Atlantic used for dumping low-level radioactive wastes: results of the research cruises of FRV *Walter Herwig. Archiv für Fischereiwissenschaft* 35: 91–195.

Fellows, M. Q. & Zedler, J. B. (1999) Effects of the non-native grass, *Parapholis incurva* (Poaceae) on the rare and endangered hemiparasite, *Cordylanthus maritimus* subsp. *maritimus* (Scrophulariaceae). *Madroño* 52: 91–98.

Felzer, B. & Heard, P. (1999) Precipitation differences amongst GCMs used for the US National Assessment. *Journal of American Water Works Association* 35: 1327–1339.

Feminella, J. W. (1996) Comparison of benthic macroinvertebrate assemblages in small streams along a gradient of flow permanence. *Journal of the North American Benthological Society* 15: 651–669.

Fenchel, T. & Blackburn, T. H. (1979) *Bacteria and Mineral Cycling*. London, UK: Academic Press.

Ferguson, N. P. & Lee, J. A. (1979) Some effects of bisulphite and sulphate on the growth of *Sphagnum* species in the field. *Environmental Pollution Series A* 21: 59–71.

Fernandes, M. B., Sicre, M.-A., Boireau, A. & Tronczynski, J. (1997) Polycyclic hydrocarbon (PAH) distributions in the Seine River and its estuary. *Marine Pollution Bulletin* 34: 857–862.

Ferreira, L. V. & Stohlgren, T. J. (1999) Effects of river level fluctuation on plant species richness, diversity, and distribution in a floodplain forest in Central Amazonia. *Oecologia* 120: 582–587.

Ferrier-Pagès, C., Gattuso, J.-P., Dallot, S. & Jaubert, J. (2000) Effect of nutrient enrichment on growth and photosynthesis of the zooxanthellate coral *Stylophora pistillata*. *Coral Reefs* 19: 103–113.

Field, C. D. (1995) Impact of expected climate change on mangroves. *Hydrobiologia* 295: 75–81.

Field, C. D., ed. (1996) Restoration of Mangrove Ecosystems. Okinawa, Japan: International Society of Mangrove Ecosystems.

Field, C. D. (1998) Rehabilitation of mangrove ecosystems: an overview. *Marine Pollution Bulletin* 37: 383–392.

Field, C. D. (1999) Mangrove rehabilitation: choice and necessity. *Hydrobiologia* 413: 47–52. Field, J. G., Griffiths, C. L., Griffiths, R. J., et al. (1980) Variation in structure and biomass of kelp communities along the south-west Cape coast. *Transactions of the Royal Society of South Africa* 44: 145–203.

Field, J. G., Griffiths, C. L., Linley, E.A, Carter R. A. & Zoutendyk, P. (1977) Upwelling in the nearshore marine ecosystem and its biological implications. *Estuarine and Coastal Marine Science* 11: 133–150.

Field, J. G., Hempel, G. & Summerhayes, C. P. (2002) Oceans 2020: Science, Trends, and the Challenge of Sustainability. Washington, DC, USA: Island Press.

Fielding, A. H. & Russell, G. (1976) The effect of copper on competition between marine algae. *Journal of Applied Ecology* 13: 871–876.

Fields, P., Graham, J., Rosenblatt, R. & Somero, G. (1993) Effects of expected global climate change on marine faunas. *Trends in Ecology and Evolution* 8: 361–367.

Figueras, M. J., Polo, F., Inza, I. & Guarro, J. (1997) Past, present and future perspectives of the EU bathing water directive. *Marine Pollution Bulletin* 34: 148–156.

Figueroa, M. E., Castillo, J. M., Redondo, S., et al. (2003) Facilitated invasion by hybridization of *Sarcocornia* species in a salt-marsh succession. *Journal of Ecology* 91: 616–626.

Findlay, R. H., Watling, L. & Mayer, L. M. (1995) Environmental impact of salmon net-pen culture on marine benthic communities in Maine: a case study. *Estuaries* 18: 145–179.

Finlayson, C. M. (1996) Information required for wetland management in the South Pacific. In: Wetland Conservation in the Pacific Islands Region: Proceedings of the Regional Workshop on Wetland Protection and Sustainable use in Oceania, Port Moresby, Papua New Guinea, 6–10 June 1994, ed. R. Jaensch, pp. 185–201. Canberra, ACT, Australia: Wetlands International–Asia/Pacific.

Finlayson, C. M. & Davidson, N. C. (1999) Global review of wetland resources and priorities for wetland inventory: summary report. Unpublished report. Wageningen, the Netherlands: Wetlands International and Jabiru.

Finlayson, C. M. & Eliot, I. (2001) Ecological assessment and monitoring of coastal wetlands in Australia's wet-dry tropics: a paradigm for elsewhere? *Coastal Management* 29: 105–115.

Finlayson, C. M. & Moser, M., eds. (1992) *Wetlands*. Oxford, UK: Facts on File.

Finlayson, C. M. & Rea, N. (1998) Reasons for the loss and degradation of Australian wetlands. *Wetlands Ecology and Management* 7: 1–11. Finlayson, C. M. & Spiers, A. G., eds. (1999) Global Review of Wetland Resources and Priorities for Wetland Inventory, Supervising Scientist Report No. 144/Wetlands International Publication No. 53. Canberra, ACT, Australia: Supervising Scientist.

Finlayson, C. M. & von Oertzen, I. (1993) Wetlands in Australia: Northern (tropical) Australia. In: Wetlands of the World, vol.1, Inventory, Ecology and Management, ed.
D. F. Whigham, D. Dykyjová & S. Hejný, pp. 195–243.
Dordrecht, the Netherlands: Kluwer.

Finlayson, C. M. & von Oertzen, I., eds. (1996) Landscape and Vegetation Ecology of the Kakadu Region, Northern Australia. Dordrecht, the Netherlands: Kluwer.

Finlayson, C. M., Begg, G. W., Howes, J., et al. (2002) A Manual for an Inventory of Asian Wetlands: Version 1.0. Kuala Lumpur, Malaysia: Wetlands International.

Finlayson, C. M., Davidson, N. C., Spiers, A. G. & Stevenson, N. J. (1999) Global wetland inventory: status and priorities. *Marine and Freshwater Research* 50: 717–727.

Finlayson, C. M., D'Cruz, R. & Davidson, N., Synthesis Team Co-Chairs (2005) *Ecosystems and Human Well-Being: Wetlands and Water Synthesis*, a report of the Millennium Ecosystem Assessment. Washington, DC, USA: World Resources Institute.

Finlayson, C. M., Hall, R. & Bayliss, B. (1998) Regional Review of Wetlands Management Issues: Wet–Dry Tropics of Northern Australia, LWRRDC Occasional Paper No. 03/97. Canberra, ACT, Australia: Supervising Scientist.

Firth, P. & Fisher, S. G., eds. (1992) *Global Climate Change and Freshwater Ecosystems.* Berlin, Germany: Springer-Verlag.

Fischer, G. & Heilig, G. K. (1997) Population momentum and the demand on land and water resources. *Philosophical Transactions of the Royal Society of London B* 352: 869–889.

Fish, C. (1960) Limnological Survey of Eastern and Central Lake Erie, 1928–29 US Fish and Wildlife Service Special Science Report, Fisheries No. 334. Washington, DC, USA: US Fish and Wildlife Service.

Fishelson, L. (1983) Population ecology and biology of *Dotilla sulcata* (Crustacea, Ocypodidae) on sandy beaches of the Red Sea. In: *Sandy Beaches as Ecosystems*, ed. A. McLachlan & T. Erasmus, pp. 643–654. The Hague, the Netherlands: Dr W. Junk.

Fisher, C. R., Urcuyo, I. A., Simpkins, M. A. & Nix, E. (1997) Life in the slow lane: growth and longevity of cold-seep vestimentiferans. *Marine Ecology – Publicazioni della Stazione Zoologica di Napoli* 18: 83–94. Fitt, W. K., Brown, B. E., Warner, M. E. & Dunne, R. P. (2001) Coral bleaching: interpretation of thermal tolerance limits and thermal thresholds in tropical corals. *Coral Reefs* 20: 51–65.

Fletcher, H. & Frid, C. (1996) Impact and management of visitor pressure on rocky intertidal algal communities. *Aquatic Conservation: Marine and Freshwater Ecosystems* 6: 287–297.

Flower, R. (1994) A review of current biological and recent environmental research on Lake Baikal from a British perspective. *Freshmater Forum* 4: 8–22.

Flower, R. (1998) Paleolimnology and recent environmental change in Lake Baikal: an introduction and overview of interrelated concurrent studies. *Journal of Paleolimnology* 20: 107–117.

Fojt, W. J. (1994) Dehydration and the threat to East Anglian fens, England. *Biological Conservation* **69**: 163–175.

Foley, J. A., Kutzbach, J. E., Coe, M. T. & Levis, S. (1994) Feedbacks between climate and boreal forests during the Holocene epoch. *Nature* 371: 52–54.

Folke, C. & Kaaberger, T., eds. (1991) Ecology, Economy and Environment, vol. 1, Linking the Natural Environment and the Economy: Essays from the Eco-Eco Group, Dordrecht, the Netherlands: Kluwer.

Folt, C. L., Chen, C. Y., Moore, M. V. & Burnaford, J. (1999) Synergism and antagonism among multiple stressors. *Limnology and Oceanography* 44: 864–877.

Fong, P. & Glynn, P. W. (1998) A dynamic size-structured population model: does disturbance control size structure of a population of the massive coral *Gardineroseris panulata* in the Eastern Pacific? *Marine Biology* **130**: 663–674.

Fong, P. & Glynn, P. W. (2000) A regional model to predict coral population dynamics in response to El Niño–Southern Oscillation. *Ecological Applications* 10: 842–854.

Fonseca, M. S., Kenworthy, W. J. & Thayer, G. W. (1998) Guidelines for the Conservation and Restoration of Seagrasses in the United States and Adjacent Waters, NOAA Coastal Ocean Program Decision Analysis Series No. 12. Silver Spring, MD, USA: NOAA Coastal Ocean Office.

Fonseca, M. S., Thayer, G. W. & Chester, A. J. (1984) Impact of scallop harvesting on eelgrass (*Zostera marina*) meadows: implications for management. *North American Journal of Fisheries Management* 4: 286–293.

Fortes, M. D. (1988) Mangrove and seagrasses of East Asia: habitats under stress. *Ambio* 17: 207–213.

Forum Biodiversität Schweiz (2004) *Biodiversität in der Schweiz: Zustand, Erhalt, Perspektiven.* Bern, Switzerland: Haupt-Verlag. Fosberg, F. R. (1971) Mangroves versus tidal waves. *Biological Conservation* 4: 38–39.

Fossa, J. H., Mortensen, P. B. & Furevik, D. M. (2002) The deep-water coral *Lophelia pertusa* in Norwegian waters: distribution and fishery impacts. *Hydrobiologia* 471: 1–12.

Foster, D. R., King, G. A., Glaser, P. H. & Wright, H. E. (1983) Origin of string patterns in boreal peatlands. *Nature* 306: 256–258.

Foster, G. & Carter, L. (1997) Mud sedimentation on the continental shelf at an accretionary margin: Poverty Bay, New Zealand. New Zealand Journal of Geology and Geophysics 40: 157–173.

Foster, I. D. L. & Charlesworth, S. M. (1996) Heavy metals in the hydrological cycle: trends and explanation. *Hydrological Processes* 10: 227–261.

Foster, M. & Schiel, D. R. (1985) The Ecology of Giant Kelp Forests in California: A Community Profile, Biological Report No. 85. Washington, DC, USA: US Fish and Wildlife Service.

Foster, M. S. (1990) Organization of macroalgal assemblages in the Northeast Pacific: the assumption of homogeneity and the illusion of generality. *Hydrobiologia* **192**: 21–33.

Foster, M. S., Tarpley, J. A. & Dern, S. L. (1990) To clean or not to clean: the rationale, methods, and consequences of removing oil from temperate shores. *Northwest Environmental Journal* 6: 105–120.

Fowler, A. M. & Hennessey, K. J. (1995) Potential impacts of global warming on the frequency and magnitude of heavy precipitation. *Natural Hazards* 11: 283–303.

Fox, J. J. (1998) Shoals and reefs in Australia–Indonesia relations: traditional Indonesian fishermen. In: *Australia in Asia: Episodes*, ed. A. Milner & M. Quilty, pp. 111–139. Melbourne, Victoria, Australia: Oxford University Press.

Francour, P., Ganteaume, A. & Poulain, M. (1999) Effects of boat anchoring in *Posidonia oceanica* seagrass beds in the Port-Cros National Park (north-western Mediterranean Sea). *Aquatic Conservation: Marine and Freshwater Ecosystems* 9: 391–400.

Frankl, R. & Schmeidl, H. (2000) Vegetation change in a south German raised bog: ecosystem engineering by plant species, vegetation switch or ecosystem level feedback mechanisms? *Flora* 195: 265–276.

Fraser, W. R., Trivelpiece, W. Z., Ainley, G. G. & Trivelpiece, S. G. (1992) Increases in Antarctic penguin populations: reduced competition with whales or loss of sea ice due to environmental warming. *Polar Biology* 11: 525–531. Freestone, A. L. & Nordstrom, K. F. (2001) Early development of vegetation in restored dune plant microhabitats on a nourished beach at Ocean City, New Jersey. *Journal of Coastal Conservation* 7: 105–116.

Frei, C., Schär, C., Lüthi, D. & Davies, H. C. (1998) Heavy precipitation processes in a warmer climate. *Geophysical Research Letters* 25: 1431–1434.

Fremlin, G., ed. (1974) The National Atlas of Canada, 4th edn. Toronto, Ontario, Canada: Macmillan.

French, G. T., Awisoka, L. F. & Ibe, C. E. (1995) Sea-level rise and Nigeria: potential impacts and consequences. *Journal of Coastal Research* Special Issue 14: 224–242.

Frenken, K., & Mharapara, I., eds. (2002) Wetland Development and Management in SADC Countries, Proceedings of a sub-regional workshop, 19–23 November 2001. Harare, Zimbabwe: Sub-regional Office for Southern and East Africa of the Food and Agriculture Organization of the United Nations.

Friday, A. & Ingram, D. S., eds. (1985) The Cambridge Encyclopedia of Life Sciences. Cambridge, UK: Cambridge University Press.

Friedlander, A. M. & Parrish, J. D. (1998) Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *Journal of Experimental Marine Biology and Ecology* 224: 1–30.

Friedman, J. M. & Auble, G. T. (2000) Floods, flood control, and bottomland vegetation. In: *Inland Flood Hazards: Human, Riparian, and Aquatic Communities*, ed. E. Wohl, pp. 219–237. Cambridge, UK: Cambridge University Press.

Friedrich, R. & Reis, S. (2000) *Tropospheric Ozone Abatement*. Berlin, Germany: Springer-Verlag.

Frissell, C. A., Liss, W. J., Warren, C. E. & Hurley, M. D. (1986) A hierarchical framework for stream habitat classification: viewing streams in a watershed context. *Environmental Management* 10: 199–214.

Froese, R. & Binohlan, C. (2000) Empirical relationships to estimate asymptotic length, length at first maturity, and length at maximum yield per recruit in fishes, with a simple method to evaluate length–frequency data. *Journal of Fish Biology* 56: 758–773.

Fryer, G. & Iles, T. D. (1972) The Cichlid Fishes of the Great Lakes of Africa. Edinburgh, UK: Oliver & Boyd.

Fujimoto, K., Miyagi, T., Kikuchi, T. & Kawana, T. (1996) Mangrove habitat formation and response to Holocene sea-level changes on Kosrae Island, Micronesia. *Mangroves* and Salt Marshes 1: 47–57.

Fujita, D. (1987) The report of interviews with fishermen on Isoyake in Taisei-cho, Hokkaido. Suisanzoshoku 35: 135–138. Fujita, D. (1998) Strongylocentrotid sea urchin-dominated barren grounds on the Sea of Japan coast of northern Japan.
In: *Echinoderms*, ed. R. Mooi & M. Telford, pp. 659–664.
Rotterdam, the Netherlands: A.A. Balkema.

Fukami, T. & Morin, P. J. (2003) Productivity–biodiversity relationships depend on the history of community assembly. *Nature* 424: 423–426.

Funnell, B. F. & Boomer, I. (1998) Microbiofacies, tidal level and age deduction in Holocene saltmarsh on the North Norfolk Coast. *Bulletin of the Geological Society of Norfolk* 46: 31–55.

Funnell, B. F. & Pearson, I. (1989) Holocene sedimentation on the North Norfolk barrier coast in relation to relative sea-level. *Journal of Quaternary Science* 4: 25–36.

Gacia, E. & Duarte, C. M. (2001) Elucidating sediment retention by seagrasses: sediment deposition and resuspension in a Mediterranean (*Posidonia oceanica*) meadow. *Estuarine*, *Coastal and Shelf Science* 52: 505–514.

Gacia, E., Duarte, C. M., Marbà, N., et al. (2003) Sediment deposition and production in SE-Asia seagrass meadows. Estuarine, Coastal and Shelf Science 56: 909–919.

Gaden, M. (1998) Phosphorus targets achieved in Lake Erie [www.document]. URL http://www.enviro-mich@ great-lakes.net

Gage, J. D. & Tyler, P. A. (1991) *Deep-Sea Biology: A Natural History of Organisms at the Deep-Sea Floor.* Cambridge, UK: Cambridge University Press.

Gagne, J. A., Mann, K. H. & Chapman, A. R. O. (1982) Seasonal patterns of growth and storage in *Laminaria longicruris* in relation to differing patterns of availability of nitrogen in the water. *Marine Biology* 69: 91–101.

Gaines, S. D. & Roughgarden, J. (1985) Larval settlement rate, a leading determinant of structure in sessile marine species. *Nature* **360**: 579–580.

Gaines, S. D. & Roughgarden, J. (1987) Fish in offshore kelp forests affect recruitment to intertidal barnacle populations. *Science* 235: 479–481.

Galatowitsch, S. M., Anderson, N. O. & Ascher, P. D. (1999) Invasiveness in wetland plants in temperate North America. *Wetlands* 19: 733–755.

Gales, R. (1997) Albatross populations: status and threats. In: *Albatross Biology and Conservation*, ed. G. Robertson &
R. Gales, pp. 20–45. Chipping Norton, NSW, Australia: Surrey Beatty.

Galloway, J. (1995) Acid deposition: perspectives in time and space. *Water, Air and Soil Pollution* 85: 15–24.

Galloway, J. N., Levy II, H. & Kasibhatia, P. S. (1994) Year 2020: consequences of population growth and development on deposition of oxidized nitrogen. *Ambio* 23: 123–132.

Gan, B. K. (1995) *A Working Plan for the Matang Mangrove Forest Reserve, Perak*, 4th revision. Ipoh, Malaysia: State Forestry Department of Perak Darul Ridzuan.

Garcia, S. & Grainger, R. (1996) Fisheries management and sustainability: a new perspective on an old problem. In: Developing and Sustaining World Fisheries Resources: The State of Science and Management, 2nd World Fisheries Congress Proceedings, ed. D. A. Hancock, D. C. Smith, A. Grant & J. P. Beumer, pp. 631–645. Collingwood, Victoria, Australia: CSIRO.

Garcia-Charton, J. A. & Pérez-Ruzafa, Á. (2001) Spatial pattern and habitat structure of a Mediterranean rocky reef fish local assemblage. *Marine Biology* **138**: 917–934.

Gardner, T. A., Cote, I. M., Gill, J. A., Grant, A. & Watkinson, A. R. (2003) Long-term region-wide declines in Caribbean corals. *Science* **301**: 958–960.

Garg, J. K., Singh, T. S. & Murthy, T. V. R., eds. (1998) *Wetlands of India*. Ahmedabad, Gujurat, India: Space Applications Centre (ISRO).

Gattuso, J. P., Frankignoulle, M. & Wollast, R. (1998) Carbon and carbonate metabolism in coastal aquatic ecosystems. *Annual Review of Ecology and Systematics* 29, 405–434.

Gawler, M., ed. (2002) Strategies for Wise Use of Wetlands: Best Practices in Participatory Management, proceedings of a workshop held at the 2nd International Conference on Wetlands and Development, November 1998, Dakar, Senegal, WWF Publication No. 56. Wageningen, the Netherlands: Wetlands International and IUCN.

Geddes, M. C. & Williams, W. D. (1987) Comments on Artemia introductions and the need for conservation. In: Artemia Research and its Applications, vol. 3, ed. P. Sorgeloos, D. A. Bengston, W. Decleir & E. Jaspers, pp. 19–26.
Wetteren, Belgium: Universa Press.

Geelen, L. H. W., Cousin, E. F. H. & Schoon, C. F. (1995) Regeneration of dune slacks in the Amsterdam Waterworks dunes. In: *Directions in European Coastal Management*, ed. M. G. Healy & P. J. Doody, pp. 525–532. Cardigan, UK: Samara Publishing.

George, D. G. (2002) Regional-scale influences on the long-term dynamics of lake plankton. In: *Phytoplankton Production*, ed. P. J. Le B. Williams, D. N. Thomas & C. S. Reynolds, pp. 265–290. Oxford, UK: Blackwell Scientific Publications. George, D. G. & Taylor, A. H. (1995) UK lake plankton and the Gulf Stream. *Nature* **378**: 139.

George, D. G., Talling, J. F. & Rigg, E. (2000) Factors influencing the temporal coherence of five lakes in the English Lake District. *Freshwater Biology* 43: 449–461.

Gerard, V. A. (1982) Growth and utilization of internal nitrogen reserves by the giant kelp *Macrocystis pyrifera* in a low nitrogen environment. *Marine Biology* **66**: 27–35.

Gerard, V. A. (1997) The role of nitrogen nutrition in high-temperature tolerance of kelp, *Laminaria saccharina*. *Journal of Phycology* 33: 800–810.

GESAMP (1982) Scientific Criteria for the Selection of Waste Disposal Sites at Sea, Reports and Studies No. 16, Joint Group of Experts in the Scientific Aspects of Marine Environmental Protection. London, UK: Inter-Governmental Maritime Consultative Organization.

GESAMP (1990) *The State of the Marine Environment*, Reports and Studies No. 39, Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection. London, UK: IMP.

GESAMP (2001) Planning and Management for Sustainable Coastal Aquaculture Development, Report and Studies No. 68, Joint Group of Experts in the Scientific Aspects of Marine Environmental Protection. Rome, Italy: Food and Agriculture Organization of the United Nations.

Getter, C. D., Ballou, T. G. & Koons, C. B. (1985) Effects of dispersed oil on mangroves, synthesis of a seven year study. *Marine Pollution Bulletin* 16: 318–324.

Ghassemi, F., Jakeman, A. J. & Nix, H. A., eds. (1995) Salinization of Land and Water Resources: Human Causes, Extent, Management and Case Studies. Sydney, NSW, Australia: University of New South Wales Press.

Ghosh, A. K. & Mukhopadhyay, R. (2000) *Mineral Wealth of the Ocean*. Rotterdam, the Netherlands: A.A. Balkema.

Gibbs, P. E. & Bryan, G. W. (1987) TBT paints and the demise of the dogwhelk, *Nucella lapillus* (Gastropoda). In: *Oceans '87, Proceedings of the International Organotin Symposium*, vol. 4, pp. 1482–1487. New York, USA: Institute of Electrical and Electronic Engineers.

Gibert, J. (2001a) Basic attributes of groundwater
ecosystems. In: *Groundwater Ecology: A Tool for Management* of Water Resources, ed. C. Griebler, D. L. Danielopol,
J. Gibert, H. P. Nachtnebel & J. Notenboom, pp. 39–54.
Luxembourg: Office for Official Publications of the European Communities.

Ghiorse, W. C. (1997) Subterranean life. *Science* 275: 789–790.

Gibert, J. (2001b) Protocols for the assessment and conservation of aquatic life in the subsurface (PASCALIS): a European project. In: *Mapping Subterranean Biodiversity*, Special Publications No. 6, ed. D. C. Culver, L. Deharveng, J. Gibert & I. D. Sasowsky, pp. 19–21. Charles Town, WV, USA: Karst Waters Institute.

Gibert, J., Danielopol, D. L. & Stanford, J. A., eds. (1994) *Groundwater Ecology*. San Diego, CA, USA: Academic Press.

Gibert, J., Malard, F., Turquin, M.-J. & Laurent, R. (2000)
Karst ecosystems in the Rhône River basin. In: *Ecosystems of* the World, vol. 30 Subterranean Ecosystems, ed. H. Wilkens, D. C. Culver & W. F. Humphreys, pp. 533–558.
Amsterdam, the Netherlands: Elsevier.

Gibson, D. J. & Looney, P. B. (1994) Vegetation colonization of dredge spoil on Perdido Key, Florida. *Journal of Coastal Research* 10: 133–143.

Gibson, D. J., Ely, J. S. & Looney, P. B. (1997) A Markovian approach to modeling succession on a coastal barrier island following beach nourishment. *Journal of Coastal Research* 13: 831–841.

Gichuki, C. M. (2000) Community participation in the protection of Kenya's wetlands. *Ostrich* 71: 122–125.

Gido, K. B. & Brown, J. H. (1999) Invasion of North American drainages by alien fish species. *Freshwater Biology* 42: 387–399.

Giesen, W., Baltzer, M. & Baruadi, R., eds. (1991) Integrating Conservation with Land-Use Development in Wetlands of South Sulawesi. Bogor, Indonesia: PHPA and Asian Wetland Bureau.

Gilding, P., Hogarth, M. & Reed, D. (2002) Single bottom line sustainability: how a value centred approach to corporate sustainability can pay off for shareholders and society [www document]. ECOS Corporation, Sydney, Australia: 28 pp. URL http://www.ecoscorporation.com/ think/essen/SingleBottomLine.htm

Gill, A. M., Ryan, P. G., Moore, P. H. R. & Gibson, M. (2000) Fire regimes of World Heritage National Park, Australia. *Austral Ecology* 25: 616–625.

Ginsburg, R. N. C. (1994) Proceedings of the Colloquium on Global Aspects of Coral Reefs: Health, Hazards and History. Miami, FL, USA: Rosenstiel School of Marine and Atmospheric Science, University of Miami.

Gitay, H., Brown, S., Easterling, W. & Jallow, B. (2001) Ecosystems and their goods and services. In: *Climate Change 2001: Impacts, Adaptations, and Vulnerability – Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, ed. J. J. McCarthy, O. F. Canziani, N. A. Leary, D. J. Dokken & K. S. White, pp. 235–342. Cambridge, UK: Cambridge University Press.

Gitay, H., Suárez, A., Watson, R. T. & Dokken, D. J. (2002) *Climate Change and Biodiversity*, IPCC Technical Paper No. 92-9169-104-7. Geneva, Switzerland: Intergovernmental Panel on Climate Change.

Glasby, G. P. (2000) Lessons learned from deep-sea mining. *Science* 289: 551–553.

Glasby, T. & Connell, S. (1999) Urban structures as marine habitats. *Ambio* 28: 595–598.

Glaser, P. H. & Janssens, J. A. (1986) Raised bogs in eastern North America: transitions in landforms and stratigraphy. *Canadian Journal of Botany* 64: 395–415.

Glaser, P. H., Hansen, B. C. S., Siegel, D. I., Reeve, A. S. & Morin, P. J. (2004a) Rates, pathways and drivers for peatland development in the Hudson Bay Lowlands, northern Ontario, Canada. *Journal of Ecology* 92: 1036–1053.

Glaser, P. H., Siegel, D. I., Reeve, A. S., Janssens, J. A. & Janecky, D. R. (2004b) Tectonic drivers for vegetation patterning and landscape evolution in the Albany River region of the Hudson Bay Lowlands. *Journal of Ecology* 92: 1054–1070.

Gleick, P. H., ed. (1993a) Water in Crisis: A Guide to the World's Fresh Water Resources. New York, USA: Oxford University Press.

Gleick, P. H. (1993b) Water in the 21st century. In: Water in Crisis: A Guide to the World's Fresh Water Resources, ed.
P. H. Gleick, pp. 105–116. New York, USA: Oxford University Press.

Gleick, P. H. (1996) Basic water requirements for human activities: meeting basic needs. *Water International* 21: 83–122.

Gleick, P. H. (1999) The human right to water. *Water Policy* 1: 487–503.

Gleick, P., Cohen, M. & Mann, A. S. (2002) The World's Water 2002–2003: The Biennial Report on Freshwater Resources. Washington, DC, USA: Island Press.

Gleick, P. H., Singh, A. & Shi, H. (2001) Threats to the World's Freshwater Resources. Oakland, CA, USA: Pacific Institute; and Nairobi, Kenya: United Nations Environmental Programme.

Glenn, E. P., Brown, J. B. & O'Leary, J. W. (1998) Irrigating crops with seawater. *Scientific American* 279: 76–81.

Glenn, E. P., O'Leary, J. W., Watson, M. C., Thompson, T. L. & Kuehl, R. O. (1991) Salicornia bigelovii Torr.: an oilseed halophyte for seawater irrigation. Science 251: 1065–1067. Gliwicz, Z. M. (2002) On the different nature of top-down and bottom-up effects in pelagic food webs. *Freshwater Biology* 47: 2296–2312.

GLOBEC (2004) Global Ocean Ecosystem Dynamics [www document]. URL http://www.globec.org

Gloersen, P. & Campbell, W. J. (1988) Variations in the Arctic, Antarctic and global sea ice covers during 1978–1987 as observed with the Nimbus 7 scanning multichannel microwave radiometer. *Journal of Geophysical Research* 93: 10666–10674.

Gloersen, P., Campbell, W. J., Cavalieri, D. J., et al. (1992) Arctic and Antarctic Sea Ice, 1978–1987: Satellite Passive-Microwave Observations and analysis, NASA Special Publication No. 511. Pasadena, CA: NASA.

Gloersen, P., Parkinson, C. L., Cavalieri, D. J., Comiso, J. C.
& Zwally, H. J. (1999) Spatial distribution of trends and seasonality in the hemispheric sea ice covers: 1978–1996. *Journal of Geophysical Research* 104: 20 827–20 835.

Glover, A. G. & Smith, C. R. (2003) The deep-sea floor ecosystem: current status and prospects for anthropogenic change by the year 2025. *Environmental Conservation* 30: 219–242.

Glover, A. G., Paterson, G., Bett, B., et al. (2001) Patterns in polychaete abundance and diversity from the Madeira Abyssal Plain, northeast Atlantic. *Deep-Sea Research I* 48: 217–236.

Glover, A. G., Smith, C. R., Paterson, G. L. J., et al. (2002) Polychaete species diversity in the central Pacific abyss: local and regional patterns, and relationships with productivity. *Marine Ecology Progress Series* 240: 157–170.

Glynn, P. W. (1994) State of coral reefs in the Galapagos Islands: natural vs. anthropogenic impacts. *Marine Pollution Bulletin* 29: 131–140.

Glynn, P. W. (2000) El Niño–Southern Oscillation mass mortalities of reef corals: a model of high temperature marine extinctions? In: *Carbonate Platform Systems: Components and Interactions*, ed. E. Insalaco, P. W. Skelton & T. J. Palmer, pp. 117–133. London, UK: Geological Society of London.

Glynn, P. W. & Ault, J. S. (2000) A biogeographic analysis and review of the far eastern Pacific coral reef region. *Coral Reefs* 19: 1–23.

Glynn, P. W., Mate, J. L., Baker, A. C. & Calderon, M. O. (2001) Coral bleaching and mortality in Panama and Ecuador during the 1997–1998 El Niño–Southern Oscillation event: spatial/temporal patterns and comparisons with the 1982–1983 event. *Bulletin of Marine Science* 69: 79–109. Godcharles, M. F. (1971) A Study of the Effects of a Commercial Hydraulic Clam Dredge on Benthic Communities in Estuarine Areas, Technical Series No. 64. Tallahassee, FL, USA: Florida Department of Natural Resources.

Godwin, H. (1981) The Archives of the Peat Bog. Cambridge, UK: Cambridge University Press.

Godwin, K. S., Shallenberger, J., Leopold, D. J. & Bedford,
B. L. (2002) Linking landscape properties to local hydrogeologic gradients and plant species occurrence in New York fens: a hydrogeologic setting (HGS) framework. *Wetlands* 22: 722–737.

Gold, T. (1999) *The Deep Hot Biosphere*. New York, USA: Springer-Verlag.

Goldberg, E. D. (1994) Diamonds and plastics are forever? Marine Pollution Bulletin 25: 1–4.

Goldberg, E. D. (1995) Emerging problems in the coastal zone for the twenty-first century. *Marine Pollution Bulletin* 31: 152–158.

Goldberg, E. D. (1998) Marine pollution: an alternative view. Marine Pollution Bulletin 36: 112–113.

Goldschmidt, T. (1996) Darwin's Dreampond. Cambridge, MA, USA: MIT Press.

Gong, W. K., Ong, J. E. & Wong, C. H. (1984) Demographic studies on *Rhizophora apiculata* Bl. in the Matang Mangrove Forest, Malaysia. In: *Ecological Effects of Increasing Human Activities on Tropical and Subtropical Forest Ecosystems*, pp. 64–67. Technical Report of the Malaysian National MAB Committee. Penang, Malaysia: Man and the Biosphere.

Goodall, D. W., ed. (1977–2000) Ecosystems of the World, vol. 1–30. Amsterdam, the Netherlands: Elsevier.

Gooday, A. J. (2002) Biological responses to seasonally varying fluxes of organic matter at the ocean floor: a review. *Journal of Oceanography* 58: 305–332.

Goodland, R. (1991) The environment as capital. In: Ecological Economics: The Science and Management of Sustainability, ed. R. Costanza, pp. 168–175. New York, USA: Columbia University Press.

Goolsby, D. A., Battalin, W. A., Lawrence, G. B., et al. (1999) Flux and Sources of Nutrients in the Mississippi-Atchafalaya

González Bernáldez, F., Pérez Pérez, C. & Sterling Carmona, A. (1985) Areas of evaporative discharge from aquifers: little known Spanish ecosystems deserving protection. *Journal of Environmental Management* 21: 321–330.

González, E. (1991) El manglar de la Cíenaga Grande de Santa Marta, ecosistema en peligro de extinción: Colombia, sus gentes y regiones. *Instituto Geográfico 'Agustín Codazzi'* 21: 2–21.

River Basin, Topic 3, Report for the Integrated Assessment on Hypoxia in the Gulf of Mexico, NOAA Coastal Ocean Program Decision Analysis Series No. 17. Silver Spring, MD, USA: National Oceanic and Atmospheric Administration.

Gopal, B. (1987) *Water Hyacinth*. Amsterdam, the Netherlands: Elsevier.

Gopal, B. (1991) Wetland (mis)management by keeping people out: two examples from India. *Landscape and Urban Planning* 20: 53–59.

Gopal, B. (1999) Natural and constructed wetlands for wastewater treatment: potentials and problems. *Water Science and Technology* 40(3): 27–35.

Gopal, B. (2002) Management issues of aquatic resources including wetlands. In: *Institutionalizing Common Pool Resources*, ed. D. K. Marothia, pp. 674–688. New Delhi, India: Concept Publishing.

Gopal, B. (2003) Aquatic biodiversity in arid and semi-arid zones of Asia and water management. In: *Conserving Biodiversity in Arid Regions: Best Practices in Developing Nations*, ed. J. Lemons, R. Victor and D. Schaeffer, pp. 199–215. Dordrecht, the Netherlands: Kluwer.

Gopal, B. (2005) Does inland aquatic biodiversity have a future in Asian developing countries? *Hydrobiologia* 542: 69–75.

Gopal, B. & Chauhan, M. (2001) South Asian wetlands and their biodiversity: the role of monsoons. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, vol. 2, ed.
B. Gopal, W. J. Junk & J. A. Davis, pp. 257–276. Leiden, the Netherlands: Backhuys.

Gopal, B. & Junk, W.J. (2001) Assessment, determinants, function and conservation of biodiversity in wetlands: present status and future needs. In: *Biodiversity in Wetlands:* Assessment, Function and Conservation, vol. 2, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 277–302. Leiden, the Netherlands: Backhuys.

Gopal, B. & Krishnamurthy, K. (1993) Wetlands of South Asia. In: Wetlands of the World, vol. 1, Inventory, Ecology and Management, ed. D. F. Whigham, D. Dykyjová & S. Heiný, pp. 345–414. Dordrecht, the Netherlands: Kluwer.

Gopal, B., Junk, W. J. & Davis, J. A., eds. (2000) Biodiversity in Wetlands: Assessment, Function and Conservation, vol. 1. Leiden, the Netherlands: Backhuys.

Gopal, B., Junk, W. J. & Davis, J. A., eds. (2001) Biodiversity in Wetlands: Assessment, Function and Conservation, vol. 2. Leiden, the Netherlands: Backhuys.

Gopal, B., Kvet, J., Löffler, H., Masing, V. & Patten, B. C. (1990) Definition and classification. In: *Wetlands and* Shallow Continental Water Bodies. vol. 1, Natural and Human Relationships, ed. B. C. Patten *et al.*, pp. 9–16. The Hague, the Netherlands: SPB Academic Publishing.

Gopal, B., Zutshi, D. P. & Van Duzer, C. (2003) Floating islands in India: control or conserve? *International Journal of Ecology and Environmental Sciences* 29: 157–169.

Gordon, C. (1992) Sacred groves and conservation in Ghana. Newsletter of the IUCN SSC African Reptile and Amphibian Specialist Group 1: 3–4.

Gordon, C. (2003) Aquatic resource management and freshwater ecosystems of West Africa. In: Conservation, Ecology and Management of African Freshwaters, ed. T. L. Crisman, L. J. Chapman, C. A. Chapman & L. S. Kaufman, pp. 62–78. Gainesville, FL, USA: University of Florida Press.

Gore, A. J. P., ed. (1983) *Ecosystems of the World*, vol. 4, *Mires*, *Swamp*, *Bog*, *Fen and Moor*. Amsterdam, the Netherlands: Elsevier.

Goreau, T., McClanahan, T., Hayes, R. & Strong, A. (2000) Conservation of coral reefs after the 1998 global bleaching event. *Conservation Biology* 14: 5–15.

Gorham, E. (1991) Northern peatlands: role in the carbon cycle and probable responses to climate warming. *Ecological Applications* 1: 182–195.

Gorke, M. (2003) *The Death of our Planet's Species*. Washington, DC, USA: Island Press.

Goss-Custard, J. D. & Moser, M. E. (1988) Rates of change in the numbers of dunlin *Calidris alpina* wintering in British estuaries in relation to the spread of *Spartina anglica*. *Journal of Applied Ecology* 25: 95–109.

Gosselink, J. G. & Maltby, E. (1990) Wetland gains and losses. In: Wetlands: A Threatened Landscape, ed. M. Williams, pp. 296–322. Oxford, UK: Blackwell Scientific Publications.

Gottgens, J. F., Perry, J. E., Fortney, R., et al. (2001) The Paraguay–Paraná Hidrovia: protecting the Pantanal with lessons from the past. *BioScience* 51: 301–308.

Goulet, T. L. (2006) Most corals may not change their symbionts. *Marine Ecology Progress Series* 321: 1–7.

Gowan, R. J., Mills, D. K., Trimmer, M. & Nedwell, D. B. (2000) Production and its fate in two coastal regions of the Irish Sea: the influence of anthropogenic nutrients. *Marine Ecology Progress Series* 208: 51–64.

Grachev, M. (1991) On the present state of the ecological system of Lake Baikal. Unpublished report. Irkutsk, USSR: Limnological Institute, Siberian Division, USSR Academy of Sciences. Graham, M. H. (1996) Effect of high irradiance on recruitment of the giant kelp *Macrocystis* (Phaeophyta) in shallow water. *Journal of Phycology* 32: 903–906.

Graham, M. H. (1997) Factors determining the upper limit of giant kelp, *Macrocystis pyrifera* Agardh, along the Monterey Peninsula, central California, USA. *Journal of Experimental Marine Biology and Ecology* 218: 127–149.

Graham, M. H., Dayton, P. K. & Erlandson, J. M. (2003) Ice ages and ecological transitions on temperate coasts. *Trends* in Ecology and Evolution 18: 33–40.

Graham, M. H., Harrold, C., Lisin, S., Light, K., Watanabe, J. M. & Foster, M. S. (1997) Population dynamics of giant kelp *Macrocystis pyrifera* along a wave exposure gradient. *Marine Ecology Progress Series* 148: 269–279.

Graham, N. E. (1995) Simulation of recent global temperature trends. *Science* 267: 666–671.

Graham, N. E. & Diaz, H. F. (2001) Evidence for intensification of North Pacific winter cyclones since 1948. Bulletin of the American Meteorological Society 82: 1869–1893.

Granger, S. L., Traver, M. S. & Nixon, S. W. (2000) Propagation of Zostera marina L. from seed. In: Seas at the Millennium: An Environmental Evaluation, vol. 3, Global Issues and Processes, ed. C. R. C. Sheppard, pp. 4–5. Amsterdam, the Netherlands: Pergamon.

Granina, L. (1997) The chemical budget of Lake Baikal: a review. *Limnology and Oceanography* **42**: 373–378.

Grant, A. & Briggs, A. D. (2002) Toxicity of sediments from around a North Sea oil platform: are metals or hydrocarbons responsible for ecological impacts? *Marine Environmental Research* 53: 95–116.

Grant, A., Hateley, J. G. & Jones, N. V. (1989) Mapping the ecological impact of heavy metals on the estuarine polychaete *Nereis diversicolor* using inherited metal tolerance. *Marine Pollution Bulletin* 20: 235–238.

Grant, J., Hatcher, A., Scott, D. B., *et al.* (1995) A multidisciplinary approach to evaluating impacts of shellfish aquaculture on benthic communities. *Estuaries* 18: 124–144.

Grassle, J. F. (1991) Effects of sewage sludge on deep-sea communities. *Eos* 72: 84 (abstract).

Grassle, J. F. (2000) The ocean biogeographic system (OBIS): an on-line, worldwide atlas for accessing, modeling and mapping marine biological data in a multidimensional geographic context. *Oceanography* **13**: 5–7.

Gray, A.J. (1977) Reclaimed land. In: *The Coastline*, ed. R.S.K. Barnes, pp. 253–270. Chichester, UK: John Wiley. Great Lakes Commission (2003) Toward a water resources management decision support system for the Great Lakes–St Lawrence River Basin. Unpublished report on CD-ROM. Ann Arbor, MI, USA: Great Lakes Commission.

Great Lakes Environmental Research Laboratory (2002) Great Lakes fish community impacted by *Diporeia* disappearance. Unpublished fact sheet. Ann Arbor, MI, USA: Great Lakes Environmental Research Laboratory.

Great Lakes Fishery Commission (2001*a*) TFM and sea lamprey control. Unpublished fact sheet no. 4. Ann Arbor, MI, USA: Great Lakes Fishery Commission.

Great Lakes Fishery Commission (2001*b*) Sea lamprey barriers. Unpublished fact sheet no. 5. Ann Arbor, MI, USA: Great Lakes Fishery Commission.

Green, E. P. & Short, F. T., eds. (2003) *World Atlas of Seagrasses.* Berkeley, CA, USA: University of California Press.

Green, M. O., Vincent, C. E., McCave, I. N., et al. (1995) Storm sediment transport: observations from the British North sea shelf. Continental Shelf Research 15: 889–912.

Greene, K. (2002) Beach Nourishment: A Review of the Biological and Physical Impacts, ASMFC Habitat Management Series No.7. Washington, DC, USA: Atlantic States Marine Fisheries Commission.

Greenstein, B. J., Curran, H. A. & Pondolfi, J. M. (1998) Shifting ecological baselines and the demise of *Acropora cervicornis* in the western North Atlantic and Caribbean Province: a Pleistocene perspective. *Coral Reefs* 17: 249–261.

Gregory, J. M., Stott, P. A., Cresswell, D. J., et al. (2002) Recent and future changes in Arctic sea ice simulated by the HadCM3 AOGCM. *Geophysical Research Letters* 29: art. no. 2175.

Gregory, S. V., Swanson, F. J., McKee, W. A. & Cummins, K. W. (1991) An ecosystem perspective of riparian zones. *BioScience* 41: 540–551.

Grevemeyer, I., Herber, R. & Essen, H. H. (2000) Microseismological evidence for changing wave climate in the northeast Atlantic Ocean. *Nature* 408: 349–352.

Grevstad, F. S., Strong, D. R., Garcia-Rossi, M., Switzer, R. W.
& Wecker, M. S. (2003) Biological control of *Spartina* alterniflora in Willapa Bay, Washington using the planthopper *Prokelisia marginata:* agent specificity and early results. *Biological Control* 27: 32–42.

Griebler, C., Danielopol, D. L., Gibert, J., Nachtnebel, H. P.
& Notenboom, J., eds. (2001) Groundwater Ecology: A Tool for Management of Water Resources. Luxembourg: Office for Official Publications of the European Communities.

Griebler, C., Mindl, B., Slezak, D. & Geiger-Kaiser, M. (2002) Distribution patterns of attached and suspended bacteria in pristine and contaminated shallow aquifers studied with an in situ sediment exposure microcosm. *Aquatic Microbial Ecology* 28: 117–129.

Griffing, S. (2003) The wild world of West Nile. Smithsonian Environmental Research Center Newsletter Spring Edition 1: 4–5.

Griffiths, C., Hockey, P., van Erkom Schurink, C. & le Roux, P. (1992) Marine invasive aliens on South African shores: implications for community structure and trophic functioning. *South African Journal of Marine Science* 12: 713–722.

Grigal, D. F., Buttleman, C. G. & Kernick, L. K. (1985) Biomass and productivity of the woody strata of forested bogs in northern Minnesota. *Canadian Journal of Botany* 63: 2416–2424.

Grigg, R. W. (1993) Precious coral fisheries of Hawaii and the US Pacific Islands. *Marine Fisheries Review* 55: 50–60.

Grigg, R. W. (1997) Paleoceanography of coral reefs in the Hawaiian–Emperor Chain revisited. *Coral Reefs* 16: S33–S38.

Grimm, N. B. & Fisher, S. G. (1992) Response of arid land streams to changing climate. In: *Global Climate Change* and Freshmater Ecosystems, ed. P. Firth & S. G. Fisher, pp. 211–233. Berlin, Germany: Springer-Verlag.

Grimm, N. B., Chacon, A., Dahm, C., et al. (1997) Sensitivity of aquatic ecosystems to climatic and anthropogenic changes: the basin and range, American southwest and Mexico. *Hydrological Processes* 11: 1023–1041.

Grizzle, R. E., Short, F. T., Hoven, H., Kindbloom, L. & Newell, C. R. (1996) Hydrodynamically induced synchronous waving of seagrasses: 'monami' and its possible effects on larval mussel settlement. *Journal of Experimental Marine Biology and Ecology* 206: 165–177.

Groombridge, B. (1992) *Global Biodiversity: Status of the Earth's Living Resources.* London, UK: World Conservation Monitoring Centre and Chapman & Hall.

Groombridge, B. & Jenkins, M. (1998) Freshwater Biodiversity: A Preliminary Global Assessment, WCMC Biodiversity Series No. 8. Cambridge, UK: World Conservation Monitoring Centre, World Conservation Press.

Grootjans, A. P., Everts, H., Bruin, K. & Fresco, L. (2001) Restoration of wet dune slacks on the Dutch Wadden Sea Islands: recolonization after large-scale sod cutting. *Restoration Ecology* 9: 137–146.

Grosholz, E. (2002) Ecological and evolutionary consequences of coastal invasions. *Trends in Ecology and Evolution* 17: 22–27. Grosholz, E. D., Ruiz, G. M., Dean, C. A., *et al.* (2000) The impacts of a non-indigenous marine predator in a California bay. *Ecology* 81: 1206–1224.

Grunawalt, R. J., King, J. E. & McClain, R. S., eds. (1996) Protection of the Environment during Armed Conflict, International Law Studies No. 69 Newport, RI, USA: Naval War College.

Guilford, S. & Hecky, R. (2000) Total nitrogen, total phosphorus and nutrient limitation in lakes and oceans: is there a common relationship? *Limnology and Oceanography* 45: 1213–1223.

Guillén, J. E., Ramos, A. A., Martinez, L. & Sanchez Lizaso, J. L. (1994) Anti-trawling reefs and the protection of *Posidonia oceanica* (L.) Delile meadows in the western Mediterranean Sea: demand and aims. *Bulletin of Marine Science* 55: 645–650.

Gunatilaka, A. & Dreher, J. (2001) Management of urban groundwater. II. Qualitative aspects. In: Groundwater Ecology: A Tool for Management of Water Resources, ed.
C. Griebler, D. L. Danielopol, J. Gibert, H. P. Nachtnebel & J. Notenboom, pp. 213–230. Luxembourg: Office for Official Publications of the European Communities.

Gunn, J. & Keller, W. (1990) Biological recovery of an acidified lake after reductions in industrial emissions of sulphur. *Nature* 345: 431–433.

Gunnarsson, U. & Rydin, H. (2000) Nitrogen fertilization reduces *Sphagnum* production in bog communities. *New Phytologist* 147: 527–537.

Gunnarsson, U., Malmer, N. & Rydin, H. (2002) Dynamics or constancy in *Sphagnum* dominated mire ecosystems: a 40-year study. *Ecography* 25: 685–704.

Gunnarsson, U., Rydin, H. & Sjörs, H. (2000) Diversity and pH changes after 50 years on the boreal mire Skattlösbergs Stormosse, central Sweden. *Journal of Vegetation Science* 11: 277–286.

Gustavson, K., Garde, K., Wangberg, S.-A. & Selmer, J.-S. (2000) Influence of UV-B radiation on bacterial activity in coastal waters. *Journal of Plankton Research* 22: 1501–1511.

Gutleb, A. C., Kranz., A, Nechay, G. & Toman, A. (1998) Heavy metal concentrations in livers and kidneys of the otter (*Lutra lutra*) from central Europe. *Bulletin* of Environmental Contamination and Toxicology 60: 273–279.

Hacker, S. D., Heimer, C. E., Hellquist, C. E., et al. (2001) A marine plant (*Spartina anglica*) invades widely varying habitats: potential mechanisms of invasion and control. *Biological Invasions* 3: 211–217. Hader, D. P. (1997) Effects of UV radiation on phytoplankton. In: Advances in Microbial Ecology, ed. J. G. Jones, pp. 1–26. New York, USA: Plenum Press.

Hader, D. P., Kumar, H. D., Smith, R. C. & Worrest, R. C. (2003) Aquatic ecosystems: effects of solar ultraviolet radiation and interactions with other climatic change factors. *Photochemical and Photobiological Sciences* 2: 39–50.

Haedrich, R. L. (1995) Structure over time of an exploited deep-water fish assemblage. In *Deep-Water Fisheries of the North Atlantic Oceanic Slope*, ed. A. G. Hopper, pp. 27–50. Dordrecht, the Netherlands: Kluwer.

Hagen, N. T. (1983) Destructive grazing of kelp beds by sea urchins in Vestfjorden, northern Norway. Sarsia 68: 177–190.

Hahn, H. J. (2002) Distribution of the aquatic meiofauna of the Marbling Brook catchment (Western Australia) with reference to land-use and hydrological features. *Archiv für Hydrobiologie* 139 (Suppl.): 237–263.

Hakenkamp, C. C. & Morin, A. (2000) The importance of meiofauna to lotic ecosystem functioning. *Freshwater Biology* 44: 165–176.

Hall, A. & Visbeck, M. (2002) Synchronous variability in the Southern Hemisphere atmosphere, sea ice, and ocean resulting from the Annular Mode. *Journal of Climate* 15: 3043–3057.

Hall, C. M. (2001) Trends in ocean and coastal tourism: the end of the last frontier? *Ocean and Coastal Management* 44: 601–618.

Hall, M. J. & Pilkey, O. H. (1991) Effects of hard stabilization on dry beach width for New Jersey. *Journal of Coastal Research* 7: 771–785.

Hall, S. & Mills, E. (2000) Exotic species in large lakes of the world. *Aquatic Ecosystem Health and Management* 3: 105–135.

Hall, S. J. (1999) The Effects of Fishing on Marine Ecosystems and Communities. Oxford, UK: Blackwell Science.

Hall, S. J. (2001) Is offshore exploration good for benthic conservation? *Trends in Ecology and Evolution* 16: 601–618.

Hall, S. J. (2002) The soft-sediment continental shelf benthic ecosystem: current status, agents for change and future prospects. *Environmental Conservation* 29: 350–374.

Hallegraeff, G. M. (1998) Transport of toxic dinoflagellates via ships' ballast water: bioeconomic risk assessment and efficacy of possible ballast water management strategies. *Marine Ecology Progress Series* 168: 297–309.

Hallerman, E. M. & Kapuscinski, A. R. (1995) Incorporating risk assessment and risk management into public policies on genetically modified finfish and shellfish. *Aquaculture* 137: 9–17. Halling-Sørensen, B., Nors Nielsen, S., Lanzky, P. F., *et al.* (1998) Occurrence, fate and effects of pharmaceutical substances in the environment: a review. *Chemosphere* **36**: 357–393.

Halsey, L. A., Vitt, D. H. & Bauer, I. E. (1998) Peatland initiation during the Holocene in continental western Canada. *Climatic Change* 40: 315–342.

Hameedi, M. J. (1997) Strategy for monitoring the environment in the coastal zone. In: *Coastal Zone Management Imperative for Maritime Developing Nations*, ed.
B. U. Haq, pp. 111–142. Dordrecht, the Netherlands: Kluwer.

Hamelink, J. L., Landrum, P. F., Bergman, H. L. & Benson, W. H., eds. (1994) *Bioavailability: Physical*, *Chemical*, and *Biological Interactions*. Boca Raton, FL, USA: Lewis Publishers.

Hamilton, L. S. & Snedaker, S. C. (1984) Handbook for Mangrove Area Management. Honolulu, HI, USA: East–West Centre, IUCN and United Nations Environmental Programme.

Hamilton, S. K. (1999) Potential effects of a major navigation project (Paraguay–Paraná–Hidrovia) on inundation in the Pantanal floodplains. *Regulated Rivers: Research and Management* 15: 289–299.

Hamilton, S. K. & Lewis Jr, W. M. (1990) Physical characteristics of the fringing floodplain of the Orinoco River, Venezuela. *Intersciencia* 15: 491–500.

Hamilton, S. K., Sippel, S. J. & Melack, J. M. (1996) Inundation patterns in the Pantanal wetland of South America determined by passive microwave sensing. *Archiv für Hydrobiologie* 137: 1–23.

Hamm, L., Capobianco, M., Dette, H. H., et al. (2002) A summary of European experience with shore nourishment. *Coastal Engineering* 47: 237–264.

Hammer, U. T. (1986) *Saline Lake Ecosystems of the World*. Dordrecht, the Netherlands: Kluwer.

Hammer, U. T. (1990) The effects of climate change on the salinity, water levels and biota of Canadian prairie saline lakes. *Verhandlungen Internationale Vereinigung für Limnologie* 24: 321–326.

Hammerstrom, K. K., Kenworthy, W. J., Fonseca, M. S. & Whitfield, P. E. (2006) Seed bank, biomass, and productivity of *Halophila decipiens*, a deep water seagrass on the west Florida continental shelf. *Aquatic Botany* 84: 110–120.

Hammerton, D. (1972) The Nile river: a case study. In: *River Ecology and Man*, ed. R. T. Oglesby, C. A. Carlson & M. J. McCann, pp. 171–214. New York, USA: Academic Press.

Handa, I. T., Harmsen, R. & Jefferies, R. L. (2002) Patterns of vegetation change and the recovery potential of degraded areas in a coastal marsh system of the Hudson Bay lowlands. *Journal of Ecology* **90**: 86–99.

Hannides, A. K. & Smith, C. R. (2003) The Northeastern Pacific abyssal plain. In: *Biogeochemistry of Marine Systems*, ed. K. D. Black & G. B. Shimmield, pp. 208–237. Sheffield, UK: Sheffield Academic Press.

Hansen, M. (1999) Lake trout in the Great Lakes: basinwide stock collapse and binational restoration. In: *Great Lakes Fisheries Policy and Management*, ed. W. W. Taylor & C. P. Ferreri, pp. 417–453. East Lansing, MI, USA: Michigan State University Press.

Hansen, P. J., Bjornsen, P. K. & Hansen, B. W. (1997) Zooplankton grazing and growth scaling within the 2–2000 µm body size range. *Limnology and Oceanography* **42**: 687–704.

Hanski, I. (1998) Metapopulation dynamics. Nature 396: 41-49.

Hansson, L.-A., Annadotter, H., Bergman, E., et al. (1998) Biomanipulation as an application of food chain theory: constraints, synthesis and recommendations for temperate lakes. *Ecosystems* 1: 558–574.

Haque, C. E. & Zaman, M. Q. (1993) Human responses to riverine hazards in Bangladesh: a proposal for sustainable development. *World Development* 21: 93–107.

Haraszthy, L. (2001) The floodplain forests in Hungary. In: *The Floodplain Forests in Europe*, ed. E. Klimo & H. Hager, pp. 17–24. Leiden, the Netherlands: E. J. Brill.

Hardin, G. (1968) The tragedy of the commons. *Science* 162: 1243–1248.

Hardy, A. (1967) Great Waters. London, UK: Collins.

Harley, C. D. G., Hughes, A. R., Hultgren, K. M., et al. (2006) The impacts of climate change in coastal marine systems. *Ecology Letters* 9: 228–241.

Harmelin-Vivien, M., Francour, P. & Harmelin, J. G. (1999) Impact of *Caulerpa taxifolia* on Mediterranean fish assemblages: a six-year study. In: *Proceedings of the Workshop on Invasive* Caulerpa *in the Mediterranean*, MAP Technical Reports Series No. 125, pp. 127–138. Athens, Greece: United Nations Environmental Programme.

Harner, M. J. & Stanford, J. A. (2003) Differences in cottonwood growth between a losing and a gaining reach of an alluvial floodplain. *Ecology* 84: 1453–1458.

Harper, D. M., Mavuti, K. M. & Muchiri, S. M. (1990) Ecology and management of Lake Naivasha, Kenya, in relation to climatic change, alien species introductions, and agricultural development. *Environmental Conservation* 17: 328–336. Harris, J. M., Branch, G. M., Sibiya, C. & Bill, C. (2003) The Sokhulu subsistence mussel-harvesting project: co-management in action. In: *Waves of Change: Co-Management of Fisheries and Coastal Resources in South Africa*, pp. 61–98. Cape Town, South Africa: University of Cape Town Press.

Harrison, I. J. & Stiassny, M. L. J. (1999) The quiet crisis:
a preliminary listing of the freshwater fishes of the world that are extinct or 'missing in action'. In: *Extinctions in Near Time: Causes, Contexts and Consequences*, ed.
R. D. E. MacPhee, pp. 271–331. New York, USA: Plenum Press.

Harrold, C. & Lisin, S. (1989) Radio-tracking rafts of giant kelp: local production and regional transport. *Journal of Experimental Marine Biology and Ecology* 30: 237–252.

Harrold, C. & Reed, D. C. (1985) Food availability, sea urchin grazing, and kelp forest community structure. *Ecology* 66: 1160–1169.

Harrold, C., Light, K. & Lisin, S. (1998) Organic enrichment of submarine-canyon and continental-shelf benthic communities by macroalgal drift imported from nearshore kelp forests. *Limnology and Oceanography* 43: 669–678.

Hart, J. (1996) Storm over Mono: The Mono Lake Battle and the California Water Future. Berkeley, CA, USA: University of California Press.

Hartney, K. I. & Grorud, K. I. (2002) The effect of sea urchins as biogenic structures on the local abundance of a temperate reef fish. *Oecologia* 131: 506–513.

Hartnoll, R. G. & Hawkins, S. J. (1985) Patchiness and fluctuations on moderately exposed rocky shores. *Ophelia* 24: 53–63.

Harvell, C. D., Kim, K., Burkholder, J. M., et al. (1999) Emerging marine diseases: climate links and anthropogenic factors. *Science* 285: 1505–1510.

Harvell, C. D., Mitchell, C. E., Ward, J. R., et al. (2002) Climate warming and disease risks for terrestrial and marine biota. Science 296: 2158–2162.

Harvey, G. R., Miklas, H. P., Bowen, V. T. & Steinhauer, W. G. (1974a) Observations on the distribution of chlorinated hydrocarbons in Atlantic Ocean organisms. *Journal of Marine Research* 32: 103–118.

Harvey, G. R., Steinhauer, W. G. & Miklas, H. P. (1974b) Decline of PCB concentrations in North Atlantic surface water. *Nature* 252: 387–388.

Harwell, M. C. & Orth, R. J. (1999) Eelgrass (*Zostera marina* L.) seed protection for field experiments and implications for large-scale restoration. *Aquatic Botany* 64: 54–61. Hassan, F. A., Reuss, M., Trottier, J., et al. (2003) History and Future of Shared Water Resources, UNESCO-IHP Technical Documents in Hydrology, PCCP Series No. 6. Paris, France: UNESCO.

Hässelholms Vatten (2007) Home page [www document]. URL http://www.hasselholmsvatten.se (in Swedish).

Hatcher, B. G. (1983) Grazing in coral reef ecosystems. In: *Perspectives on Coral Reefs*, ed. D. J. Barnes, pp. 164–179. Canberra, ACT, Australia: Brian Clouster.

Hatcher, B. G., Kirkman, H. & Wood, W. F. (1987) Growth of the kelp *Ecklonia radiata* near the northern limit of its range in Western Australia. *Marine Biology* 95: 63–73.

Hateley, J. G., Grant, A., Taylor, S. M. & Jones, N. V. (1992) Morphological and other evidence on the degree of genetic differentiation between populations of *Nereis diversicolor*. *Journal of the Marine Biological Association of the United Kingdom* 72: 365–381.

Hatton, H. (1938) Essais de bionomie explicative sur quelques espèces intercotidals d'algues et d'animaux. *Annales de l'Institut Océanographique de Monaco* 17: 241–348.

Hatton, T. & Salama, R. (1999) Is it feasible to restore the salinity affected rivers of the Western Australian wheatbelt?
In: *Proceedings of the 2nd Australian Stream Management Conference*, vol. 1, ed. I. R. Rutherfurd & R. Bartley, pp. 313–317. Adelaide, SA, Australia: Cooperative Research Centre for Catchment Hydrology.

Hawkins, J. P. & Roberts, C. M. (1992) Effects of recreational SCUBA diving on fore-reef slope communities of coral reefs. *Biological Conservation* 62: 171–178.

Hawkins, S. J. (1999) Experimental ecology and coastal conservation: conflicts on rocky shores. *Aquatic Conservation: Marine and Freshwater Ecosystems* 9: 565–572.

Hawkins, S. J. & Hartnoll, R. G. (1985) Factors determining the upper limits of intertidal canopy forming algae. *Marine Ecology Progress Series* 20: 265–271.

Hawkins, S. J. & Southward, A. J. (1992) The *Torrey Canyon* oil spill: recovery of rocky shore communities. In: *Restoring the Nation's Environment*, ed. G. W. Thayer, pp. 583–631. College Park, MD, USA: Maryland Sea Grant College.

Hawkins, S. J., Allen, J. R., Fielding, N. J., Wilkinson, S. B. & Wallace, I. D. (1999) Liverpool Bay and the estuaries: human impact, recent recovery and restoration. In: *Ecology* and Landscape Development: A History of the Mersey Basin, Conference Proceedings, ed. E. F. Greenwood, pp. 155–165. Liverpool, UK: Liverpool University Press.

Hawkins, S. J., Allen, J. R., Ross, P. M. & Genner, M. J. (2002a) Marine and coastal ecosystems. In *Handbook of Ecological Restoration*, vol. 2, *Restoration in Practice*, ed. M. R. Perrow & A. J. Davy, pp. 121–148. Cambridge, UK: Cambridge University Press.

Hawkins, S. J., Corte-Real, H. B. S. M., Pannacciulli, F. G., Weber, L. C. & Bishop, J. D. D. (2000) Thoughts on the ecology and evolution of the intertidal biota of the Azores and other Atlantic Islands. *Hydrobiologia* 440: 3–17.

Hawkins, S. J., Gibbs, P. E., Pope, N. D., et al. (2002b) Recovery of polluted coastal ecosystems: the case for longterm studies. *Marine Environmental Research* 54: 215–222.

Hawkins, S. J., Proud, S. V., Spence, S. K. & Southward, A. J. (1994) From the individual to the community and beyond: water quality, stress indicators and key species in coastal ecosystems. In: Water Quality and Stress Indicators in Marine and Freshwater Ecosystems: Linking Levels of Organisation (Individuals, Populations, Communities), ed. D. W. Sutcliffe, pp. 35–62. Ambleside, UK: Freshwater Biological Association.

Hawkins, S. J., Southward, A. J. & Barrett, R. L. (1983) Population structure of *Patella vulgata* during succession on rocky shores in southwest England. *Oceanologica Acta* Special Issue: 103–107.

Haworth, E. Y. (1969) The diatoms of a sediment core from Blea Tarn, Langdale. *Journal of Ecology* 57: 429–439.

Haworth, E. Y. (1972) Diatom succession in a core from Pickerel Lake, north-eastern South Dakota. *Geological Society of America Bulletin* 83: 157–172.

Hay, M. E. (1984) Patterns of fish and urchin grazing on Caribbean coral reefs: are previous results typical? *Ecology* 65: 446–454.

Hay, M. E., Colburn, T. & Downing, D. (1983) Spatial and temporal patterns in herbivory on a Caribbean fringing reef: the effects on plant distribution. *Oecologia* 58: 299–308.

Haycock, N. E., Burt, T., Goulding, K. & Pinay, G. (1997) Buffer Zones: Their Processes and Potential in Water Protection. Hertford, UK: Quest Environmental.

Hazen, T. C. (1997) Bioremediation. In: *The Microbiology* of the Deep Subsurface, ed. P. S. Amy & D. L. Haldeman, pp. 247–266. Boca Raton, FL, USA: Lewis Publishers.

Headland, R. (1984) The Island of South Georgia. Cambridge, UK: Cambridge University Press.

Hearn, C. J., Atkinson, M. J. & Falter, J. L. (2001) A physical derivation of nutrient-uptake rates in coral reefs: effects of roughness and waves. *Coral Reefs* 20: 347–356.

Heathwaite, A. L., Johnes, P. J. & Peters, N. (1996) Trends in nutrients. *Hydrological Processes* 10: 263–293.

Heck, K. L. & Valentine, J. F. (1995) Sea urchin herbivory: evidence for long-lasting effects in subtropical seagrass meadows. *Journal of Experimental Marine Biology and Ecology* 189: 205–217.

- Hecky, R. (1993) The eutrophication of Lake Victoria. Verhandlungen Internationale Vereinigung für Limnologie 25: 39–48.
- Hecky, R. & Bugenyi, F. (1992) Hydrology and chemistry of the African Great Lakes and water quality issues: problems and solutions. *Mitteilungen Internationale Vereinigung für Limnologie* 23: 45–54.
- Hecky, R., Botsma, H., Mugidde, R. & Bugenyi, F. (1996)
 Phosphorus pumps, nitrogen sinks, silicon drains: plumbing nutrients in the African Great Lakes. In: *The Limnology, Climatology, and Paleoclimatology of the East African Lakes*, ed. T. C. Johnson & E. Odada, pp. 205–224. Toronto, Ontario, Canada: Gordon & Breach.
- Hecky, R., Bugenyi, F., Ochumba, P., et al. (1994)Deoxygenation of the deep water of Lake Victoria. Limnology and Oceanography 39: 1476–1480.
- Hecky, R., Smith, R., Barton, D., et al. (2004) The near shore phosphorus shunt: a consequence of ecosystem engineering by dreissenids in the Lawrentian Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 61: 1285–1293.
- Hector, A., Schmid, B., Beierkuhnlein, C., et al. (1999) Plant diversity and productivity experiments in European grasslands. Science 286: 1123–1127.
- Heeg, J. & Breen, C. M., eds. (1982) Man and the Pongolo Floodplain, South African National Scientific Programs Report No. 56. CSIR, Pretoria, South Africa:
- Heemsbergen, D. A., Berg, M. P., Loreau, M., et al. (2004) Biodiversity effects on soil processes explained by interspecific functional dissimilarity. *Science* 306: 1019–1020.
- Heidelbaugh, W. S. & Nelson, W. G. (1996) A power analysis of methods for assessing change in seagrass cover. *Aquatic Botany* 53: 227–233.
- Hein, M. & Sand-Jensen, K. (1997) CO₂ increases oceanic primary production. *Nature* 388: 526–527.
- Heinz Center (2002) Dam Removal: Science and Decision Making. Washington, DC, USA: The Heinz Center.
- Hellberg, M. E., Balch, D. P. & Roy, K. (2001) Climate-driven range expansion and morphological evolution in a marine gastropod. *Science* 292: 1707–1710.
- Hellström, C., Chen, D. L., Archberger, C. & Räisänen, J. (2001) Comparison of climate change scenarios for Sweden based on statistical and dynamical downscaling of monthly precipitation. *Climate Research* 19: 45–55.
- Helms, M., Buchele, B., Merkel, U. & Ihringer, J. (2002) Statistical analysis of the flood situation and assessment of the impact of diking measures along the Elbe (Labe) river. *Journal of Hydrology* 267: 94–114.

- Helmuth, B., Mieszkowska, N., Moore, P., Hawkins, S. J. (2006) Living on the edge of two changing worlds: forecasting responses of rocky intertidal ecosystems to climate change. *Annual Review of Ecology, Evolution and Systematics* 37: 373–404.
- Hemminga, M. A. & Duarte, C. M. (2000) Seagrass Ecology. Cambridge, UK: Cambridge University Press.
- Hemond, H. F. & Benoit, J. (1988) Cumulative impacts on water quality functions of wetlands. *Environmental Management* 12: 639–653.
- Henley, W. J. & Dunton, K. H. (1997) Effects of nitrogen supply and continuous darkness on growth and photosynthesis of the arctic kelp *Laminaria solidungula*. *Limnology and Oceanography* 42: 209–216.
- Hensen, V. (1887) Über die Bestimmung des Planktons oder des im Meere treibenden Materials an Pflanzen und Thieren. Kommission zur wissenschaftlichen Untersuchungen der deutschen Meere in Kiel, 1882–1886. V. Bericht, Jahrgang 12–16.
- Herbert, R. J. H., Hawkins, S. J., Sheader, M. & Southward, A. J. (2003) Range extension and reproduction of the barnacle *Balanus perforatus* in the Eastern English Channel. *Journal of the Marine Biological Association of the United Kingdom* 83: 73–82.
- Herdendorf, C. (1982) Large lakes of the world. Journal of Great Lakes Research 8: 379–412.
- Herdendorf, C. E. (1990) Distribution of the world's large lakes. In: *Large Lakes: Ecological Structure and Function*, ed. M. M. Tilzer & C. Serruya, pp. 3–38. New York, USA: Springer-Verlag.
- Herman, J. S., Culver, D. C. & Salzman, J. (2001) Groundwater ecosystems and the service of water purification. *Stanford Environmental Law Journal* 20: 479–498.
- Hermsen, J. M., Collie, J. S. & Valentine, P. C. (2003) Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series* 260: 97–108
- Hershey, A. (1985) Effects of predatory sculpin on the chironomid communities in an Arctic lake. *Ecology* 66: 1131–1138.
- Herut, B., Krom, M., Pan, G. & Mortimer, R. (1999) Atmospheric input of nitrogen and phosphorus to the Southeast Mediterranean: sources, fluxes, and possible impact. *Limnology and Oceanography* 44: 1683–1692.
- Hessen, D. O., Hindar, A. & Holtan, G. (1997) The significance of nitrogen runoff for eutrophication of freshwater and marine recipients. *Ambio* 26: 312–320.

Heuveline, P. (1999) The global impact of mortality and fertility transitions, 1950–2000. *Population Development Review* December: 681–702.

Hewes, L. & Frandson, P. E. (1952) Occupying the wet prairie: the role of artificial drainage in Story County, Iowa. Annals of the Association of American Geographers 42: 24–50.

Hiaasen, C. (1990) Double Whammy. London, UK: Pan Books.

Hickel, W., Mangelsdorf, P. & Berg, J. (1993) The human impact in the German Bight: eutrophication during 3 decades (1962–1991). *Helgoländer Meeresuntersuchungen* 47: 243–263.

Highsmith, R. C. (1980) Geographic patterns of coral bioerosion: a productivity hypothesis. *Journal of Experimental Marine Biology and Ecology* **46**: 55–67.

Hildrew, A. G. & Giller, P. S. (1994) Patchiness, species interactions and disturbance in the stream benthos. In: *Aquatic Ecology: Scale, Pattern and Process*, ed. P. S. Giller, A. G. Hildrew & D. G. Raffaelli, pp. 21–62. Oxford, UK: Blackwell Scientific Publications.

Hildrew, A. G. & Ormerod, S. J. (1995) Acidification: causes, consequences and solutions. In: *The Ecological Basis for River Management*, ed. D. M. Harper & A. Ferguson, pp. 147–160. Chichester, UK: John Wiley.

Hill, B. J. (1977) The effect of heated effluent on egg production in the estuarine prawn Upogebia africana (Ortmann). Journal of Experimental Marine Biology and Ecology 29: 291–302.

Hilton-Taylor, C., compiler (2000) *IUCN Red List of Threatened Species.* Gland, Switzerland and Cambridge, UK: IUCN.

Hily, C. (1991) Is the activity of benthic suspension feeders a factor controlling water quality in the Bay of Brest? *Marine Ecology Progress Series* 69: 179–188.

Hily, C., van Katwijk, M. M. & den Hartog, C. (2003) The seagrasses of western Europe. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 46–55. Berkeley, CA, USA: University of California Press.

Himmelman, J. H. (1980) The role of the green sea urchin Strongylocentrotus droebachiensis in the rocky subtidal region of Newfoundland. In: Proceedings of the Workshop on the Relationship between Sea Urchin Grazing and Commercial Plant/Animal Harvesting, ed. J. D. Pringle, G. J. Sharp & J. F. Caddy, pp. 92–119. Halifax, NS, Canada: Resource Branch, Fisheries and Oceans Canada.

Hiscock, K., Southward, A., Tittley, I., Hawkins, S. (2004) Effects of changing temperature on benthic marine life in Britain and Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 14: 333–362. Hjorleifsson, E., Kassa, O. & Gunnarsson, K. (1995) Grazing of kelp by green sea urchins in Eyyjafjordu, North Iceland.
In: Ecology of Fjords and Coastal Waters: Proceedings of the Mare Nor Symposium on the Ecology of Fjords and Coastal Waters, Tromso, Norway, 5–9 December 1994, ed.
H. R. Skjoldal, C. Hopkins, K. E. Erikstad & H. P. Leinass,

pp. 593-597. Amsterdam, the Netherlands: Elsevier.

Hobday, A. J. (2000) Abundance and dispersal of drifting kelp Macrocystis pyrifera rafts in the Southern California Bight. Marine Ecology Progress Series 195: 101–116.

Hockey, P. A. R. & Bosman, A. L. (1986) Man as an intertidal predator in Transkei: disturbance, community convergence and management of a natural food resource. *Oikos* 46: 3–14.

Hockey, P. A. R. & van Erkom Schurink, C. (1992) The invasive biology of the mussel *Mytilus galloprovincialis* on the southern African Coast. *Transactions of the Royal Society* of South Africa 48: 123–139.

Hodgkin, E. P. & Hamilton, B. (1998) Changing estuarine wetlands: a long-term perspective for management. In: *Wetlands for the Future*, ed. A. J. McComb & J. A. Davis, pp. 243–255. Adelaide, SA, Australia: Gleneagles Publishing.

Hodgson, G. & Dixon, J. A. (1992) Sedimentation damage to marine resources: environmental and economic analysis.
In: *Resources and Environment in Asia Marine Sector*, ed.
J. B. March, pp. 421–426. Washington, DC, USA: Taylor & Francis.

Hoegh-Guldberg, O. (1999) Climate change, coral bleaching and the future of the world's coral reefs. *Marine and Freshwater Research* 50: 839–866.

Hoegh-Guldberg, O., Jones, R. J., Ward, S. & Loh, W. K. (2002) Is coral bleaching really adaptive? *Nature* 415: 601–602.

Hoekstra, A. Y. & Hung, P. Q. (2002) Virtual Water Trade: a Quantification of Virtual Water Flows between Nations in Relation to International Crop Trade, Value of Water Research Report Series No. 11. Delft, the Netherlands: National Institute for Public Health and the Environment (IHE).

Hofmann, E. E., Klinck, J. M., Locarnini, R. A., Fach, B. & Murphy, E. J. (1998) Krill transport in the Scotia Sea and environs. *Antarctic Science* 10: 406–415.

Hogarth, P.J. (1999) *The Biology of Mangroves*. Oxford, UK: Oxford University Press.

Holbrook, S. J., Schmitt, R. J. & Ambrose, R. F. (1990) Biogenic habitat structure and characteristics of temperate reef fish assemblages. *Australian Journal of Ecology* 15: 489–503.

- Holbrook, S. J., Swarbrick, S. L., Schmitt, R. J. & Ambrose, R. F. (1993) Reef architecture and reef fish: correlations of population densities with attributes of subtidal rocky environments. In: *The Ecology of Temperate Reefs: Proceedings of the 2nd International Temperate Reef Symposium*, Auckland, pp. 99–106. Wellington, New Zealand: National Institute of Water and Atmospheric Research.
- Holdridge, L. R., Grenke, W. C., Hatheway, W. H., Liang, T. & Tosi Jr, J. A. (1971) *Forest Environments in Tropical Life Zones*. New York, USA: Pergamon Press.
- Holland, R. (1993) Changes in planktonic diatoms and water transparency in Hatchery Bay, Bass Island area, western Lake Erie since the establishment of the zebra mussel. *Journal of Great Lakes Research* 19: 617–624.
- Holland, R. & Beeton, A. (1972) Significance to eutrophication of spatial differences in nutrients and diatoms in Lake Michigan. *Limnology and Oceanography* 17: 88–96.
- Holland, R., Johengen, T. & Beeton, A. (1995) Trends in nutrient concentrations in Hatchery Bay, Western Lake Erie, before and after *Dreissena polymorpha*. *Canadian Journal of Fisheries and Aquatic Sciences* 52: 1202–1209.
- Hollis, G. E. (1998) Future wetlands in a world short of water. In: Wetlands for the Future, ed. A. J. McComb & J. A. Davis, pp. 5–18. Adelaide, SA, Australia: Gleneagles Publishing.
- Holloway, G. & Sou, T. (2002) Has Arctic sea ice rapidly thinned? *Journal of the American Meteorological Society* 15: 1691–1701.
- Holmer, M. (1999) The effect of oxygen depletion on anaerobic organic matter degradation in marine sediments. *Estuarine, Coastal and Shelf Science* 48: 383–390.
- Holmer, M., Andersen, F. O., Nielsen, S. L. & Boschker, H. T. S. (2001) The importance of mineralization based on sulfate reduction for nutrient regeneration in tropical seagrass sediments. *Aquatic Botany* **71**: 1–17.
- Holmes, K. E., Edinger, E. N., Limmon, H. G. V. & Risk, M. J. (2000) Bioerosion of live massive corals and branching coral rubble on Indonesian coral reefs. *Marine Pollution Bulletin* 40: 606–617.
- Holmes, N. T. H. (1999) Recovery of headwater stream flora following the 1989–1992 groundwater drought. *Hydrological Processes* 13: 341–354.
- Holm-Hansen, O., Helbling, E. W. & Lubin, D. (1993) Ultraviolet radiation in Antarctica: inhibition of primary production. *Photochemistry and Photobiology* 58: 567–570.
- Holmquist, C. (1959) Problems on marine-glacial relicts on account of investigations on the genus *Mysis*. Unpublished report. Lund, Sweden: Berlindska.

- Hood, G. W. & Naiman, R. J. (2000) Vulnerability of riparian zones to invasion by exotic vascular plants. *Plant Ecology* 148: 105–114.
- Hooper, R. (1980) Observations on algal–grazer interactions in Newfoundland and Labrador. In: Proceedings of the Workshop on the Relationship between Sea Urchin Grazing and Commercial Plant/Animal Harvesting, ed. J. D. Pringle, G. J. Sharp & J. F. Caddy, pp. 120–124. Halifax, NS, Canada: Resource Branch, Fisheries and Oceans Canada.
- Hopkins, J. S., Sandifer, P. A., DeVoe, M. R., *et al.* (1995) Environmental impacts of shrimp farming with special reference to the situation in the continental United States. *Estuaries* 18: 25–42.
- Hopkinson, C. (1992) A comparison of ecosystem dynamics in fresh-water wetlands. *Estuaries* 15: 549–562.
- Horner, R., Ackley, S. F., Dieckmann, G. S., et al. (1992) Ecology of sea ice biota. I. Habitat, terminology and methodlogy. *Polar Biology* 12: 417–427.
- Hornung, H., Krom, M. D., Cohen, Y. & Bernhard, M. (1993) Trace metal content in deep-water sharks from the eastern Mediterranean Sea. *Marine Biology* 115: 331–338.
- Houde, E., Coleman, F., Dayton, P., et al. (2001) Marine Protected Areas: Tools for Sustaining Ocean Ecosystems.Washington, DC, USA: National Academy Press.
- Houghton, J. (1999) *Global Warming: The Complete Briefing*, 2nd edn. Cambridge, UK: Cambridge University Press.
- Houghton, J. T., Jenkins, G. J. & Ephrauns, J. J., eds. (1990) Scientific Assessment of Climate Change: Report of Working Group 1. Cambridge, UK: Cambridge University Press.
- Houghton, J. T., Meira Filho, L. G., Bruce, J., et al. (1996) Climate Change 1995: World Meteorological Organization/ Intergovernmental Panel on Climate Change. Cambridge, UK: Cambridge University Press.
- Houlahan, J. E., Findlay, C. S., Schmidt, B. R., Meyer, A. H. & Kuzmin, S. L. (2000) Quantitative evidence for global amphibian population declines. *Nature* 404: 752–755.
- Houston, J. R. (2002) The economic value of beaches: a 2002 update. *Shore and Beach* 70: 9–12.
- Howard-Williams, C. & Thompson, K. (1985) The conservation and management of African wetlands. In: *The Ecology and Management of African Wetland Vegetation*, ed. P. Denny, pp. 203–230. The Hague, the Netherlands: Dr W. Junk.
- Howarth, R. W., Anderson, D. M., Church, T. M., (2000a) Clean Coastal Waters: Understanding and Reducing the Effects of Nutrient Pollution. Washington, DC, USA: Ocean Studies Board and Water Science and Technology Board, National Academy Press.

Howarth, R. W., Anderson, D., Cloern, J., *et al.* (2000b) Nutrient pollution of coastal rivers, bays, and seas. *Issues in Ecology* 7: 1–15.

Howarth, R. W., Sharpley, A. & Walker, D. (2002) Sources of nutrient pollution to coastal waters in the United States: implications for achieving coastal water quality goals. *Estuaries* 25: 656–676.

Hu, S., Chapin, F. S., Firestone, M. K., Field, C. B. & Chlariello, N. R. (2001) Nitrogen limitation of microbial decomposition in a grassland under elevated CO₂. *Nature* 409: 188–191.

Hudson, P. F. & Colditz, R. R. (2003) Flood delineation in a large and complex alluvial valley, lower Pánuco basin, Mexico. *Journal of Hydrology* 280: 229–245.

Hughes, F. M. R. (1990) The influence of flooding regimes on forest distribution and composition in the Tana River floodplain, Kenya. *Journal of Applied Ecology* 27: 475–491.

Hughes, F. M. R., ed. (2003) The Flooded Forest: Guidance for Policy Makers and River Managers in Europe on the Restoration of Floodplain Forests, FLOBAR2. Cambridge, UK: European Union and Department of Geography, University of Cambridge.

Hughes, F. M. R. & Rood, S. B. (2003) The allocation of river flows for the restoration of floodplain forest ecosystems: a review of approaches and their applicability in Europe. *Environmental Management* 32: 12–33.

Hughes, P. D. M. & Barber, K. E. (2003) Mire development across the fen-bog transition on the Teifi floodplain at Tregaron Bog, Ceredigion, Wales, and a comparison with 13 other raised bogs. *Journal of Ecology* **91**: 253–264.

Hughes, T. P. (1994) Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science* **265**: 1547–1551.

Hughes, T. P. & Tanner, J. E. (2000) Recruitment failure, life histories, and long-term decline of Caribbean corals. *Ecology* 81: 2250–2263.

Hughes, T. P., Baird, A. H., Bellwood, D. R., *et al.* (2003) Climate change, human impacts, and the resilience of coral reefs. *Science* **301**: 929–933.

Hughes, T. P., Bellwood, D. R. & Connolly, S. R. (2002) Biodiversity hotspots, centres of endemicity, and the conservation of coral reefs. *Ecology Letters* 5: 775–784.

Hukkinen, J. (1998) Institutions, environmental management and long-term ecological sustenance. *Ambio* 27: 112–117.

Hulme, M., Jenkins, G. J., Lu, X., et al. (2002) Climate Change Scenarios for the United Kingdom: The UKCIP02 Scientific Report. Norwich, UK: Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia.

- Hultén, E. (1968) Flora of Alaska and Neighboring Territories: A Manual of Vascular Plants. Stanford, CA, USA: Stanford University Press.
- Humborg, C., Ittekkot, V., Cociasu, A. & VonBodungen, B. (1997) Effect of Danube River dam on Black Sea biogeochemistry and ecosystem structure. *Nature* 386: 385–388.

Humphries, P. & Baldwin, D. S. (2003) Drought and aquatic ecosystems: an introduction. *Freshwater Biology* 48: 1141–1146.

Humphries, S. E., Groves, R. H. & Mitchell, D. S. (1991) Plant invasions of Australian ecosystems: a status review and management directions. Unpublished report, Kowari 2. Canberra, ACT, Australia: Australian National Parks and Wildlife Service.

Hunt, R. J. & Wilcox, D. A. (2003) Ecohydrology: why hydrologists should care. *Ground Water* **41**: 289.

Huntington, H. P. (2000) Sustainable fisheries management. WWF Arctic Bulletin 1: 1–18.

Hurem, A. K. 1979. Rubble-mound structures as artificial reefs. In: *Coastal Structures 79*, pp. 1042–1051. New York, USA: American Society of Civil Engineers.

Hurrell, J. W. (1995) Decadal trends in the North Atlantic oscillation: regional temperatures and precipitation. *Science* 269: 676–679.

Huszar, P., Petermann, P., Leite, A., et al. (1999) Fact or Fiction: A Review of the Hidrovia Paraguay–Paraná Official Studies. Toronto, Ontanio, Canada: WWF.

Hutchin, P. R., Press, M. C., Lee, J. A. & Ashenden, T. W. (1995) Elevated concentrations of CO₂ may double methane emissions from mires. *Global Change Biology* 1: 125–128.

Hutchings, J. A. (2000) Collapse and recovery of marine fishes. *Nature* **406**: 882–885.

Hutchinson, G. E. (1957) *A Treatise on Limnology*, vol. 1. New York, USA: John Wiley.

Hwang, Y. H. & Morris, J. T. (1994) Whole plant gas exchange responses of *Spartina alterniflora* (Poaceae) to a range of constant and transient salinities. *American Journal* of Botany 81: 659–665.

Ingole, B., Ansari, Z. A., Rathod, V. & Rodrigues, N. (2001) Response of deep-sea macrobenthos to a small-scale environmental disturbance. *Deep-Sea Research II* 48: 3401–3410.

Ingram, H. A. P. (1978) Soil layers in mires: function and terminology. *Journal of Soil Science* 29: 224–227.

- Ingrid, G., Andersen, T. & Vadstein, O. (1996) Pelagic food webs and eutrophication of coastal waters: impact of grazers on algal communities. *Marine Pollution Bulletin* 33: 22–35.
- Inman, D. J., Masters, P. M. & Stone, K. E. (1991) Induced subsidence: environmental and legal implications. In: *Coastal Zone 91*, pp. 16–27. New York, USA: American Society of Civil Engineers.
- Innocenti, L. & Pranzini, E. (1993) Geomorphological evolution and sedimentology of the Ombrone River delta, Italy. *Journal of Coastal Research* 9: 481–493.
- International Board of Inquiry for the Great Lakes Fisheries (1943) Report and Supplement. Unpublished report. Washington, DC, USA: US Government Printing Office.
- International Joint Commission (1989) Revised Great Lakes water quality agreement of 1978. Unpublished report. Washington, DC, USA and Ottawa, Ontario, Canada: International Joint Commission. United States and Canada.
- International Joint Commission (2000) Tenth biennial report on Great Lakes water quality. Unpublished report.
 Washington, DC, USA and Ottawa, Ontario, Canada: International Joint Commission, United States and Canada.
- International Organization for Migration (2000) World Migration Report. New York, USA: IOM.
- International Tanker Owners Pollution Federation Limited (2000) Historical data [www.document]. URL http://www. itopf.com/stats.html
- Invers, O., Zimmerman, R. C., Alberte, R. S., Pérez, M. & Romero, J. (2001) Inorganic carbon sources for seagrass photosynthesis: an experimental evaluation of bicarbonate use in species inhabiting temperate waters. *Journal of Experimental Marine Biology and Ecology* 265: 203–217.
- IPCC (1995): Climate Change 1995: The IPCC Second Assessment Synthesis of Scientific–Technical Information Relevant to Interpreting Article 2 of the UN Framework Convention on Climate Change. Geneva, Switzerland: IPCC.
- IPCC (1996) Climate Change 1995: Impacts, Adaptations, and Mitigation of Climate Change – Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change, ed.
 R. T. Watson, M. C. Zinoyowera & R. H. Moss. Cambridge, UK: Cambridge University Press.
- IPCC (2001a) Climate Change 2001: The Scientific Basis Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, ed. J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell & C. A. Johnson. Cambridge, UK: Cambridge University Press.

- IPCC (2001b) Climate Change 2001: Impacts, Adaptation, and Vulnerability – Contribution of Working Group II to the Third Assessment Report of the Intergovernmental Panel on Climate Change, ed. J. J. McCarthy, O. F. Canziani, N. A. Leary, D. J. Dokken & K. S. White. Cambridge, UK: Cambridge University Press.
- IPCC (2001c) Climate Change 2001: Mitigation Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change, ed. B. Metz, O. Davidson, R. Swart & J. Pan. Cambridge, UK: Cambridge University Press.
- IPCC (2001d) Climate Change 2001: Synthesis Report A Contribution of Working Groups I, II, and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change, ed. R. T. Watson and the Core Writing Team. Cambridge, UK: Cambridge University Press.
- IPCC (2001e) IPCC Third Assessment Report: Climate Change 2001. Cambridge, UK: Cambridge University Press.
- IPCC (2002) Climate Change and Biodiversity, Intergovernmental Panel on Climate Change Technical Paper No. V, ed. H. Gitay, A. Suárez, R. T. Watson & D. J. Dokken. Geneva, Switzerland: IPCC. Available online at www.ipcc.ch/pub/tpbiodiv.pdf
- ISSLR (2007) International Society for Salt Lake Research: home page [www document]. URL http://www.isslr.org.
- Ittekkot, V., Jilan, S., Miles, E., et al. (1996) Oceans. In: Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change – Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change, ed. R. T. Watson, M. C. Zinoyowera & R. H. Moss, pp. 267–288. Cambridge, UK: Cambridge University Press.
- IUCN (1996) *IUCN Red List of Threatened Animals*. Gland, Switzerland: IUCN (World Conservation Union).
- IUCN (2000) Vision for Water and Nature: A World Strategy for Conservation and Sustainable Management of Water Resources in the Twenty-First Century. Gland, Switzerland: IUCN (World Conservation Union).
- Ivanov, K. E. (1981) *Water Movement in Mirelands*. London, UK: Academic Press.
- Ivanov, V. P., Kamakin, A. M., Ushintzev, V. B., et al. (2000) Invasion of the Caspian Sea by the comb jellyfish *Mnemiopsis leidyi* (Ctenophora). *Biological Invasions* 2: 255–258.
- Ivanov, V. P., Vlasenko, A., Dkhodorevskaya, R. P. & Raspopov, V. M. (1999) Contemporary status of Caspian sturgeon (Acipenseridae) stock and its conservation. *Journal* of Applied Ichthyology – Zeitschrift für Angewandte Ichthyologie 15: 103–105.

Iversen, T. M., Grant, R. & Nielsen, K. (1998) Nitrogen enrichment of European inland and marine waters with special attention to Danish policy measures. *Environmental Pollution* **102**: 771–780.

Iverson, R. L. (1990) Control of marine fish production. Limnology and Oceanography 35: 1593–1604.

Iwata, H., Shinsuke, T., Ueda, K. & Tatsukawa, R. (1995) Persistent organochlorine residues in air, water, sediments, and soils from the Lake Baikal region, Russia. *Environmental Science and Technology* 29: 792–801.

Jackson, G. (1998) Currents in the high drag environment of a coastal kelp stand off California. *Continental Shelf Research* 17: 1913–1928.

Jackson, J. B. C. (1997) Reefs since Columbus. *Coral Reefs* 16: S23–S32.

Jackson, J. B. C., Kirby, M. X., Berger, W. H., et al. (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293: 629–637.

Jackson, N. L. & Nordstrom, K. F. (1993) Depth of sediment activation by plunging waves on a steep sand beach. *Marine Geology* 115: 143–151.

Jackson, N. L., Nordstrom, K. F. & Smith, D. R. (2002) Geomorphic–biotic interactions on sandy estuarine beaches. *Journal of Coastal Research* Special Issue **36**: 414–424.

Jacobs, S. S. & Comiso, J. C. (1993) A recent sea-ice retreat west of the Antarctic Peninsula. *Geophysical Research Letters* 20: 1171–1174.

Jagtap, T. G., Komarpant, D. S. & Rodrigues, R. (2003) The seagrasses of India. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 113–120. Berkeley, CA, USA: University of California Press.

Jaksic, F. M. (1998) The multiple facets of El Niño Southern Oscillation in Chile. *Revista Chilena de Historia Natural* 71: 121–131.

Jaramillo, C. & Bayona, G. (2000) Mangrove distribution during the Holocene in Tribugá Gulf, Colombia. *Biotropica* 32: 14–22.

Jefferies, R. L. (1997) Long-term damage to sub-arctic coastal ecosystems by geese: ecological indicators and measures of ecosystem dysfunction. In: *Disturbance* and Recovery in Arctic Lands: An Ecological Perspective, ed. R. M. M. Crawford, pp. 151–166. Boston, MA, USA: Kluwer.

Jefferies, R. L. & Perkins, N. (1977) The effects on the vegetation of the additions of inorganic nutrients to salt marsh soils at Stiffkey, Norfolk. *Journal of Ecology* 65: 847–865. Jefferies, R. L., Henry, H. & Abraham, K. F. (2002) Agricultural nutrient subsidies to migratory geese and ecological change to Arctic coastal habitats. In: *Food Webs at the Landscape Level*, ed. G. A. Polis, M. E. Power & G. R. Huxel, pp. 268–283. Chicago, IL, USA: University of Chicago Press.

Jeglum, J. K. (1990) Peatland forestry in Canada: an overview. In: Biomass Production and Element Fluxes in Forested Peatland Ecosystems, ed. B. Hanell, pp. 19–28. Umeå, Sweden: Swedish University of Agricultural Science.

Jellison, R. & Melack, J. M. (1993) Algal photosynthetic activity and its response to meromixis in hypersaline Mono Lake, California. *Limnology and Oceanography* 38: 818–837.

Jenkins, M., Green, R. E. & Madden, J. (2003) The challenge of measuring global change in wild nature: are things getting better or worse? *Conservation Biology* **17**: 20–23.

Jenkins, S. R., Arenas, F., Arrontes, J., et al. (2001) European-scale analysis of seasonal variability in limpet grazing activity and microalgal abundance. *Marine Ecology Progress Series* 211: 193–203.

Jennings, S. & Kaiser, M. J. (1998) The effects of fishing on marine ecosystems. In *Advances in Marine Biology*, vol. 34, ed. J. Blaxter, A. J. Southward & P. A. Tyler, pp. 201–352. San Diego, CA, USA: Academic Press.

Jennings, S. & Lock, J. M. (1996) Population and ecosystem effects of fishing. In: *Reef Fisheries*, ed. N. V. C. Polunin & C. M. Roberts, pp. 193–218. London, UK: Chapman & Hall.

Jennings, S., Dinmore, T. A., Duplisea, D. E., Warr, K. J. & Lancaster, J. E. (2001) Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology* 70: 459–475.

Jennings, S., Greenstreet, S. P. R. & Reynolds, J. D. (1999) Structural change in an exploited fish community: a consequence of differential fishing effects on species with contrasting life histories. *Journal of Animal Ecology* 68: 617–627.

Jennings, S., Reynolds, J. D. & Mills, S. C. (1998) Life history correlates of responses to fisheries exploitation. *Proceedings* of the Royal Society of London B 265: 333–339.

Jeppesen, E. & Sammalkorpi, I. (2002) Lakes. In: Handbook of Restoration Ecology, vol. 2, ed. Restoration in Practice, ed. M. Perrow & T. Davy, pp. 297–324. Cambridge, UK: Cambridge University Press.

Jeppesen, E., Christoffersen, K., Landkildehus, F., Lauridsen, T. & Amsinck, S. (2001) Fish and crustaceans in northeast Greenland lakes with special emphasis on interactions between Arctic charr (*Salvelinus alpinus*), *Lepidurus arcticus* and benthic chydorids. *Hydrobiologia* 442: 329–337. Jeppesen, E., Jensen, J. P., Jensen, C., *et al.* (2003) The impact of nutrient state and lake depth on top-down control in the pelagic zone of lakes: study of 466 lakes from the temperate zone to the Arctic. *Ecosystems* 6: 313–325.

Jeremiason, J., Hornbuckle, K. & Eisenreich, S. (1994) PCBs in Lake Superior, 1978–1992: decreases in water concentrations reflect loss by volatilization. *Environmental Science and Technology* 28: 903–914.

JGOFS (2008) Joint Global Ocean Flux Study: home page [www document]. URL http://ijgofs.whoi.edu

Ji, C. Y., Liu, Q., Sun, D., et al. (2001) Monitoring urban expansion with remote sensing in China. International Journal of Remote Sensing 22: 1441–1455.

Jickells, T. D. (1998) Nutrient biogeochemistry of the coastal zone. Science 281: 217–222.

Joabsson, A. & Christensen, T. R. (2001) Methane emissions from wetlands and their relationship with vascular plants: an Arctic example. *Global Change Biology* 7: 919–932.

Joabsson, A., Christensen, T. R. & Wallén, B. (1999) Vascular plant controls on methane emissions from northern peat forming wetlands. *Trends in Ecology and Evolution* 14: 385–388.

Johannesburg Summit (2002) World summit on sustainable development, Johannesburg, 26 August–4 September 2002. UN News Service [www document]. URL http://www.un. org/events/wssd/

Johannessen, O. M., Miles, M. & Bjørgo, E. (1995) The Arctic's shrinking sea ice. *Nature* 376: 126–127.

Johannessen, O. M., Shalina, E. & Miles, M. W. (1999) Satellite evidence for and Arctic sea ice cover in transition. *Science* 286: 1937–1939.

Johannesson, K. & Warmoes, T. (1990) Rapid colonisation of Belgian breakwaters by the direct developer *Littorina* saxatilis (Olivi) (Prosobranchia, Mollusca). *Hydrobiologia* 193: 99–108.

Johns Hopkins (1998) Solutions for a Water-Short World, Population Report Vol. XXVI, No. 1. Baltimore, MD, USA: Johns Hopkins Population Information Program. Available online at www.infoforhealth.org/pr/m14edsum. shtml

Johnson, L. C. & Damman, A. W. H. (1993) Decay and its regulation in *Sphagnum* peatlands. *Advances in Bryology* 5: 249–296.

Johnson, L. C., Damman, A. W. H. & Malmer, N. (1990) Sphagnum macrostructure as an indicator of decay and compaction in peat cores from an ombrotrophic South Swedish peat bog. Journal of Ecology 78: 633–647. Johnson, M. P., Burrows, M. T. & Hawkins, S. J. (1998) Individual based simulations of the direct and indirect effects of limpets on a rocky shore *Fucus* mosaic. *Marine Ecology Progress Series* 169: 179–188.

Johnson, N., Revenga, C. & Echeverria, J. (2001) Managing water for people and nature. *Science* 292: 1071–1072.

Johnson, T., Scholz, C., Talbot, M., et al. (1996) Late Pleistocene desiccation of Lake Victoria and rapid evolution of cichlid fishes. *Science* 273: 1091–1093.

Johnston, E. L. & Keough, M. J. (2003) Competition modifies the response of organisms to toxic disturbance. *Marine Ecology Progress Series* 215: 15–26.

Jokiel, P. L., Hunter, C. L., Taguchi, S. & Watarai, L. (1993) Ecological impact of a fresh-water 'reef kill' in Kaneohe bay, Oahu, Hawaii. *Coral Reefs* 12: 177–184.

Jolly, I. D. (1996) The effects of river management on the hydrology and hydroecology of arid and semi-arid floodplains. In: *Floodplain Processes*, ed. M. G. Anderson, D. E. Walling & P. D. Bates, pp. 577–609. Chichester, UK: John Wiley.

Jonasson, S. & Chapin III, F. S. (1985) Significance of sequential leaf development for nutrient balance of the cotton sedge, *Eriophorum vaginatum* L. *Oecologia* 67: 511–518.

Jones, C. G., Lawton, J. H. & Shachak, M. (1994) Organisms as ecosystem engineers. *Oikos* 69: 373–386.

Jones, D., Cocklin, C. & Cutting, M. (1995) Institutional and landowner perspectives on wetland management in New Zealand. *Journal of Environmental Management* 45: 143–161.

Jones, D. A., Watt, I., Woodhouse, T. D. & Richmond, M. D. (1994) Intertidal recovery in the Dawhat ad Dafi and Dawhat al Mussallamiya region (Saudi Arabia) after the Gulf War oil spill. *Courier Forschungsinstitut Senckenberg* **166**: 27–33.

Jones, D. K. C. (1993) Global warming and geomorphology. Geographical Journal 159: 124–130.

Jones, G. (1994) Global warming, sea level change and the impact on estuaries. *Marine Pollution Bulletin* 28: 7–14.

Jones, G. P. & Andrew, N. L. (1990) Herbivory and patch dynamics on rocky reefs in temperate Australasia: the roles of fish and sea urchins. *Australian Journal of Ecology* 15: 505–520.

Jones, G. P. & Andrew, N. L. (1993) Temperate reefs and the scope of seascape ecology. In: *The Ecology of Temperate Reefs, Proceedings of the 2nd International Temperate Reef Symposium*, Auckland, pp. 63–76. Wellington, New Zealand: NIWA Publications. Jones, G. P. (1988) Ecology of rocky reef fish of north-eastern New Zealand. New Zealand Journal of Marine and Freshwater Research 22: 445–462.

Jones, N. S. (1946) Browsing of Patella. Nature 158: 557.

Jones, N. S. & Kain, J. M. (1967) Subtidal algal colonization following the removal of *Echinus*. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 15: 160–166.

Jonsson, M. & Malmqvist, B. (2000) Ecosystem process rate increases with animal species richness: evidence from leaf-eating, aquatic insects. *Oikos* 89: 519–523.

Jonsson, M. (2003*a*) Mechanisms behind diversity effects on ecosystem functioning: testing the facilitation and interference hypotheses. *Oecologia* 134: 554–559.

Jonsson, M. (2003b) Importance of species identity and number for process rates within different functional feeding groups. *Journal of Animal Ecology* 72: 453–459.

Jonsson, M., Dangles, O., Malmqvist, B. & Guérold, F. (2002) Simulating species loss following disturbance: assessing the effects on process rates. *Proceedings of the Royal Society of London B* 269: 1047–1052.

Jonsson, M., Malmqvist, B. & Hoffsten, P. O. (2001) Leaf litter breakdown rates in boreal streams: does shredder species richness matter? *Freshmater Biology* 46: 161–172.

Jorgensen, S., Riccardo, B., Ballatore, T. & Muhandiki, V. (2003) Lake Watch 2003: the changing state of the world's lakes. Unpublished report. Kusatsu, Japan: International Environment Committee Foundation.

Jumars, P. A. (1981) Limits in predicting and detecting benthic community responses to manganese nodule mining. *Marine Mining* 3: 213–229.

Junk, W. J. & da Silva, V. M. F. (1997) Mammals, reptiles and amphibians. In: *The Central Amazon Floodplain: Ecology of a Pulsing System*, ed. W. J. Junk, pp. 409–417. Berlin, Germany: Springer-Verlag.

Junk, W. J. & Wantzen, K. M. (2004) The flood pulse concept: new aspects, approaches, and applications – an update. In: *Proceedings of the 2nd Large River Symposium (LARS)*, ed. R. Welcomme & C. Barow, pp. 117–149. Phnom Penh, Cambodia: Mekong River Commission and Food and Agriculture Organization of the United Nations.

Junk, W. J. (1993) Wetlands of tropical South America. In: Wetlands of the World, vol. 1, Inventory, Ecology and Management, ed. D. F. Whigham, D. Dykyjová & S. Hejný, pp. 679–739. Dordrecht, the Netherlands: Kluwer. Junk, W. J. (1995) Human impact on neotropical wetlands: historical evidence, actual status and perspectives. *Scientia Guianae* 5: 299–311.

Junk, W. J., ed. (1997) The Central Amazon Floodplain: Ecology of a Pulsing System. Berlin, Germany: Springer-Verlag.

Junk, W. J. (2000) Mechanisms for development and maintenance of biodiversity in neotropical floodplains. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, vol. 1, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 119–139. Leiden, the Netherlands: Backhuys.

Junk, W. J. (2002) Long-term environmental trends and the future of tropical wetlands. *Environmental Conservation* 29: 414–435.

Junk, W. J. & Welcomme, R. L. (1990). Floodplains. In: Wetlands and Shallow Continental Water Bodies, ed.
B. C. Patten et al. pp. 491–524. The Hague, the Netherlands: SPB Academic Publishing.

Junk, W. J., Bayley, P. B. & Sparks, R. E. (1989) The flood pulse concept in river–floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences* 106: 110–127.

Junk, W. J., Brown, M., Campbell, I. S., et al. (2006) Comparative biodiversity of large wetlands: a synthesis. Aquatic Sciences 68: 254–277.

Junk, W. J., Ohly, J. J., Piedade, M. T. F. & Soares, M. G. M., eds. (2000) The Central Amazon Floodplain: Actual Use and Options for a Sustainable Management. Leiden, the Netherlands: Backhuys.

Jutila, H. M. (1997) Vascular plant species richness in grazed and ungrazed coastal meadows, SW Finland. *Annales Botanici Fennici* 34: 245–263.

Kabat, P., Schulze, R. E., Hellmuth, M. E. & Veraart, J. A., eds. (2002) Coping with impacts of climate variability and climate change in water management: a scoping paper. Unpublished report. Wageningen, the Netherlands: Dialogue on Water and Climate.

Kaczmarek, Z., Arnell, N. W., Stakhiv, E. Z., et al. (1996)
Water resources management. In: Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change – Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change, ed. R. T. Watson, M. C. Zinoyowera & R. H. Moss, pp. 469–486. Cambridge, UK: Cambridge University Press.

Kaczynski, V. M. & Fluharty, D. L. (2002) European policies in West Africa: who benefits from fisheries agreements? *Marine Policy* 26: 75–93. Kadioğlu, M., Şen, Z. & Batur, E. (1997) The greatest soda-water lake in the world and how it is influenced by climatic change. *Annales Geophysicae: Atmospheres, Hydrospheres and Space Sciences* 15: 1489–1497.

Kain, J. M. (1975) Algal recolonization of some cleared subtidal areas. *Journal of Ecology* 63: 739–765.

Kaiser, M. J. & de Groot, S. J., eds. (2000) *The Effects of Fishing on Non-Target Species and Habitats*. Oxford, UK: Blackwell Science.

Kaiser, M. J., Ramsay, K., Richardson, C. A., Spence, F. E. & Brand, A. R. (2000) Chronic fishing disturbance has changed shelf sea benthic community structure. *Journal of Animal Ecology* 69: 494–503.

Kania, H. J. & O'Hara, J. (1974) Behavioral alterations in a simple predator-prey system due to sub-lethal exposure to mercury. *Transactions of the American Fisheries Society* 103: 134–136.

Kannan, K., Smith Jr, R. G., Lee, R. F., et al. (1998) Distribution of total mercury and methylmercury in water, sediment, and fish from South Florida estuaries. Archives of Environmental Contamination and Toxicology 34: 109–118.

Kapuscinski, A. M. & Hallerman, E. R. (1991) Implications of introduction of transgenic fish into natural ecosystems. *Canadian Journal of Fisheries and Aquatic Science* 48: 99–107.

Karl, D. M. (1999) A sea of change: biogeochemical variability in the North Pacific Subtropical Gyre. *Ecosystems* 2: 181–214.

Karl, D. M., Bidigare, R. R. & Letelier, R. M. (2001) Long-term changes in plankton community structure and productivity in the North Pacific Subtropical Gyre: the domain shift hypothesis. *Deep-Sea Research* II 48: 1449–1470.

Karlson, K., Rosenberg, R. & Bonsdorff, E. (2002) Temporal and spatial large-scale effects of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic waters: a review. *Oceanography and Marine Biology Annual Review* 40: 427–489.

Karlsson, H., Muir, D., Teixiera, C., et al. (2000) Persistent chlorinated pesticides in air, water, and precipitation from the Lake Malawi area, Southern Africa. Environmental Science and Technology 34: 4490–4495.

Karoly, D., Risbey, J. & Reynolds, A. (2003) Global warming contributes to Australia's worst drought. WWF Australia [www document]. URL http://www.wwf.org.au/News_ and_information/Publications/PDF/Report/drought_ report.pdf. Karr, J. R. (1991) Biological integrity: a long-neglected aspect of water resource management. *Ecological Applications* 1: 66–84.

Karr, J. R., Fausch, K. D., Angermeier, P. L., Yant, P. R. & Schlosser, I. J. (1986) Assessing the Biological Integrity of Running Waters: A Method and Its Rationale. Champaign, IL, USA: Illinois Natural History Survey.

Kathiresan, K. & Bingham, B. L. (2001) Biology of mangroves and mangrove ecosystems. *Advances in Marine Biology* 40: 81–251.

Katunin, D. N. (2000) Trans-boundary diagnostic analysis of relevantly important commercial bioresources. Caspian Environment Programme [www document]. URL http:// www.caspianenvironment.org/report_miscell3.htm.

Kaufman, L. (1992) Catastrophic change in species-rich freshwater ecosystems. *BioScience* 42: 846–858.

Kautsky, N., Kautsky, H., Kautsky, U. & Waern, M. (1986) Decreased depth penetration of *Fucus vesiculosus* (L.) since the 1940s indicates eutrophication of the Baltic Sea. *Marine Ecology Progress Series* 28: 1–8.

Kawall, H. G., Torres, J. J. & Geiger, S. P. (2001) Effects of the ice-edge bloom and season on the metabolism of copepods in the Weddell Sea, Antarctica. *Hydrobiologia* 453: 67–77.

Keddy, P. A. (1981) Experimental demography of a dune annual: *Cakile edentula* growing along an environmental gradient in Nova Scotia. *Journal of Ecology* 69: 615–630.

Keddy, P. A. (2000) Wetland Ecology: Principles and Conservation. Cambridge, UK: Cambridge University Press.

Keddy, P. A. & Wisheu, I. C. (1989) Ecology, biogeography, and conservation of coastal plain plants: some general principles from the study of Nova Scotian wetlands. *Rhodora* 91: 72–94.

Keefer, D. K., Defrance, S. D., Moseley, M. E., *et al.* (1998) Early maritime economy and El Niño Events at Quebrada Tacahuay, Peru. *Science* 281: 1833–1835.

Keenan, C. P. (1994) Recent evolution of population structure in Australian barramundi, *Lates calcarifer* (Bloch): an example of isolation by distance in one dimension. *Australian Journal of Marine and Freshwater Research* 45: 1123–1148.

Kelleher, G., Bleakley, C. & Wells, S. (1995) A Global Representative System of Marine Protected Areas, vol. 1. Washington, DC, USA: World Bank.

Kelly, L. C., Rundle, S. D. & Bilton, D. T. (2002) Genetic population structure and dispersal in Atlantic Island caddisflies. *Freshmater Biology* 47: 1642–1650. Kelsey, K. A. & West, S. D. (1998) Riparian wildlife. In: *River Ecology and Management: Lessons from the Pacific Coastal Ecoregion*, ed. R. J. Naiman & R. E. Bilby, pp. 235–258. New York, USA: Springer-Verlag.

Kendall, M. A., Bowman, R. S., Williamson, P. & Lewis, J. R. (1985) Annual variation in recruitment of *Semibalanus* balanoides on the North Yorkshire coast 1969–1981. *Journal* of the Marine Biological Association of the United Kingdom 65: 1009–1030.

Kendrick, G. A., Eckersley, J. & Walker, D. I. (1999) Landscape scale changes in seagrass distribution over time: a case study from Success Bank, Western Australia. *Aquatic Botany* 65: 293–309.

Kennish, M. J. (1986) *Ecology of Estuaries: Physical and Chemical Aspects.* Boca Raton, FL, USA: CRC Press.

Kennish, M. J. (1992) Ecology of Estuaries: Anthropogenic Effects. Boca Raton, FL, USA: CRC Press.

Kennish, M. J. ed. (1997) Practical Handbook of Estuarine and Marine Pollution. Boca Raton, FL, USA: CRC Press.

Kennish, M. J. (1998a) Pollution Impacts on Marine Biotic Communities. Boca Raton, FL, USA: CRC Press.

Kennish, M. J. (1998b) Trace metal-sediment dynamics in estuaries: pollution assessment. *Reviews in Environmental Contamination and Toxicology* 155: 69–110.

Kennish, M. J. ed. (2000) Estuary Restoration and Maintenance: The National Estuary Programme. Boca Raton, FL, USA: CRC Press.

Kennish, M. J. ed. (2001a) Practical Handbook of Marine Science, 3rd edn. Boca Raton, FL, USA: CRC Press.

Kennish, M. J. (2001b) Coastal salt marsh systems: a review of anthropogenic impacts. *Journal of Coastal Research* 17: 731–748.

Kennish, M. J. ed. (2001c) Barnegat Bay–Little Egg Harbor, New Jersey: estuary and watershed assessment. *Journal of Coastal Research* Special Issue 32.

Kennish, M. J. ed. (2002a) Impacts of motorized watercraft on shallow estuarine and coastal marine environments. *Journal* of Coastal Research Special Issue 37.

Kennish, M. J. (2002b) Sediment contaminant concentrations in estuarine and coastal marine environments: potential for remobilization by boats and personal watercraft. *Journal of Coastal Research* Special Issue 37: 150–177.

Kennish, M. J. (2002c) Environmental threats and environmental future of estuaries. *Environmental Conservation* 29: 78–107.

Kennish, M. J. & Lutz, R. A., eds. (1984) Ecology of Barnegat Bay, New Jersey. New York, USA: Springer-Verlag.

Kentula, M. E. (2000) Perspectives on setting success criteria for wetland restoration. *Ecological Engineering* 15: 199–209. Kenworthy, W. J., Fonseca, M. S., Whitfield, P. E. & Hammerstrom, K. K. (2002) Analysis of seagrass recovery in experimental excavations and propeller-scar disturbances in the Florida Keys National Marine Sanctuary. *Journal of Coastal Research* Special Issue 37: 150–177.

Kerr, R. A. (2002) A warmer Arctic means change for all. Science 297: 1490–1492.

Kestner, F. J. T. (1962) The old coastline of the Wash: a contribution to the understanding of loose boundary processes. *Geographical Journal* 128: 457–487.

Kickert, R. N., Tonella, G., Simonov, A. & Krupa, S. V. (1999) Predictive modeling of effects under global change. *Environmental Pollution* 100: 87–132.

Kidd, K., Bootsma, H., Hesslein, R., Muir, D. & Hecky, R. (2001) Biomagnification of DDT through the benthic and pelagic food webs of Lake Malawi, East Africa: importance of trophic level and carbon source. *Environmental Science* and Technology 39: 14–25.

Kiehl, K., Esselink, P. & Bakker, J. P. (1997) Nutrient limitation and plant species composition in temperate salt marshes. *Oecologia* 111: 325–330.

Kimura, M. & Weiss, G. H. (1964) The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics* 49: 561–574.

King, J. C. (1994) Recent climate variability in the vicinity of the Antarctic Peninsula. *International Journal of Climatology* 14: 357–369.

King, J. C. & Harangozo, S. A. (1998) Climate change in the western Antarctic Peninsula since 1945: observations and possible causes. *Annals of Glaciology* 27: 571–575.

King, J. M. & Louw, D. (1998) Instream flow assessment for regulated rivers in South Africa using the building block methodology. *Aquatic Ecosystem Health and Management* 1: 109–124.

Kingett, P. D. & Choat, J. H. (1981) Analysis of density and distribution patterns in *Chrysophrys auratus* (Pisces: Sparidae) within a reef environment: an experimental approach. *Marine Ecology Progress Series* 5: 283–290.

Kingsford, M. J., Underwood, A. J. & Kennelly, S. J. (1991) Humans as predators on rocky reefs in New South Wales, Australia. *Marine Ecology Progress Series* 72: 1–2.

Kingsford, R. T. (2000) Ecological impacts of dams, water diversions and river management on floodplain wetlands in Australia. *Austral Ecology* 25: 109–127.

Kingsford, R. T. & Thomas, R. F. (1995) The Macquarie Marshes in arid Australia and their waterbirds: a 50-year history of decline. *Environmental Management* 19: 867–878. Kingsford, R. T., Boulton, A. J. & Puckridge, J. T. (1998) Challenges in managing dryland rivers crossing political boundaries: lessons from Cooper Creek and the Paroo River, central Australia. *Aquatic Conservation: Marine and Freshwater Ecosystems* 8: 361–378.

Kingsley, C. (1890) *Glaucus, or the Wonders of the Shore.* London, UK: Macmillan.

Kinner, N. E., Harvey, R. W., Blakeslee, K., Novarino, G. & Meeker, L. D. (1998) Size-selective predation on groundwater bacteria by nanoflagellates in an organic-contaminated aquifer. *Applied and Environmental Microbiology* 64: 618–625.

Kirby, A. (2002) Lake level fuels climate concern. BBC News World Edition, 28 October 2002 [www document]. URL http://news.bbc.co.uk/1/hi/sci/tech/2369333.stm

Kirkby, J. & O'Keefe, P. (1998) Conservation of inland deltas: a case study of the Gash Delta, Sudan. In: *The Sustainable Management of Tropical Catchments*, ed. D. Harper & A. G. Brown, pp. 209–223. Chichester, UK: John Wiley.

Kitching, J. A. & Ebling, F. J. (1961) The ecology of Lough Ine. XI. The control of algae by *Paracentrotus lividus* (Echinoidea). *Journal of Animal Ecology* **30**: 373–383.

Kitching, J. A. & Thain, V. M. (1983) The ecological impact of the sea urchin *Paracentrotus lividus* (Lamarck) in Lough Ine, Ireland. *Philosophical Transactions of the Royal Society* of London B 300: 513–552.

Kivinen, E. & Pakarinen, P. (1981) Geographical distribution of peat resources and major peatland complex types in the world. Annales Academiae Scientiarum Fennicae, Series A, III, Geologica–Geographica 132: 1–28.

Kjerfve, B. (1989) Estuarine geomorphology and physical oceanography. In: *Estuarine Ecology*, ed. J. W. Day Jr, C. A. S. Hall, W. M. Kemp & A. Yañez-Arancibia, pp. 47–78. New York, USA: John Wiley.

Klarquist, M., Bohlin, E. & Nilsson, M. (2001) Long-term decline in apparent peat carbon accumulation in boreal mires in northern Sweden. *Acta Universitatis Agriculturae Suediae Silvestria* 203(3): 1–22.

Klein, R. J. T., Smit, M. J., Goosen, H. & Hulsbergen, C. H. (1998) Resilience and vulnerability: coastal dynamics or Dutch dikes? *Geographical Journal* 164: 259–268.

Kleypas, J. A., Buddemeier, R. W., Archer, D., et al. (1999a) Geochemical consequences of increased atmospheric carbon dioxide on coral reefs. *Science* 284: 118–120.

Kleypas, J. A., McManus, J. W. & Menez, L. A. B. (1999b) Environmental limits to coral reef development: where do we draw the line? *American Zoologist* 39: 146–159. Klijn, F. & Witte, J.-P. M. (1999) Eco-hydrology: groundwater flow and site factors in plant ecology. *Hydrogeology Journal* 7: 65–77.

Klimo, E. & Hager, H., eds. (2001) *The Floodplain Forests in Europe*. Leiden, the Netherlands: E. J. Brill.

Knapp, R., Matthews, K. R. & Sarnelle, O. (2001) Resistance and resilience of alpine lake fauna to fish introductions. *Ecological Monographs* 71: 401–421.

Knopf, F. L. & Samson, F. B. (1994) Scale perspectives on avian diversity in western riparian ecosystems. *Conservation Biology* 8: 669–676.

Knowlton, N. (1993) Sibling species in the sea. Annual Review of Ecology and Systematics 24: 189–216.

Knowlton, N. (2000) Molecular genetic analyses of species boundaries in the sea. *Hydrobiologia* 420: 73–90.

Knowlton, N., Lang, J. C. & Keller, B. D. (1988) Fates of staghorn coral fragments on hurricane-damaged reefs in Jamaica: the role of predators. In: *Proceedings of the 6th International Coral Reef Symposium*, Townsville, Australia, 8–12 August 1988, vol. 2, ed. J. H. Choat, D. Barnes, M. A. Borowitzka, *et al.*, pp. 83–88. Townsville, Queensland, Australia: 6th International Coral Reef Symposium Executive Committee.

Knutson, T. R., Tuleya, R. E. & Kurihara, Y. (1998) Simulated increase of hurricane intensities in a CO₂warmed climate. *Science* 279: 1018–1020.

Koch, E. W. & Orth, R. J. (2003) The seagrasses of the Mid-Atlantic Coast of the United States. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 234–243. Berkeley, CA, USA: University of California Press.

Kock, K.-H. (1992) Antarctic Fish and Fisheries. Cambridge, UK: Cambridge University Press.

Kock, K.-H. (2001) The direct influence of fishing and fishery-related activities on non-target species in the Southern Ocean with particular emphasis on longline fishing and its impact on albatrosses and petrels: a review. *Reviews in Fish Biology and Fisheries* 11: 31–56.

Koike, K. (1993) The countermeasures against coastal hazards in Japan. *Geojournal* 38: 301–312.

Kolar, C. S. & Lodge, D. M. (2001) Progress in invasion biology: predicting invaders. *Trends in Ecology and Evolution* 16: 199–204.

Kolpin, D. W., Furlong, E. T., Meyer, M. T., et al. (2002) Pharmaceuticals, hormones, and other organic wastewater contaminants in US streams, 1999–2000: a national reconnaissance. *Environmental Science and Technology* 36: 1202–1211. Kooijman, A. M. & de Haan, M. W. A. (1995) Grazing as a measure against grass encroachment in Dutch dry dune grassland: effects on vegetation and soil. *Journal of Coastal Conservation* 1: 127–134.

Koop, K., Booth, D., Broadbent, A. D., et al. (2001) ENCORE: the effect of nutrient enrichment on coral reefs – synthesis of results and conclusions. *Marine Pollution Bulletin* 42: 91–120.

Kosarev, A. N. & Yablonskaya, E. A. (1994) *The Caspian Sea*. The Hague, the Netherlands: SPB Academic Publishing.

Koslow, J. A. (1997) Seamounts and the ecology of deep-sea fisheries. *American Scientist* 85: 168–176.

Koslow, J. A., Bell, J., Virtue, P. & Smith, D. C. (1995) Fecundity and its variability in orange roughy: effects of population density, condition, egg size, and senescence. *Journal of Fish Biology* 47: 1063–1080.

Koslow, J. A., Boehlert, G. W., Gordon, J. D. M., et al. (2000) Continental slope and deep-sea fisheries: implications for a fragile ecosystem. *ICES Journal of Marine Science* 57: 458–557.

Koslow, J. A., Gowlett-Holmes, K., Lowry, J. K., et al. (2001) Seamount benthic macrofauna off southern Tasmania: community structure and impacts of trawling. *Marine Ecology Progress Series* 213: 111–125.

Kota, S., Borden, R. C. & Barlaz, M. A. (1999) Influence of protozoan grazing on contaminant biodegradation. *FEMS Microbiology Ecology* 29: 179–189.

Kothari, A., Pande, P., Singh, S. & Variava, D. (1989) Management of National Parks and Sanctuaries in India: A Status Report. New Delhi, India: Ministry of Environment and Forests.

Kotze, D. C. & Breen, C. M. (1994) Agricultural Land-Use Impacts on Wetland Functional Values, WRC Report No. 501/3/94. Pretoria, South Africa: Water Research Commission.

Kotze, D. C. & Breen, C. M. (2000) WETLAND-USE: A Wetland Management Decision Support System for South African Freshwater Palustrine Wetlands. Pretoria, South Africa: Department of Environmental Affairs and Tourism. Available online at www.ccwr.ac.za/wetlands/

Kozhov, M. (1963) Lake Baikal and its Life. The Hague, the Netherlands: Dr W. Junk.

Kozhova, O. & Silow, E. (1998) The current problems of Lake Baikal ecosystem. Lakes and Reservoirs: Research and Management 3: 19–33.

Kraus, N. C. & Rankin, K. L., eds. (2004) Functioning and design of coastal groins: the interaction of groins and the beach-process and planning. *Journal of Coastal Research* Special Issue 33. Krieger, K. J. (2001) Coral (*Primnoa*) impacted by fishing gear in the Gulf of Alaska. In: *Proceedings of the 1st International Symposium on Deep-Sea Corals*, ed. J. Willison, J. Hall, S. Gass, *et al.*, pp. 106–116. Halifax, NS, Canada: Ecology Action Centre.

Kroes, D. & Brinson, M. M. (2004) Occurrence of riverine wetlands on floodplains along a climatic gradient. *Wetlands* 24: 167–177.

Kroeze, C. & Seitzinger, S. P. (1998) Nitrogen inputs to rivers, estuaries and continental shelves and related nitrous oxide emissions in 1990 and 2050: a global model. *Nutrient Cycling in Agroecosystems* 52: 195–212.

Kucklick, J. R. & Bidleman, T. F. (1994) Organic contaminants in Winyah Bay, South Carolina. I. Pesticides and polycyclic aromatic hydrocarbons in subsurface and microlayer waters. *Marine Environmental Research* 37: 63–78.

Kucklick, J., Bidleman, T., McConnell, L., Walla, M. & Ivanov, G. (1994) Organochlorines in water and biota of Lake Baikal, Siberia. *Environmental Science and Technology* 28: 31–37.

Kuffner, I. B. (2001) Effects of ultraviolet radiation and water motion on the reef coral *Porites compressa* Dana: a flume experiment. *Marine Biology* 138: 467–476.

Kuhn, G. G. & Shepard, F. P. (1980) Coastal erosion in San Diego County, California. In: *Coastal Zone 80*, pp. 1899–1918. New York, USA: American Society of Civil Engineers.

Kuhnert, H., Patzold, J., Hatcher, B. G., *et al.* (1999) A 200-year coral stable oxygen isotope record from a high-latitude reef off Western Australia. *Coral Reef* 18: 1–12.

Kumar, K. K., Rajagopalan, B. & Cane, M. A. (1999) On the weakening relationship between the Indian monsoon and ENSO. *Science* 284: 2156–2159.

Kuriandewa, T. E., Kiswara, W., Hutomo, M., Soemodihardjo, S. & Hermana, I. (2003) The seagrasses of Indonesia. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 187–198. Berkeley, CA, USA: University of California Press.

Kurlansky, M. (1997) Cod: A Biography of the Fish that Changed the World. New York, USA: Walker & Co.

Kusler, J. & Larson, L. (1993) Beyond the ark: a new approach to US floodplain management. *Environment* 35: 7–35.

Kwok, R. & Comiso, J. C. (2002) Southern ocean climate and sea ice anomalies associated with the Southern Oscillation. *Journal of Climate* 15: 487–501.

- Kyle, R., Pearson, B., Fielding, P. J., Robertson, W. D. & Birnie, S. L. (1997) Subsistence shellfish harvesting in the Maputaland Marine Reserve in northern KwaZulu-Natal, South Africa: rocky shore organisms. *Biological Conservation* 82: 183–192.
- La Peyre, M. K., Mendelssohn, I. A., Reams, M. A., Templet, P. H. & Grace, J. B. (2001) Identifying determinants of nations' wetland management programs using structural equation modeling: an exploratory analysis. *Environmental Management* 27: 859–868.
- Lacambra, C., Cutts, N., Allen, J., Burd, F. & Elliott, M. (2004) Spartina anglica: A Review of its Status, Dynamics and Management., English Nature Research Report No.527. Peterborough, UK: English Nature.
- Lack, M. & Sant, G. (2001) Patagonian toothfish: are conservation and trade measures working? *TRAFFIC Bulletin* 19: 1–18.
- Lae, R. (1994) Effects of drought, dams, and fishing pressure on the fisheries of the central delta of the Niger River. *International Journal of Ecology and Environmental Sciences* 20: 119–128.
- Lamb, M. & Zimmerman, M. H. (1964) Marine vegetation of Cape Ann, Essex County, Massachusetts. *Rhodora* 66: 217–254.
- Lambert, G. & Steinke, T. D. (1986) Effects of destroying juxtaposed mussel-dominated and coralline algal communities at Umdoni Park, Natal South coast. *South African Journal of Marine Science* 4: 203–217.
- Lambert, R. A., Hotchkiss, P. F., Roberts, N., *et al.* (1990) The use of wetlands (dambos) for micro-scale irrigation in Zimbabwe. *Irrigation and Drainage Systems* 4: 17–28.
- Lambert, W. J., Levin, P. S. & Berman, J. (1992) Changes in the structure of a New England (USA) kelp bed: the effects of an introduced species? *Marine Ecology Progress Series* 88: 303–307.
- Lambs, L. & Muller, E. (2002) Sap flow and water transfer in the Garonne River riparian woodland, France: first results on poplar and willow. *Annals of Forest Science* 59: 301–315.
- Lamers, L. P. M., Smolders, A. J. P. & Roelofs, J. G. M. (2002) The restoration of fens in the Netherlands. *Hydrobiologia* 478: 107–130.
- Lamers, L. P. M., Tomassen, H. B. M. & Roelofs, J. G. M. (1998) Sulfate-induced eutrophication and phytotoxicity in freshwater wetlands. *Environmental Science and Technology* 32: 199–205.
- Lamparelli, C. C., Rodrigues, F. O. & de Moura, D. O. (1997) Long-term assessment of an oil spill in a mangrove forest in São Paulo, Brazil. In: *Mangrove Ecosystem Studies in Latin*

America and Africa, ed. B. Kjerfve, L. D. Lacerda & E. H. S. Diop, pp. 191–203. Paris, France: UNESCO.

- Lancaster, J., Real, M., Juggins, S., et al. (1996) Monitoring temporal changes in the biology of acid waters. *Freshwater Biology* 36: 179–201.
- Landesman, L. (1994) Negative impacts of coastal tropical aquaculture developments. World Aquaculture 25(2): 12–17.
- Lane, R. R., Day Jr, J. W. & Thibodeaux, B. (1999) Water quality analysis of a freshwater diversion at Caernarvon, Louisiana. *Estuaries* 22: 327–336.
- Langdon, C. (2002) Overview of experimental evidence for effects of CO₂ on calcification of reef builders. In: *Proceedings of the 9th International Coral Reef Symposium*, 23–27 October 2000, vol. 2, ed. K. M. Moosa, S. Soemodihardjo, A. Soegiarto, *et al.*, pp. 1091–1098. Bali, Indonesia: Ministry of Environment, Indonesian Institute of Sciences, International Society for Reef Studies.
- Langdon, C., Takahashi, T., Sweeney, C., Chapman, D. & Goddard, J. (2000) Effect of calcium carbonate saturation state on the calcification rate of an experimental coral reef. *Global Biogeochemical Cycle* 14: 639–654.
- Langis, R., Zalejko, M. & Zedler, J. B. (1991) Nitrogen assessments in a constructed and a natural salt marsh of San Diego Bay, California. *Ecological Applications* 1: 40–51.
- Lanier-Graham, S.D. (1993) The Ecology of War: Environmental Impacts of Weaponry and Warfare. New York, USA: Walker & Co.
- Lant, C. L., Kraft, S. E. & Gillman, K. R. (1995) The 1990 farm bill and water quality in Corn Belt watersheds: conserving remaining wetlands and restoring farmed wetlands. *Journal of Soil and Water Conservation* 50: 201–205.
- Lanyon, J. M., Limpus, C. J. & Marsh, H. (1989) Dugongs and turtles: grazers in seagrass systems. In: Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region, ed. A. W. D. Larkum, A. J. McComb & S. A. Shepherd, pp. 182–210. Amsterdam, the Netherlands: Elsevier/North Holland.
- Lapointe, B. E. (1999) Simultaneous top-down and bottom-up forces control macroalgal blooms on coral reefs. *Limnology* and Oceanography 44: 1586–1592.
- Larkum, A. W. D. & West, R. J. (1990) Long-term changes of seagrass meadows in Botany Bay, Australia. *Aquatic Botany* 37: 55–70.
- Larned, S. T., Kinzie, R. A., Covich, A. P. & Chong, C. T. (2003) Detritus processing by endemic and non-native Hawaiian stream invertebrates: a microcosm study of species-specific effects. *Archiv für Hydrobiologie* 156: 241–254.

Larson, J. S. (1992) North America. In: *Wetlands*, ed. M. Finlayson & M. Moser, pp. 57–84. Oxford, UK: Facts on File.

Laurance, W. F., Cochrane, M. A., Bergen, S., et al. (2001) The future of the Brazilian Amazon. Science 291: 438–439.

Lawrence, G. B., David, M. B., Lovett, G. M., et al. (1999) Soil calcium status and the response of stream chemistry to changing acidic deposition rates. *Ecological Applications* 9: 1059–1072.

Lawrence, J. M. (1975) On the relationships between marine plants and sea urchins. Oceanography and Marine Biology Annual Review 13: 213–286.

Lawton, J. H. (2000) Community Ecology in a Changing World. Oldendorf/Kuhe, Germany: Ecology Institute.

Le Maitre, D. L., Scott, D. F. & Colvin, C. (1999) A review of information on interactions between vegetation and groundwater. *Water SA* **25**: 137–152.

Leach, J., Mills, E. & Dochoda, M. (1999) Non-indigenous species in the Great Lakes: ecosystem impacts, binational policies, and management. In: *Great Lakes Fisheries Policy and Management*, ed. W. W. Taylor & C. P. Ferreri, pp. 185–207. East Lansing, MI, USA: Michigan State University Press.

Leach, S. J. (1988) Rediscovery of *Halimione pedunculata* (L.) Aellen in Britain. *Watsonia* 17: 170–171.

Leadbitter, D., Lee Long, W. & Dalmazzo, P. (1999)
Seagrasses and their management implications for research.
In: *Seagrass in Australia*, ed. A. Butler & P. Jernakoff,
pp. 140–171. Canberra, ACT, Australia: CSIRO.

Leaman, B. M. & Beamish, R. J. (1984) Ecological and management implications of longevity in some Northeast Pacific groundfishes. *International North Pacific Fisheries Commission Bulletin* 42: 85–97.

Lean, G. (2003) 'Ecological meltdown': huge dust cloud threatens Asia. *The Independent* 26 January.

Lear, C. H., Elderfield, H. & Wilson, P. A. (2000) Cenozoic deep-sea temperatures and global ice volumes from Mg/Ca in benthic foraminiferal calcite. *Science* **287**: 269–272.

Leatherland, T. M., Burton, J. D. C. F., McCartney, M. J. & Morris, R. J. (1973) Concentrations of some trace metals in pelagic organisms and of mercury in Northeast Altantic Ocean water. *Deep-Sea Research* 20: 679–685.

Ledger, M. E. & Hildrew, A. G. (2001) Recolonization by the benthos of an acid stream following a drought. *Archiv für Hydrobiologie* 152: 1–17.

Ledger, M. E. & Hildrew, A. G. (2005) The ecology of acidification and recovery: changes in herbivore–algal food web linkages across a stream pH gradient. *Environmental Pollution* 137: 103–118. Ledley, T. S. (1993) Variations in snow on sea ice: a mechanism for producing climate variations. *Journal of Geophysical Research* 98: 10401–10410.

Ledley, T. S., Sundquist, E. T., Schwartz, S. E., *et al.* (1999) Climate change and greenhouse gases. *Eos* 80: 453–458.

Lee, J. A. (1998) Unintentional experiments with terrestrial ecosystems: ecological effects of sulphur and nitrogen pollutants. *Journal of Ecology* 86: 1–12.

Lee, K.-S. & Lee, S. Y. (2003) The seagrasses of the Republic of Korea. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 211–216. Berkeley, CA, USA: University of California Press.

Lee, S. Y. & Kwok, P. W. (2002) The importance of mangrove species association to the population biology of the sesarmine crabs, *Perasesarma affinis* and *Parisesarma bidens*. *Wetland Ecology and Management* 10: 215–226.

Legendre, L., Ackley, S. F., Dieckmann, G. S., et al. (1992) Ecology of sea ice biota. II. Global significance. Polar Biology 12: 429–444.

Leggett, D., Bubb, J. C. & Lester, J. N. (1995) The role of pollutants and sedimentary processes in flood defence: a case study – salt marshes on the Essex coast, UK. *Environmental Toxicology* 16: 457–466.

Lehmusluoto, P., Machbub, B., Terangna, N., et al. (1999) Limnology in Indonesia. In: Limnology in Developing Countries, vol. 2, ed. R. G. Wetzel & B. Gopal, pp. 119–234. New Delhi, India: International Association of Theoretical and Applied Limnology.

Leighton, D. (1998) Control of sabellid infestation in green and pink abalone, *Haliotis fulgens* and *H. corrugata*, by exposure to elevated water temperatures. *Journal of Shellfish Research* 17: 701–705.

Leighton, D. L., Jones, L. G. & North, W. (1966) Ecological relationships between giant kelp and sea urchins in southern California. In: *Proceedings of the 5th International Seaweed Symposium*, ed. E. G. Young & J. L. McLachlan, pp. 141–153. Oxford, UK: Pergamon Press.

Leiva, G. E. & Castilla, J. C. (2001) A review of the world marine gastropod fishery: evolution of catches, management and the Chilean experience. *Reviews in Fish Biology and Fisheries* 11: 283–300.

Leland, A. (2002) A new apex predator in the Gulf of Maine: large, mobile Jonah crabs (*Cancer borealis*) control benthic community structure. M.Sc. thesis, University of Maine, Orono, ME, USA.

Lemmens, J. W. T. J, Clapin, G., Lavery, P. & Cary, J. (1996) Filtering capacity of seagrass meadows and other habitats of Cockburn Sound, Western Australia. *Marine Ecology Progress Series* 143: 187–200.

Lenihan, H. S., Micheli, F., Shelton, S. W. & Peterson, C. H. (1999) The influence of multiple environmental stressors on susceptibility to parasites: an experimental determination with oysters. *Limnology and Oceanography* 44: 910–924.

Leonard, G. (2000) Latitudinal variation in species interactions: a test in the New England rocky intertidal zone. *Ecology* 81: 1015–1030.

Leopold, A. (1924) Canada: journal entries dated June 11–25. Republished in Leopold, L. B., ed. (1953) *Round River: From the Journals of Aldo Leopold*. Oxford, UK: Oxford University Press.

Leopold, A. (1939) The farmer as conservationist. American Forests 45: 294–299.

Leopold, A. (1941) Lakes in relation to terrestrial life patterns. In: A Symposium on Hydrobiology, ed. J. G. Needham, pp. 17–22. Madison, WI, USA: University of Wisconsin Press.

Leopold, L. B., Wolman, M. G. & Miller, J. P. (1964) Fluvial Processes in Geomorphology. San Francisco, CA, USA: W. H. Freeman.

Lessios, H. A. (1988) Mass mortality of *Diadema antillarum* in the Caribbean: what have we learned? *Annual Review of Ecology and Systematics* **19**: 371–393.

Lessios, H. A., Garrido, M. J. & Kessing, B. D. (2001) Demographic history of *Diadema antillarum*, a keystone herbivore on Caribbean reefs. *Proceedings of the Royal Society of London B* 268: 1–7.

Letolle, R. & Mainguet, M. (1993) *L'Aral.* Paris, France: Springer-Verlag.

Levin, L. A. & Gooday, A. J. (2003) The deep Atlantic Ocean. In: *Ecosystems of the World*, vol. 28, *Ecosystems of the Deep Oceans*, ed. P. A. Tyler, pp. 111–178. Amsterdam, the Netherlands: Elsevier.

Levin, L. A., Blair, N. E., Martin, C. M., et al. (1999) Macrofaunal processing of phytodetritus at two sites on the Carolina margin: *in situ* experiments using C-13 labeled diatoms. *Marine Ecology Progress Series* 182: 37–54.

Levin, P. S. (1994) Fine-scale temporal variation in recruitment of a temperate demersal fish: the importance of settlement versus post-settlement loss. *Oecologia* **97**: 124–133.

Levin, P. S. & Hay, M. E. (1996) Responses of temperate reef fishes to alterations in algal structure and species composition. *Marine Ecology Progress Series* 134: 37–47.

Levin, P. S., Coyer, J. A., Petrik, R. & Good, T. P. (2002) Community-wide effects of non-indigenous species on temperate rocky reefs. *Ecology* 83: 3182–3193. Levin, R. B., Epstein, P. R., Ford, T. E., et al. (2002) US drinking water challenges in the twenty-first century. *Environmental Health Perspectives* 110 (Suppl. 1): 43–52.

Levitus, S., Antonov, J. I., Boyer, T. P. & Stephens, C. (2000) Warming of the world ocean. *Science* 287: 2225–2229.

Levitus, S., Antonov, J. I., Wang, J., et al. (2001) Anthropogenic warming of Earth's climate systems. Science 292: 267–270.

Lewis, J. R. (1964) *The Ecology of Rocky Shores*. London, UK: English Universities Press.

Lewis, J. R. (1976) Long-term ecological surveillance: practical realities in the rocky littoral. *Oceanography and Marine Biology Annual Review* 14: 371–390.

Lewis, J. R. (1996) Coastal benthos and global warming: strategies and problems. *Marine Pollution Bulletin* 32: 698–700.

Lewis, S. A. (1986) The role of herbivorous fishes in the organization of a Caribbean reef community. *Ecological Monographs* 56: 183–200.

Li, M. S. & Lee, S. Y. (1997) Mangroves of China: a brief review. Forest Ecology and Management 96: 241–259.

Liddle, M. J. & Greig-Smith, P. (1975) A survey of tracks and paths in a sand dune ecosystem. *Journal of Applied Ecology* 12: 893–908.

Likens, G. E. & Bormann, F. H. (1974) Acid rain: a serious regional environmental problem. *Science* 184: 1176–1179.

Lima, F. P., Queiroz, N., Ribeiro, P. A., Hawkins, S. J. & Santos, A. M. (2006) Recent changes in the distribution of a marine gastropod, *Patella rustica* Linnaeus, 1758, and their relationship to unusual climatic events. *Journal of Biogeography* 33: 812–822.

Lindeman, K. C. & Snyder, D. B. (1999) Nearshore hardbottom fishes of southeast Florida and effects of habitat burial caused by dredging. *Fishery Bulletin* 97: 508–525.

Lindig-Cisneros, R., Desmond, J., Boyer, K. & Zedler, J. B. (2003) Wetland restoration thresholds: can a degradation transition be reversed with increased effort? *Ecological Applications* 13: 193–205.

Linley, E. A.S., Newell, R. C. & Bosma, S. A. (1981) Hetertrophic utilization of mucilage released during fragmentation of Kelp (*Ecklonia maxima* and *Laminaria pallida*) development of microbial communities associated with the degradation of kelp mucilage. *Marine Ecology Progress Series* 4: 31–41.

Lipcius, R. N. & Stockhausen, W. T. (2002) Concurrent decline of spawning stock, recruitment, larval abundance and size of the blue crab *Callinectes sapidus* in Chesapeake Bay. *Marine Ecology Progress Series* 226: 45–61. Lipkin, Y. (1975) *Halophila stipulacea*, a review of a successful immigration. *Aquatic Botany* 1: 203–215.

Lipkin,Y., Beer S. & Zakai, D. (2003) The seagrasses of the Eastern Mediterranean and the Red Sea. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 75–83. Berkeley, CA, USA: University of California Press.

Little, C. & Kitching, J. A. (1996) *The Biology of Rocky Shores*. Oxford, UK: Oxford University Press.

Little, C. & Mettam, C. (1994) Rocky shore zonation in the Rance tidal power basin. *Biological Journal of the Linnean Society* 51: 169–182.

Littler, M. M. & Littler, D. S. (1980) The evolution of thallus form and survival strategies in benthic marine macroalgae: field and laboratory tests of a functional form model. *American Naturalist* 116: 25–44.

Littler, M. M., Littler, D. S. & Titlyanov, E. A. (1991) Comparisons of N- and P-limited productivity between high granitic islands versus low carbonate atolls in the Seychelles Archipelago: a test of the relative-dominance paradigm. *Coral Reefs* 10: 199–209.

Littler, M. M., Martz, D. R. & Littler, D. (1983) Effects of recurrent sand deposition on rocky intertidal organisms: the importance of substrate heterogeneity in a fluctuating environment. *Marine Ecology Progress Series* 11: 129–139.

Livingston, R. J. (1996) Eutrophication in estuaries and coastal systems: relationships of physical alterations, salinity stratification, and hypoxia. In: *Sustainable Development in the Southeastern Coastal Zone*, ed. F. J. Vernberg, W. B. Vernberg & T. Siewicki, pp. 285–318. Columbia, SC, USA: University of South Carolina Press.

Livingston, R. J. (1997) Trophic response of estuarine fishes to long-term changes of river runoff. *Bulletin of Marine Science* 60: 984–1004.

Livingston, R. J. (2000) Eutrophication Processes in Coastal Systems: Origin and Succession of Plankton Blooms and Secondary Production in Gulf Coast Estuaries. Boca Raton, FL, USA: CRC Press.

Livingston, R. J. (2001) Management of the Apalachicola River Estuary: Relationships of River Flow and River-Bay Productivity, Technical Report. Tallahassee, FL, USA: Florida State University.

Livingston, R. J. (2002) Trophic Organization in Coastal Systems: Natural Food Web Processes and Effects of Natural Loading, Plankton Blooms, and Toxic Substances. Boca Raton, FL, USA: CRC Press.

Livingston, R. J., Niu, X.-F., Lewis, F. G. & Woodsum, G. C. (1997) Freshwater input to a Gulf estuary: long-term control of trophic organization. *Ecological Applications* 7: 277–299. Llewellyn, D. W., Shaffer, G. P., Craig, N. J., et al. (1996) A decision-support system for prioritizing restoration sites on the Mississippi River alluvial plain. *Conservation Biology* 10: 1446–1455.

Locarnini, S. J.P. & Presley, B. J. (1996) Mercury concentrations in benthic organisms from a contaminated estuary. *Marine Environmental Research* **41**: 225–239.

Loeb, V., Siegel, V., Holm-Hansen, O., *et al.* (1997) Effects of sea-ice extent and krill or salp dominance on the Antarctic food web. *Nature* 387: 897–900.

Löffler, M. & Coosen, J. (1995) Ecological impact of sand replenishment. In: *Directions in European Coastal Management*, ed. M. G. Healy & P. J. Doody, pp. 291–299. Cardigan, UK: Samara Publishing.

Lofgren, B., Quinn, F., Clites, A., Assel, R. & Eberhardt, A. (2000) Impacts, challenges and opportunities: water resources, Great Lakes. In: *Preparing for a Changing Climate*, ed. P. J. Sousounis & J. M. Bisanz, pp. 29–37. Washington, DC, USA: US Environmental Protection Agency.

Lomborg, B. (2001) The Skeptical Environmentalist: Measuring the Real State of the World. Cambridge, UK: Cambridge University Press.

Long, E. R., Robertson, A., Wolfe, D. A., Hameedi, J. & Sloane, G. M. (1996) Estimates of the spatial extent of sediment toxicity in major US estuaries. *Environmental Science and Technology* **30**: 3585–3592.

Longhurst, A. R. (1998) *Ecological Geography of the Sea*. San Diego, CA, USA: Academic Press.

Longhurst, A., Sathyendranath, S., Platt, T. & Caverhill, C. (1995) An estimate of global primary production in the ocean from satellite radiometer data. *Journal of Plankton Research* 17: 1245–1271.

Looney, P. B. & Gibson, D. J. (1993) Vegetation monitoring of beach nourishment. In: *Beach Nourishment: Engineering and Management Considerations*, ed. D. K. Stauble & N. C. Kraus, pp. 226–241. New York, USA: American Society of Civil Engineers.

Lorance, P. (1998) Structure du peuplement ichthyologique du talus continental à l'ouest des Iles Britanniques et impact de la pêche. *Cybium* **22**: 209–231.

Lord, D. A., Paling, E. & Gordon, D. G. (1999) Review of seagrass rehabilitation and restoration programs in Australia. In: *Seagrass in Australia*, ed. A. Butler & P. Jernakoff, pp. 65–139. Canberra, ACT, Australia: CSIRO.

Loreau, M., Naeem, S., Inchausti, P., et al. (2001) Biodiversity and ecosystem functioning: current knowledge and future challenges. Science 294: 804–808. Loreau, M., Naeem, S. & Inchausti, P., eds. (2002) Biodiversity and Ecosystem Functioning: Synthesis and Perspectives. Oxford, UK: Oxford University Press.

Lough, J. & Barnes, D. (1997) Several centuries of variation in skeletal extension, density and calcification in massive *Porites* colonies from the Great Barrier Reef: a proxy for seawater temperature and a background of natural variability against which to identify unnatural change. *Journal of Experimental Marine Biology and Ecology* 211: 29–67.

Løvas, S. M. & Tørum, A. (2001) Effect of the kelp Laminaria hyperborea upon sand dune erosion and water particle velocities. Coastal Engineering 44: 37–63.

Lowe-McConnell, R. H., ed. (1987) *Ecological Studies in Tropical Fish Communities*. New York, USA: Cambridge University Press.

Lowe-McConnell, R. (1994) The changing ecosystem of Lake Victoria, East Africa. *Freshwater Forum* **4**: 76–89.

Loya, Y. & Rinkevich, B. (1980) Effects of oil pollution on coral reef communities. *Marine Ecology Progress Series* 3: 167–180.

Lu, J. (1995) Ecological significance and classification of Chinese wetlands. *Vegetatio* 118: 49–56.

Lubchenco, J., Olson, A. M., Brubaker, L. B., *et al.* (1991) The sustainable biosphere initiative: an ecological research agenda. *Ecology* **72**: 371–412.

Lubin, D., Booth, C. R., Lucas, T., Neuschuler, D. & Frederick, J.E (1989) Measurements of enhanced springtime ultraviolet radiation at Palmer Station, Antarctica. *Geophysical Research Letters* **16**: 783–785.

Ludgate, J. W. (1987) The economic and technical impact of TBT legislation on the USA marine industry. In: Oceans '87 Proceedings, vol. 4, International Organotin Symposium, pp. 1309–1313. New York, USA: Institute of Electrical and Electronic Engineers.

Ludwig, D. F., Iannuzzi, T. J. & Esposito, A. N. (2003) *Phragmites* and environmental management: a question of values. *Estuaries* 26: 624–630.

Lugo, A. E. & Snedaker, S. C. (1974) The ecology of mangroves. Annual Reviews of Ecology and Systematics 5: 39–64.

Lugo, A. E., Brinson, M. & Brown, S., eds. (1990) *Ecosystems* of the World, vol. 15, Forested Wetlands. Amsterdam, the Netherlands: Elsevier.

Lundquist, J. (1998) Avert looming hydrocide. *Ambio* 27: 428–433.

Lutz, R. A., Shank, T. M., Fornari, D. J., et al. (1994) Rapid growth at deep-sea vents. Nature 371: 663–664.

Lynam, C., Gibbons, M., Axelsen, B., et al. (2006) Jellyfish overtake fish in a heavily fished ecosystem. *Current Biology* 16: R492–R493. MacIntyre, S., Flynn, K., Jellison, R. & Romero, J. (1999) Boundary mixing and nutrient fluxes in Mono Lake, California. *Limnology and Oceanography* **44**: 512–529.

Mack, R. N., Simberloff, D., Lonsdale, W. M., et al. (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689–710.

Mackay, A. W. & Tallis, J. T. (1996) Summit-type blanket mire erosion in the Forest of Bowland, Lancashire, UK: predisposing factors and implications for conservation. *Biological Conservation* 76: 31–44.

MacKenzie, S. (1993) Ecosystem management in the Great Lakes: some observations from three RAP sites. *Journal of Great Lakes Research* 19: 136–144.

Mackereth, F. J.H. (1965) Chemical investigations of lake sediments and their interpretation. *Proceedings of the Royal Society of London B* **161**: 293–375.

Madon, S. P., Williams, G. D., West, J. M. & Zedler, J. B. (2001) The importance of marsh access to growth of the California killifish, *Fundulus parvipinnis*, evaluated through bioenergetics modelling. *Ecological Modelling* 136: 149–165.

Madsen, T. V. & Sand-Jensen, K. (1991) Photosynthetic carbon assimilation in aquatic macrophytes. *Aquatic Botany* 41: 5–40.

Magnusson, J. J., Benson, B. J. & Kratz, T. K. (1990) Temporal coherence in the limnology of a suite of lakes in Wisconsin, USA. *Freshmater Biology* 23: 145–159.

Magnuson, J., Robertson, D., Benson, B., et al. (2000) Historical trends in lake and river ice cover in the Northern Hemisphere. Science 289: 1743–1746.

Magnuson, J. J., Webster, K. E., Assel, R. A., et al. (1997) Potential effects of climate changes on aquatic systems: Laurentian Great Lakes and Precambrian Shield Region. *Hydrological Processes* 11: 825–871.

Magoulick, D. D. (2000) Spatial and temporal variation in fish assemblages of drying stream pools: the role of abiotic and biotic factors. *Aquatic Ecology* **34**: 29–41.

Maguire, T. L., Saenger, P., Baverstock, P. & Henry, R. (2000) Microsatellite analysis of genetic structure in the mangrove species *Avicennia marina* (Forsk.) Vierh. (Avicenniaceae). *Molecular Ecology* 9: 1853–1862.

Mäkilä, M., Saarnisto, M. & Kankainen, T. (2001) Aapa mires as a carbon sink and source during the Holocene. *Journal of Ecology* 89: 589–599.

Malakoff, D. (2002) Miscue raises doubt about survey data. *Science* **298**: 515.

Malmer, N. (1962) Studies on mire vegetation in the archaean area of Southwestern Götaland (South Sweden). II. Distribution and seasonal variation in elementary constituents on some mire sites. *Opera Botanica* 7: 2–15. Malmer, N. (1988) Patterns in the growth and the accumulation of inorganic constituents in the *Sphagnum*-cover on ombrotrophic bogs in Scandinavia. *Oikos* 53: 105–120.

Malmer, N. & Wallén, B. (1993) Accumulation and release of organic matter in ombrotrophic bog hummocks: processes and regional variation. *Ecography* **16**: 193–211.

Malmer, N. & Wallén, B. (1996) Peat formation and mass balance in sub-arctic ombrotrophic peatlands around Abisko, northern Scandinavia. *Ecological Bulletins* 45: 79–92.

Malmer, N. & Wallén, B. (1999) The dynamics of peat accumulation on bogs: mass balance of hummocks and hollows and its variation throughout a millennium. *Ecography* 22: 736–750.

Malmer, N. & Wallén, B. (2004) Input rates, decay losses and accumulation rates of carbon in bogs during the last millennium: internal processes and environmental changes. *Holocene* 14: 111–117.

Malmer, N. & Wallén, B. (2005) Nitrogen and phosphorus in mire plants: variation during 50 years in relation to supply rate and vegetation type. *Oikos* 109: 539–554.

Malmer, N., Albinsson, C., Svensson, B. & Wallén, B. (2003) Interferences between *Sphagnum* and vascular plants: effects on plant community structure and peat formation. *Oikos* 100: 469–482.

Malmer, N., Johansson, T., Olsrud M. & Christensen, T. R. (2005) Vegetation, climatic changes and net carbon sequestration in a North-Scandinavian sub-arctic mire over 30 years. *Global Change Biology* 11: 1895–1909.

Malmer, N., Svensson, G. & Wallén, B. (1997) Mass balance and nitrogen accumulation in hummocks on a South Swedish bog during Late Holocene. *Ecography* 20: 535–549.

Malmqvist, B. (2002) Aquatic invertebrates in riverine landscapes. *Freshwater Biology* **47**: 679–694.

Malmqvist, B. & Rundle, S. R. (2002) Threats to the running water ecosystems of the world. *Environmental Conservation* 29: 134–153.

Malone, T. C., Màlej, A. & Smodlaka, N. (1996) Trends in land-use, water quality, and fisheries: a comparison of the northern Adriatic Sea and the Chesapeake Bay. *Periodicum Biologorum* 98: 137–148.

Maltby, E. (1986) Waterlogged Wealth: Why Waste the World's Wet Places? London, UK: Earthscan Publications.

Maltby, E., Hogan, D. V., Immirzi, C. P., Tellam, J. H. & van der Peijl, M. J. (1994) Building a new approach to the investigation and assessment of wetland ecosystem functioning. In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 637–658. Amsterdam, the Netherlands: Elsevier. Manabe, S., Stouffer, R. J., Spelman, M. J. & Bryan, K. (1991) Transient responses of a coupled ocean–atmosphere model to gradual changes in atmospheric CO2. I. Annual mean response. *Journal of Climate* 4: 785–818.

Mann, K. H. (1973) Seaweeds: their productivity and strategy for growth. *Science* 182: 975–981.

Mann, K. H. (1977) Destruction of kelp beds by sea urchins: a cyclical phenomenon or irreversible degradation? *Helgoländer Wissenschaftliche Meeresuntersuchungen* 30: 455–467.

Mann, K. H. (2000) Ecology of Coastal Waters, with Implications for Management, vol. 2. Oxford, UK: Blackwell Science.

Mann, M. E., Bradley, R. S. & Hughes, M. K. (1998) Global-scale temperature patterns and climate forcing over the past six centuries. *Nature* 392: 779–787.

Marbà, N. & Duarte, C. M. (1995) Coupling of seagrass (Cymodocea nodosa) patch dynamics to subaqueous dune migration. Journal of Ecology 83: 381–389.

Marbà, N. & Duarte, C. M. (1997) Interannual changes in seagrass (*Posidonia oceanica*) growth and environmental change in the Spanish Mediterranean littoral. *Limnology and Oceanography* 42: 800–810.

Marbà, N. & Duarte, C. M. (1998) Rhizome elongation and seagrass clonal growth. *Marine Ecology Progress Series* 174: 269–280.

Marbà, N., Duarte, C. M., Cebrián, J., *et al.* (1996) Growth and population dynamics of *Posidonia oceanica* on the Spanish Mediterranean coast: elucidating seagrass decline. *Marine Ecology Progress Series* 137: 203–213.

Marbà, N., Duarte, C. M., Holmer, M., et al. (2002) Assessing the effectiveness of protection on *Posidonia oceanica* populations in the Cabrera National Park (Spain). *Environmental Conservation* 29: 509–518.

Marchant, C. J. (1968) Evolution in *Spartina* (Gramineae). II. Chromosomes, basic relationships and the problem of the *S.* × *townsendii* agg. *Botanical Journal of the Linnean Society* **60**: 381–409.

Marsh, P., Lesack, L. F.W. & Roberts, A. (1999) Lake sedimentation in the Mackenzie Delta, NWT. *Hydrological Processes* 13: 2519–2536.

Martikainen, P. J., Nykanen, H., Crill, P. & Silvola, J. (1993) Effect of a lowered water table on nitrous oxide fluxes from northern peatlands. *Nature* 366: 51–53.

Martin, S., Munoz, E. & Drucker, R. (1997) Recent observations of a spring–summer surface warming over the Arctic Ocean. *Geophysical Research Letters* 24: 1259–1262.

Martinez Austria, P. & van Hofwegen, P., eds. (2006) Synthesis of the 4th World Water Forum 2006. Mexico DF, Mexico: Comisión National de Agua [www document]. URL http://www.worldwatercouncil.org.

Maser, C. & Sedell, J. R. (1994) From the Forest to the Sea: The Ecology of Wood in Streams, River, Estuaries, and Oceans. Delray Beach, FL, USA: St Lucie Press.

Masing, V., Svirezhev, Y. M., Loffler, H. & Patten, B. C. (1990) Wetlands in the biosphere. In: *Wetlands and Shallow Continental Water Bodies*, vol. 1, ed. B. C. Patten, pp. 313–344. The Hague, the Netherlands: SPB Academic Publishing.

Maslanik, J. A., Serreze, M. C. & Barry, R. G. (1996) Recent decreases in summer Arctic ice cover and linkages to atmospheric circulation anomalies. *Geophysical Research Letters* 23: 1677–80.

Mason, C. F., Underwood, G. J.C., Baker, N. R., et al. (2003) The role of herbicides in the erosion of salt marshes in eastern England. *Environmental Pollution* 122: 41–49.

Mason, R. P., Fitzgerald, W. F. & Morel, F. M.M. (1994) The biogeochemical cycling of elementary mercury: anthropogenic influences. *Geochima et Cosmochimica Acta* 58: 3191–3198.

Mason, R. P., Reinfelder, J. R. & Morel, F. M.M. (1995) Bioaccumulation of mercury and methylmercury. *Water*, *Air, and Soil Pollution* **80**: 915–921.

Mather, A. S. & Ritchie, W. (1977) *The Beaches of the Highlands and Islands of Scotland*. Perth, UK: Countryside Commission for Scotland.

Matsunaga, K., Kawaguchi, T., Suzuki Y. & Nigi, G. (1999) The role of terrestrial humic substances on the shift of kelp community to crustose coralline algae community of the southern Hokkaido Island in the Japan Sea. *Journal of Experimental Marine Biology and Ecology* 241: 193–205.

Matthiessen, P. & Gibbs, P. E. (1998) Critical appraisal of the evidence for tributyltin-mediated endocrine disruption. *Environmental Toxicology and Chemistry* **17**: 37–43.

Mattice, A. (2003) Eliminating fishing subsidies as a way to promote conservation. *Economic Perspectives: An Electronic Journal of the US Department of State* 8 [www document]. URL http://usinfo.state.gov/journals/ites/0103/ijee/ mattice.htm

Mauri, M., Polimeni, R., Modica, A. & Ferraro, M. (1998) Heavy metal bioaccumulation associated with drilling and production activities in middle Adriatic Sea. *Fresenius Environmental Bulletin* 7: 60–70.

Maykut, G. A. (1985) The ice environment. In: *Sea Ice Biota*, ed. R. A. Horner, pp. 21–82. Boca Raton, FL, USA: CRC Press.

Mazumder, A., Taylor, W. D., McQueen, D. J. & Lean, D. R.S. (1990) Effects of fish and plankton on lake temperature and mixing depth. *Science* 247: 312–315.

Mbewe, D. M.N. (1992) Agriculture as a component of wetlands conservation. In: *Managing the Wetlands of the Kafue Flats and Bangweulu Basin*, ed. R. C.V. Jeffery, H. N. Chabwela, G. Howard & P. J. Dugan, pp. 57–64. Gland, Switzerland: WWF and IUCN.

McAllister, D. E., Hamilton, A. L. & Harvey, B. (1997) Global freshwater biodiversity: striving for the integrity of freshwater ecosystems. *Sea Wind: Bulletin of Ocean Voice International* 11: 1–140.

McAllister, M., Craig, J. F., Davidson, N., Delany, S. & Seddon, M. (2000) Biodiversity impacts of large dams [www.document]. URL http://www.dams.org/kbase/ thematic/tr21.htm

McCabe, G. J. & Dettinger, M. D. (1999) Decadal variations in the strength of ENSO teleconnections with precipitation in the western United States. *International Journal of Climatology* **19**: 1399–1410.

McCartney, M. P., Masiyandima, M. & Houghton-Carr, H. A. (2005) Working Wetlands: Classifying Wetland Potential for Agriculture, Research Report No. 90. Colombo, Sri Lanka: International Water Management Institute.

McClanahan, T. R. (1988) Seasonality in East Africa's coastal waters. *Marine Ecology Progress Series* 44: 191–199.

McClanahan, T. R. (1992) Resource utilization, competition and predation: a model and example from coral reef grazers. *Ecological Modelling* **61**: 195–215.

McClanahan, T. R. (1995) A coral reef ecosystem–fisheries model: impacts of fishing intensity and catch selection on reef structure and processes. *Ecological Modelling* 80: 1–19.

McClanahan, T. R. (1999) Is there a future for coral reef parks in poor tropical countries? *Coral Reefs* 18: 321–325.

McClanahan, T. R. (2000a) Recovery of the coral reef keystone predator, *Balistapus undulatus*, in East African marine parks. *Biological Conservation* 94: 191–198.

McClanahan, T. R. (2000b) Bleaching damage and recovery potential of Maldivian coral reefs. *Marine Pollution Bulletin* 40: 587–597.

McClanahan, T. R. (2002a) The near future of coral reefs. Environmental Conservation 29: 460–483.

McClanahan, T. R. (2002b) The effects of time, habitat and fisheries management on Kenyan coral-reef associated gastropods. *Ecological Application* 12: 1484–1495. McClanahan, T. R. & Kurtis, J. D. (1991) Population regulation of the rock-boring sea urchin *Echinometra mathaei* (de Blainville). *Journal of Experimental Marine Biology and Ecology* 147: 121–146.

McClanahan, T. R. & Maina, J. (2003) Response of coral assemblages to the interaction between natural temperature variation and rare warm-water events. *Ecosystems* 6: 551–563.

McClanahan, T. R. & Mangi, S. (2000) Spillover of exploitable fishes from a marine park and its effect on the adjacent fishery. *Ecological Application* **10**: 1792–1805.

McClanahan, T. R. & Mangi, S. (2001) The effect of closed area and beach seine exclusion on coral reef fish catches. *Fisheries Management and Ecology* 8: 107–121.

McClanahan, T. R. & Obura, D. (1997) Sedimentation effects on shallow coral communities in Kenya. *Journal of Experimental Marine Biology and Ecology* 209: 103–122.

McClanahan, T. R., Cokos, B. A. & Sala, E. (2002*a*) Algal growth and species composition under experimental control of herbivory, phosphorus and coral abundance in Glovers Reef, Belize. *Marine Pollution Bulletin* **44**: 441–451.

McClanahan, T. R., Graham, N. A.J., Calnan, J. M. & MacNeil, M. A. (2007) Towards pristine biomass: reef fish recovery in coral reef marine protected areas in Kenya. *Ecological Applications* 17: 1055–1067.

McClanahan, T. R., Maina, J. & Pet-Soede, L. (2002*b*) Effects of the 1998 coral mortality event on Kenyan coral reefs and fisheries. *Ambio* 31: 543–550.

McClanahan, T. R., Maina, J., Moothien Pillay, R. & Baker, A. C. (2005) Effects of geography, taxa, water flow, and temperature variation on coral bleaching intensity in Mauritius. *Marine Ecology Progress Series* **298**: 131–142.

McClanahan, T. R., Muthiga, N. A., Kamukuru, A. T., Machano, H. & Kiambo, R. (1999) The effects of marine parks and fishing on the coral reefs of northern Tanzania. *Biological Conservation* 89: 161–182.

McClanahan, T. R., Muthiga, N. A. & Mangi, S. (2001) Coral and algal response to the 1998 coral bleaching and mortality: interaction with reef management and herbivores on Kenyan reefs. *Coral Reefs* **19**: 380–391.

McClanahan, T. R., Polunin, N. V.C. & Done, T. (2002c) Ecological states and the resilience of coral reefs. *Conservation Ecology* 6: 18.

McClanahan, T. R., Sheppard, C. R. & Obura, D. (2000) Coral Reefs of the Indian Ocean: Their Ecology and Conservation. New York, USA: Oxford University Press.

McClanahan, T. R., Uku, J. N. & Machano, H. (2002d) Effect of macroalgal reduction on coral-reef fish in the Watamu Marine National park, Kenya. *Marine and Freshwater Research* 53: 223–231.

McComb, A. J. & Lukatelich, R. J. (1995) The Peel-Harvey estuarine system, Western Australia. In: *Eutrophic Shallow Estuaries and Lagoons*, ed. A. J. McComb, pp. 5–17. Boca Raton, FL, USA: CRC Press.

McComb, A. J., ed. (1995) *Eutrophic Shallow Estuaries and Lagoons*. Boca Raton, FL, USA: CRC Press.

McConnaughey, R. A., Mier, K. L. & Dew, C. B. (2000) An examination of chronic trawling effects on soft-bottom benthos of the eastern Bering Sea. *ICES Journal of Marine Science* 57: 1377–1388.

McCoy, M. B. & Rodriguez, J. M. (1994) Cattail (*Typha domingensis*) eradication methods in the restoration of a tropical, seasonal, freshwater marsh. In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 469–483. Amsterdam, the Netherlands: Elsevier.

McCulloch, M., Fallon, S., Wyndham, T., et al. (2003) Coral record of increased sediment flux to the inner Great Barrier Reef since European settlement. *Nature* 421: 727–730.

McDowell, J. E. (1993) How marine animals respond to toxic chemicals in coastal ecosystems. *Oceanus* 36: 56–63.

McGowan, J. A., Cayan, D. R. & Dorman, L. M. (1998) Climate–ocean variability and ecosystem response in the Northeast Pacific. *Science* 281: 210–217.

McGrady-Steed, J., Harris, P. M. & Morin, P. J. (1997) Biodiversity regulates ecosystem predictability. *Nature* **390**: 162–165.

McGrath, D., Castro, F. de, Câmara, E. & Futemma, C. (1999) Community management of floodplain lakes and the sustainable development of Amazonian fisheries. In: Várzea: Diversity, Development, and Conservation of Amazonia's Whitewater Floodplains, Advances in Economic Botany, vol. 13, ed. C. Padoch, J. M. Ayres, M. Pinedo-Vasquez & A. Henderson, pp. 59–82. New York, USA: New York Botanical Garden Press.

McGuire, D., Melillo, J. M. & Joyce, L. A. (1995) The role of nitrogen in the response of forest net primary production to elevated atmospheric carbon dioxide. *Annual Review of Ecology and Systematics* **26**: 473–503.

McIntyre, A. D. (1992) The current state of the oceans. *Marine Pollution Bulletin* 25: 1–4.

McIntyre, A. D. (1995) Human impact on the oceans: the 1990s and beyond. *Marine Pollution Bulletin* **31**: 147–151.

McKay, W. A., Johnson, C. E. & Branson, J. R. (1986)
Environmental radioactivity in Caithness and Sutherland.
III. Initial measurements and modeling of inshore waters. *Nuclear Energy* 27: 321–335. McKee, D., Atkinson, D., Collings, S. E., et al. (2003) Response of freshwater microcosm communities to nutrients, fish and elevated temperature during winter and summer. *Limnology and Oceanography* 48: 707–722.

McKenzie, L. J., Lee Long, W. J., Coles, R. G. & Roder, C. A. (2000) Seagrass-Watch: community based monitoring of seagrass resources. *Biologia Marina Mediterranea* 7: 393–396.

McLachlan, A. (1980) Exposed sandy beaches as semi-closed ecosystems. *Marine Environmental Research* **4**: 59–63.

McLachlan, A. (1996) Physical factors in benthic ecology: effects of changing sand particle size on beach fauna. *Marine Ecology Progress Series* 131: 205–217.

McLachlan, A. & Harty, B. (1981) Effects of crude oil pollution on the supralittoral meiofauna of a sandy beach. *Marine Environmental Research* 7: 71–80.

McLachlan, A. & Young, N. (1982) Effects of low temperatures on the burrowing rates of four sandy beach mollusks. *Journal of Experimental Marine Biology and Ecology* 65: 275–284.

McLaren, A. S., Bourke, R. H., Walsh, J. E. & Weaver, R. L. (1994). Variability in sea-ice thickness over the North Pole from 1958 to 1992. In: *The Polar Oceans and Their Role in Shaping the Global Environment: The Nansen Centennial Volume*, ed. O. M. Johannessen, R. D. Muench & J. E. Overland, Geophysical Monograph No. 85, pp. 363–371. Washington, DC, USA: American Geophysical Union.

McMahon, K. & Walker, D. I. (1998) Fate of seasonal, terrestrial nutrient inputs to a shallow seagrass dominated embayment. *Estuarine Coastal and Shelf Science* 46: 15–25.

McMahon, T. A. & Finlayson, B. L. (2003) Droughts and anti-droughts: the low flow hydrology of Australian rivers. *Freshwater Biology* **48**: 1147–1160.

McManus, J. W., Menez, L. A.B., Kesner-Reyes, K. N., Vergara, S. G. & Ablan, M. C. (2000) Coral reef fishing and coral-algal phase shifts: implications for global reef status. *Journal of Marine Science* 57: 572–578.

McMillan, C. & Moseley, F. N. (1967) Salinity tolerances of five marine spermatophytes of Redfish Bay, Texas. *Ecology* 46: 503–506.

McMurtry, G. (2001) Authigenic deposits. In: *Encyclopedia of Ocean Sciences*, ed. S. A. Thorpe & K. K. Turekian, pp. 201–220. London, UK: Academic Press.

McNeil, P. & Waddington, J. M. (2003) Moisture controls on Sphagnum growth and CO₂ exchange on a cutover bog. Journal of Applied Ecology **40**: 354–367.

McNeill, W. H. (1976) *Plagues and Peoples*. New York, USA: Anchor Books.

McQuaid, C. D. & Dower, K. M. (1990) Enhancement of habitat heterogeneity and species richness on rocky shores inundated by sand. *Oecologia* 84: 142–143.

McRoy, C. P. & Helfferich, C. (1977) Seagrass Ecosystems: A Scientific Perspective. New York, USA: Marcel Dekker.

MDBMC (1995) Floodplain Wetlands Management Strategy for the Murray–Darling Basin. Canberra, ACT, Australia: Murray–Darling Basin Ministerial Council.

MDBMC (1999) The Salinity Audit of the Murray–Darling Basin: A 100-Year Perspective. Canberra, ACT, Australia: Murray–Darling Basin Commission.

Meade, R. H. (1969) Errors in using modern stream-loading data to estimate natural rates of denudation. *Bulletin of the Geological Society of America* 80: 1265–1274.

Meakins, N. C., Bubb, J. C. & Lester, J. N. (1985) The mobility, partitioning and degradation of atrazine and simazine in the salt marsh environment. *Marine Pollution Bulletin* 30: 812–819.

Medio, D., Sheppard, C. R.C. & Gasgoine, J. (2000) The Red Sea. In: Coral Reefs of the Indian Ocean: Their Ecology and Conservation, ed. T. R. McClanahan, C. R.C. Sheppard & D. O. Obura, pp. 231–255. New York, USA: Oxford University Press.

Mee, L. D. (2001) Eutrophication in the Black Sea and a basin-wide approach to its control. In: *Science and Integrated Coastal Management*, ed. B. von Bodungen & R. K. Turner, pp. 71–91. Berlin, Germany: Dahlem University Press.

Meehl, G. A. (1992) Effect of tropical topography on global climate. *Annual Review of Earth and Planet Planetary Science* **20**: 85–112.

Mees, C. C., Pilling, G. M. & Barry, C. J. (1999) Commercial inshore fishing activity in the British Indian Ocean Territory. In: *Ecology of the Chagos Archipelago*, ed. C. R.C. Sheppard & M. R.D. Seaward, pp. 327–345. London, UK: Linnean Society.

Meeuwig, J. J., Rasmussen, J. B. & Peters, R. H. (1998) Turbid waters and clarifying mussels: their moderation of empirical chl:nutrient relations in estuaries in Prince Edward Island, Canada. *Marine Ecology Progress Series* 171: 139–150.

Meggers, B. J. (1987) The early history of man in Amazonia. In: *Biogeography and Quaternary History in Tropical America*, ed. T. C. Whitmore & G. T. Prance, pp. 151–174. Oxford, UK: Clarendon Press.

Mehner, T., Benndorf, J., Kasprzak, P. & Koschel, R. (2002) Biomanipulation of lake ecosystems: successful applications and expanding complexity in the underlying science. *Freshmater Biology* 47: 2453–2465. Meijer, M.-L., de Boois, I., Scheffer, M., Portielje, R. & Hosper, H. (1999) Biomanipulation in shallow lakes in the Netherlands: an evaluation of 18 case studies. *Hydrobiologia* 408/409: 13–30.

Meinesz, A., de Vaugelas, J., Hesse, B. & Mari, X. (1993) Spread of the introduced tropical green alga *Caulerpa taxifolia* in northern Mediterranean waters. *Journal of Applied Phycology* 5: 141–147.

Mekong River Commission (2002) International River Prize [www document]. URL http://www.mrcmekong.org/ new_events/press_release/2002/

Melack, J. M. & Kilham, P. (1974) Photosynthetic rates of phytoplankton in East-African alkaline, saline lakes. *Limnology and Oceanography* 19: 743–755.

Melillo, J. M., McGuire, A. D., Kicklighter, D. W., et al. (1993) Global climate change and terrestrial net primary production. *Nature* 363: 234–240.

Mendelssohn, I. A. & Morris, J. T. (2000) Ecophysiological controls on the growth of *Spartina alterniflora*. In: *Concepts and Controversies in Tidal Marsh Ecology*, ed. M. P. Weinstein & D. P. Kreeger, pp. 59–80. Dordrecht, the Netherlands: Kluwer.

Mendelssohn, I. A., Hester, M. W., Monteferrante, F. J. & Talbot, F. (1991) Experimental dune building and vegetative stabilization in a sand-deficient barrier island setting on the Louisiana coast, USA. *Journal of Coastal Research* 7: 137–149.

Menge, B. A. (1991) Relative importance of recruitment and other causes of variation in rocky intertidal community structure. *Journal of Experimental Marine Biology and Ecology* 145: 69–100.

Menge, B. A. (1995) Indirect effects in marine rocky intertidal interaction webs: patterns and importance. *Ecological Monographs* 65: 21–74.

Menge, B. A. (2000) Top-down and bottom-up community regulation in marine rocky intertidal habitats. *Journal of Experimental Marine Biology and Ecology* **250**: 257–289.

Menge, B. A., Daley, B. A., Wheeler, P. A., et al. (1997) Benthic-pelagic links and rocky intertidal communities: bottom-up effects on top-down control?. Proceedings of the National Academy of Sciences of the USA 94: 14 530–14 535.

Menge, B. A., Lubchenco, J., Bracken, M. E.S., et al. (2003) Coastal oceanography sets the pace of rocky intertidal community dynamics. *Proceedings of the National Academy* of Sciences of the USA 100: 12 229–12 234.

Merrett, N. R. & Haedrich, R. L. (1997) *Deep-Sea Demersal Fish and Fisheries.* London, UK: Chapman & Hall. Merron, G. S., Bruton, M. N. & de Lalouviere, P. la H. (1993) Implications of water release from the Pongolapoort Dam for the fish and fishery of the Pongolo floodplain, Zululand. *South African Journal of Aquatic Sciences* 19: 34–49.

Mertes, L. A.K. (2000) Inundation hydrology. In: *Inland Flood Hazards*, ed. E. E. Wohl, pp. 145–166. Cambridge, UK: Cambridge University Press.

Meyer, J. L., Sale, M. J., Mulholland, P. J. & Poff, N. L. (1999) Impacts of climate change on aquatic ecosystem functioning and health. *Journal of the American Water Resources Association* 35: 1373–1386.

Meyerhoff, J. & Dehnhardt, A. (2002) Nachhaltige Entwicklung der Elbe. *Ökologisches Wirtschaften* 5: 27–28.

Meyer-Reil, L. A. & Koster, M. (2000) Eutrophication of marine waters: effects on benthic microbial communities. *Marine Pollution Bulletin* **41**: 255–263.

Meyers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A.B. & Kent, J. (2000) Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858.

Meyerson, L. A., Saltonstall, K., Windham, L., Kiviat, E. & Finlay, S. (2000) A comparison of *Phragmites australis* in freshwater and brackish marsh environments in North Australia. *Wetlands Ecology and Management* 8: 89–103.

Michigan Department of Environmental Quality (2000) Great Lakes trends: into the New Millennium. Unpublished report. Lansing, MI, USA: Michigan Department of Environmental Quality.

Micklin, P. P. (1988) Desiccation of the Aral Sea: a water management disaster in the Soviet Union. *Science* 241: 1170–1175.

Micklin, P. P. (1998) International and regional responses to the Aral crisis: an overview of efforts and accomplishments. *Post-Soviet Geography and Economics* **39**: 399–416.

Middleton, B. A. (1998) The water buffalo controversy in Keoladeo National Park, India. *Ecological Modelling* 106: 93–98.

Middleton, B. A. (2002a) Winter burning and the reduction of Cornus sericea in sedge meadows in southern Wisconsin. Restoration Ecology 10: 1–8.

Middleton, B. A. (2002b) Nonequilibrium dynamics of sedge meadows grazed by cattle in southern Wisconsin. *Plant Ecology* 161: 89–110.

Middleton, B. A. (2003) Ecology and objective based management: case study of the Keoladeo National Park, Bharatpur, Rajasthan. In: *Battles over Nature: Science and the Politics of Conservation*, ed. V. Saberwal & M. Rangarajan, pp. 86–116. New Delhi, India: Permanent Black Publishers.

Middleton, B. A. (1999) Wetland Restoration, Flood Pulsing and Disturbance Dynamics. New York, USA: John Wiley.

- Midun, Z. & Lee, S. -C. (1995) Implications of a greenhouseinduced sea-level rise: a national assessment for Malaysia. *Journal of Coastal Research* Special Issue 14: 96–115.
- Mieszkowska, N., Leaper, R., Moore, P., et al. (2005) Marine biodiversity and climate change: assessing and predicting the influence of climatic change using intertidal rocky shore biota. Occasional Publications, Marine Biological Association of the United Kingdom 20: 1–53.
- Miliman, J. D. & Meade, R. H. (1993) World-wide delivery of river sediment to the oceans. *Journal of Geology* 91: 1–21.
- Millard, A. V. & Evans, P. R. (1984) Colonization of mudflats by *Spartina anglica*: some effects on invertebrate and shore bird populations at Lindisfarne. In: Spartina anglica *in Great Britain*, ed. P. Doody, pp. 41–48. Attingham Park, UK: Nature Conservancy Council.
- Miller, M. W., Hay, M. E., Miller, S. L., et al. (1999) Effects of nutrients versus herbivores on reef algae: a new method for manipulating nutrients on coral reefs. *Limnology and Oceanography* 44: 1847–1861.
- Miller, R. J., Smith, C. R., DeMaster, D. J. & Fornes, W. L. (2000) Feeding selectivity and rapid particle processing by deep-sea megafaunal deposit feeders: a Th-234 tracer approach. *Journal of Marine Research* 58: 653–673.
- Miller, W. R. & Egler, F. E. (1950) Vegetation of the Wequetequock–Pawcatuck tidal marshes, Connecticut. *Ecological Monographs* **20**: 143–172.
- Milliman, J. D. & Meade, R. H. (1983) World-wide delivery of river sediment to the oceans. *Journal of Geology* 91: 1–21.
- Mills, C. E. (2001) Jellyfish blooms: are populations increasing globally in response to changing ocean conditions? *Hydrobiologia* 451: 55–68.
- Milner, A. M. (1994) System recovery. In: *The Rivers Handbook*, vol. 2, ed. P. Calow & G. E. Petts, pp. 76–97. Oxford, UK: Blackwell Scientific Publications.
- Milner, J. (1874) Report on the fisheries of the Great Lakes: the results of inquiries prosecuted in 1871 and 1872. In: *Report of the Commissioner for 1872 and 1873*. Washington, DC, USA: US Commission of Fish and Fisheries.
- Milton, K. (1998) Alternative futures to environmental disaster. Global Ecology and Biogeography Letters 7: 150–152.
- Minchinton, T. & Bertness, M. D. (2003) Soil salinity, nitrogen availability and the expansion of *Phragmites australis* in New England salt marshes. *Ecological Applications* 13: 1400–1416.
- Ministerio del Medio Ambiente (2001) *Politica nacional para* humedales interiores de Colombia: estrategias para su conservación y uso racional. Bogotá, Colombia: Ministerio del Medio Ambiente, Consejo Nacional Ambiental.

- Minshall, G. W. (1992) Troubled waters of greenhouse earth: summary and synthesis. In: *Global Climate Change and Freshwater Ecosystems*, ed. P. Firth & S. G. Fisher, pp. 308–318. Berlin, Germany: Springer-Verlag.
- Mirza, A. M. Q. (2002) Global warming and changes in the probability of occurrence of floods in Bangladesh and implications. *Global Environmental Change* 12: 127–138.
- Mirza, M. M. (2003) Three recent extreme floods in Bangladesh: a hydro-meteorologic analysis. *Natural Hazards* 28: 35–64.
- MIT (2003) Joint Program on the Science and Policy of Global Change, Massachusetts Institute of Technology [www document]. URL http://web.mit.edu/globalchange/www/
- Mitchell, B. D. & Geddes, M. C. (1977) Distribution of the brine shrimps *Parartemia zietziana* (Sayce) and *Artemia* salina (L.) along a salinity and oxygen gradient in a South Australian salt-field. *Freshwater Biology* 7: 461–467.
- Mitchell, J. K. (2003) European river floods in a changing world. *Risk Analysis* 23: 567–574.
- Mitchell, R. B., Clark, W. C. & Cash, D. W. (2006)
 Information and influence. In: *Global Environmental* Assessments: Information and Influence, ed. R. B. Mitchell, W. C. Clark, D. W. Cash & N. M. Dickson, pp. 307–338.
 Cambridge, MA, USA: MIT Press.
- Mitsch, W. J. & Gosselink, J. G. (2000) *Wetlands*, 3rd edn. New York, USA: John Wiley.
- Mitsch, W. J. (1993) Ecological engineering: a cooperative role with the planetary life-support system. *Environmental Science and Technology* 27: 438–445.
- Mitsch, W. J., Mitsch, R. H. & Turner, R. E. (1994) Wetlands of the Old and New Worlds: ecology and management. In: *Global Wetlands: Old World and New*, ed. W. J. Mitsch, pp. 3–56. Amsterdam, the Netherlands: Elsevier.
- Moberg, F., Nystrom, M., Kautsky, N., Tedengren, M. & Jarayabhand, P. (1997) Effects of reduced salinity on the rates of photosynthesis and respiration in the hermatypic corals *Porites lutea* and *Pocillopora damicornis*. *Marine Ecology Progress Series* 157: 53–59.
- Mono Lake Committee (2006) Home page [www document]. URL http://www.monolake.org
- Monroe, M. W. & Kelly, J. (1992) *State of the Estuary*, Technical Report. Oakland, CA, USA: San Francisco Estuary Project.
- Monson, D. H., Doak, D. F., Ballachey, B. E., Johnson, A. M. & Bodkin, J. L. (2000) Long-term impacts of the Exxon Valdez oil spill on sea otters, assessing through age-dependent mortality patterns. *Proceedings of the National Academy of Sciences of the USA* 97: 6562–6567.

Monteiro, L. R., Costa, V., Furness, R. W. & Santos, R. S. (1996) Mercury concentrations in prey fish indicate enhanced bioaccumulation in mesopelagic environments. *Marine Ecology Progress Series* 141: 21–25.

Monteith, D. T., Hildrew, A. G., Flower, R. J., et al. (2005) Biological responses to the chemical recovery of acidified fresh waters in the UK. *Environmental Pollution* 137: 83–101.

Mooney, H. A. & Hobbs, R. J., eds. (2000) *Invasive Species in a Changing World*. Washington, DC, USA: Island Press.

Moore, P. D. (1982) How to reproduce in bogs and fens. *New Scientist* 5 August: 369–371.

Moore, P. D. (1990) Soils and ecology: temperate wetlands. In: *Wetlands: A Threatened Landscape*, ed.
M. Williams, pp. 95–114. Oxford, UK: Blackwell Scientific Publications.

Moore, P. D. (1993) The origin of blanket mire, revisited. In: *Climate Change and Human Impact on the Landscape*, ed. F. M. Chambers, pp. 217–224. London, UK: Chapman & Hall.

Moore, P. D. (2002) The future of cool temperate bogs. *Environmental Conservation* **29**: 3–20.

Moore, P. D. & Bellamy, D. J. (1973) *Peatlands*. London, UK: Paul Elek.

Moore, P. D., Chaloner, W. & Stott, P. (1996) *Global Environmental Change*. Oxford, UK: Blackwell Scientific Publications.

Moore, P. D., Merryfield, D. L. & Price, M. D.R. (1984) The vegetation and development of blanket mires. In: *European Mires*, ed. P. D. Moore, pp. 203–235. London, UK: Academic Press.

Moran, P. J. (1986) The *Acanthaster* phenomenon. *Oceanography and Marine Biology Annual Review* 24: 379–480.

Moreno, C. A., Sutherland, J. P. & Jara, H. F. (1984) Man as a predator in the intertidal zone of southern Chile. *Oikos* 42: 155–160.

Moreno-Casasola, P. (1986) Sand movement as a factor in the distribution of plant communities in a coastal dune system. *Vegetatio* **65**: 67–76.

Morgan, C. L. (2000) Resource estimates of the Clarion-Clipperton manganese nodule deposits. In: *Handbook of Marine Mineral Deposits*, ed. D. S. Cronan, pp. 145–170. Boca Raton, FL, USA: CRC Press.

Morgan, P. A. & Short, F. T. (2002) Using functional trajectories to track constructed salt marsh development in the Great Bay Estuary, Maine/New Hampshire, USA. *Restoration Ecology* 10: 461–473. Morita, K. & Yamamoto, S. (2002) Effects of habitat fragmentation by damming on the persistence of stream-dwelling charr populations. *Conservation Biology* 16: 1318–1323.

Moritz, R. E., Bitz, C. M. & Steig, E. J. (2002) Dynamics of recent climate change in the Arctic. *Science* 297: 1497–1502.

Mork, M. (1996) The effects of kelps on wave dumping. *Sarsia* 80: 323–327.

Mormede, S. & Davies, I. M. (2001) Heavy metal concentrations in commercial deep-sea fish from the Rockall Trough. *Continental Shelf Research* 21: 899–916.

Morris, B. L., Lawrence, A. R.L., Chilton, P. J.C., et al. (2003) Groundwater and its Susceptibility to Degradation: A Global Assessment of the Problem and Options for Management, Early Warning and Assessment Report Series Rs 03-3. Nairobi, Kenya: United Nations Environment Programme.

Morris, J. T. (1995) The salt and water balance of intertidal sediments: results from North Inlet, South Carolina. *Estuaries* 18: 556–567.

Morris, J. T., Sundareshwar, P. V., Nietch, C. T., Kjerfve, B. & Cahoon, D. R. (2002) Responses of coastal wetlands to rising sea level. *Ecology* 83: 2869–2877.

Morrison, J., Quick, M. C. & Foreman, M. G.G. (2002) Climate change in the Fraser River watershed: flow and temperature projections. *Journal of Hydrology* 263: 230–244.

Mortensen, P. B., Buh-Mortensen, L., Gordon Jr, D. C., et al. (2005) Effects of fisheries on deep-water gorgonian corals in the Northeast Channel, Nova Scotia (Canada). In: Benthic Habitats and the Effects of Fishing, ed. P. Barnes & J. Thomas, pp. 369–382. Bethesda, MD, USA: American Fisheries Society.

Mortimer, C. (2004) Lake Michigan in Motion: Responses Of An Inland Sea To Weather, Earth Spin, and Human Activities. Madison, WI, USA: University of Wisconsin Press.

Morton, B. (1996) Protecting Hong Kong's marine biodiversity: present proposals, future challenges. *Environmental Conservation* 23: 55–65.

Morton, S. R. & Brennan, K. G. (1991) Birds. In: Monsoonal Australia: Landscape, Ecology and Man in Northern Australia, ed. C. D. Haynes, M. G. Ridpath & M. A.J. Williams, pp. 133–149. Rotterdam, the Netherlands: A. A. Balkema.

Moser, F. C. & Bopp, R. F. (2001) Particle-associated contaminants in the Barnegat Bay–Little Egg Harbor Estuary. *Journal of Coastal Research* Special Issue 32: 229–242.

Moser, M., Prentice, C. & Frazier, S. (1996) A global overview of wetland loss and degradation. In: *Proceedings* of the 6th Meeting of the Conference of the Contracting Parties, Brisbane, Australia, 19–27 March 1996, vol. 10B/12: Technical Sessions B and D. Gland, Switzerland: Ramsar Convention Bureau. Available online of www.ramsar.org/about/about_wetland_loss.htm

Moser, M., Prentice, C. & Frazier, S. (2003) A global overview of wetland loss and degradation. Ramsar, Gland, Switzerland [www document]. URL http://www.ramsar. org/about/about_wetland_loss.htm

Moss, B. (1995) The emperor's clothes of knowledge and the seamless cloth of wisdom. In: *Science for the Earth*, 9th edn, ed. T. Wakeford & M. Walters, pp. 321–348. Chichester, UK: John Wiley.

Moss, B. (1998) Ecology of Fresh Waters: Man and Medium, Past to Future, 3rd edn. Oxford, UK: Blackwell Science.

Moss, B., McKee, D., Atkinson, D., *et al.* (2003) How important is climate? Effects of warming, nutrient addition and fish on phytoplankton in shallow lake microcosms. *Journal of Applied Ecology* **40**: 782–792.

Moss, R. H. & Schneider, S. H. (1997) Characterizing and communicating scientific uncertainty: building on the IPCC 2nd Assessment. In: *Elements of Change*, ed. S. J. Hassol & J. Katzenberger, pp. 90–135. Aspen, CO, USA: Aspen Global Change Institute.

Moy, L. D. & Levin, L. A. (1991) Are Spartina marshes a replaceable resource? A functional approach to evaluation of marsh creation efforts. *Estuaries* 14: 1–16.

Moyle, P. B. (1999) Effects of invading species on freshwater and estuarine ecosystems. In: *Invasive Species and Biodiversity Management*, ed. O. T. Sandlund, P. J. Schei & Å. Viken, pp. 177–191. Dordrecht, the Netherlands: Kluwer.

Moyle, P. B. & Leidy, R. A. (1992) Loss of biodiversity in aquatic ecosystems: evidence from fish faunas. In: Conservation Biology: The Theory and Practice of Nature Conservation, Preservation and Management, ed.
P. L. Fiedler & S. K. Jain, pp. 127–169. New York, USA:

Chapman & Hall.

MSC (2004) Marine Stewardship Council: homepage [www document]. URL http://www.msc.org

Muchlstein, L. K., Porter, D. & Short, F. T. (1988) Labyrinthula sp., a marine slime mold producing the symptoms of wasting disease in eelgrass, Zostera marina. Marine Biology 99: 465–472.

Muehlstein, L. K., Porter, D. & Short F.T, (1991) Labyrinthula zosterae sp. nov., the causative agent of wasting disease of eelgrass, Zostera marina. Mycologia 83: 180–191.

Mugidde, R. (1993) The increase in phytoplankton primary productivity and biomass in Lake Victoria (Uganda). *Verhandlungen Internationale Vereinigung für Limnologie* 25: 846–849. Mulder, T. & Syvitski, J. P.M. (1995) Turbidity currents generated at river mouths during exceptional discharges to the world's oceans. *Journal of Geology* 103: 285–299.

Muller, R. A. & MacDonald, G. J. (1997) Glacial cycles and astronomical forcing. *Science* 277: 215–218.

Mullineaux, L. S. (1987) Organisms living on manganese nodules and crusts: distribution and abundance at three North Pacific sites. *Deep-Sea Research* 34: 165–184.

Mumby, P. J., Chisholm, J. R. M., Edwards, A. J., et al. (2001) Unprecedented bleaching-induced mortality in *Porites* spp. at Rangiroa Atoll, French Polynesia. *Marine Biology* 139: 183–189.

Munro, D. C. & Touron, H. (1997) The estimation of marshland degradation in southern Iraq using multitemporal Landsat TM images. *International Journal of Remote Sensing* 18: 1597–1606.

Munro, J. L. (1989) Fisheries for giant clams (Tridacnidae: Bivalvia) and prospects for stock enhancement. In: *Marine Invertebrate Fisheries: Their Assessment and Management*, ed. J. F. Caddy, pp. 541–558. New York, USA: John Wiley.

Munro, J. L. (1996) The scope of tropical reef fisheries and their management. In: *Reef Fisheries*, vol. 20, ed. N. V. C. Polunin & C. M. Roberts, pp. 1–14. London, UK: Chapman & Hall.

Murawski, S. (2000) Definitions of overfishing from an ecosystem perspective. *ICES Journal of Marine Science* 57: 649–658.

Murdoch, P. S., Baron, J. S. & Miller, T. L. (2000) Potential effects of climate change on surface-water quality in North America. *Journal of the American Water Resources Association* **36**: 347–366.

Murphy, E. J., Clarke, A., Symon, C. & Priddle, J. (1995) Temporal variation in Antarctic sea-ice: analysis of a long-term fast-ice record from the South Orkney Islands. *Deep-Sea Research I* 42: 1045–1062.

Murphy, E. J., Watkins, J. L., Reid, K., et al. (1998) Interannual variability of the South Georgia marine ecosystem: biological and physical sources of variation in the abundance of krill. *Fisheries Oceanography* 7: 381–390.

Murphy, G. I. (1968) Patterns in life history and the environment. *American Naturalist* 102: 391–403.

Muthiga, N. A., Riedmiller, S., Carter, E., et al. (2000)
Management status and case studies. In: Coral Reefs of the Indian Ocean: Their Ecology and Conservation, ed.
T. R. McClanahan, C. S. Sheppard & D. Obura,

pp. 473-505. New York, USA: Oxford University Press.

Muthuri, F. M., Jones, M. B. & Imbamba, S. K. (1989) Primary productivity of papyrus (*Cyperus papyrus*) in a tropical swamp: Lake Naivasha, Kenya. *Biomass* 18: 1–14.

Myers, R. A. & Worm, B. (2003) Rapid worldwide depletion of predatory fish communities. *Nature* **423**: 280–283.

Myers, R. A., Hutchings, J. A. & Barrowman, N. J. (1997) Why do fish stocks collapse? The example of cod in Atlantic Canada. *Ecological Applications* 7: 91–106.

Nachtnebel, H.-P. (2000) The Danube river basin environmental programme: plans and actions for a basin wide approach. *Water Policy* 2: 113–129.

Naeem, S. (1998) Species redundancy in ecosystem reliability. Conservation Biology 12: 39–45.

Naeem, S., Knops, J. M. H., Tilman, D., et al. (2000) Plant diversity increases resistance to invasion in experimental grassland plots. Oikos 91: 97–108.

Naganobu, M., Kutsuwada, K., Sasai, Y., Taguchi, S. & Siegel, V. (1999) Relationships between Antarctic krill (*Euphausia superba*) variability and westerly fluctuations and ozone depletion in the Antarctic Peninsula area. *Journal of Geophysical Research: Oceans* 104: 20651–20665.

Naiman, R. J., ed. (1992) Watershed Management: Balancing Sustainability and Environmental Change. New York, USA: Springer-Verlag.

Naiman, R. J. & Décamps, H. (1997) The ecology of interfaces: riparian zones. *Annual Review of Ecology and Systematics* 28: 621–658.

Naiman, R. J., Bunn, S. E., Nilsson, C., et al. (2002) Legitimizing fluvial ecosystems as users of water: an overview. *Environmental Management* 30: 455–467.

Naiman, R. J., Décamps, H. & McClain, M. E. (2005) *Riparia*. San Diego, CA, USA: Academic Press.

Naiman, R. J., Décamps, H. & Pollock, M. (1993) The role of riparian corridors in maintaining regional biodiversity. *Ecological Applications* 3: 209–212.

Nakamura, T. & van Woesik, R. (2001) Differential survival of corals during the 1998 bleaching event is partially explained by water-flow rates and passive diffusion. *Marine Ecology Progress Series* 212: 301–304.

Nalepa, T., Fahnenstiel, G. & Johengen, T. (1999) Impact of the zebra mussel (*Dreissena polymorpha*) on water quality: a case study in Saginaw Bay, Lake Huron. In: *Nonindigenous Freshwater Organisms, Vectors, Biology, and Impacts*, ed.
R. Claudi & J. H. Leach, pp. 255–271. Boca Raton, FL, USA: Lewis Publishers.

Nalepa, T., Hartson, D., Buchanan, J., et al. (2000a) Spatial variation in density, mean size and physiological condition

of the holarctic amphipod *Diporeia spp* in Lake Michigan. *Freshwater Biology* **43**: 107–119.

Nalepa, T., Lang, G. & Fanslow, D. (2000b) Trends in the macroinvertebrate populations in southern Lake Michigan. *Verhandlungen Internationale Vereinigung für Limnologie* 27: 2540–2545.

Nasholm, T., Ekblad, A., Nordin, A., et al. (1998) Boreal forest plants take up organic nitrogen. Nature 392: 914–916.

National Academy Press (1993) *Managing Wastewater in Coastal Urban Areas*. Washington, DC, USA: National Academy Press.

National Estuarine Research Reserve System (2001) The National Estuarine Research Reserve's System-Wide Monitoring Program (SWMP): A Scientific Framework and Plan for Detection of Short-Term Variability and Long-Term Change in Estuaries and Coastal Habitats of the United States, Technical Report, National Estuarine Research Reserve System. Silver Spring, MD, USA: National Oceanic and Atmospheric Administration.

National Estuary Program (1997*a*) Nutrient Overloading, Technical Report. Washington, DC, USA: US Environmental Protection Agency.

National Estuary Program (1997b) Management Approaches Being Used to Address Critical Issues, Technical Report. Washington, DC, USA: US Environmental Protection Agency.

National Estuary Program (1997c) Declines in Fish and Wildlife Populations, Technical Report. Washington, DC, USA: US Environmental Protection Agency.

Nauman, J. & Cory, R. L. (1969) Thermal additions and epifaunal organisms at Chalk Point, Maryland. *Chesapeake Science* 18: 218–226.

Navodaru, I., Staras, M. & Cerniseucu, I. (2001) The challenge of sustainable use of the Danube Delta Fisheries, Romania. *Fisheries Management and Ecology* 8: 323–332.

Naylor, E. (1965) Effects of heated effluents upon marine and estuarine organisms. Advances in Marine Biology 3: 63–103.

Naylor, R. L., Goldberg, J., Primavera, J. H., *et al.* (2000) Effect of aquaculture on world fish supplies. *Nature* **405**: 1017–1024.

Naylor, R., Hindar, K., Fleming, I. A., et al. (2005) Fugitive salmon: assessing the risks of escaped fish from net-pen aquaculture. *BioScience* 55: 427–437.

Neckles, H. A. & Neill, C. (1994) Hydrologic control of litter decomposition in seasonally flooded prairie marshes. *Hydrobiologia* 286: 155–165.

Nedwell, D. B. (1975) Inorganic nitrogen metabolism in an eutrophicated tropical estuary. *Water Research* 9: 221–231. Neiff, J. J. (2001) Diversity in some tropical wetland systems of South America. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, vol. 2, ed. B. Gopal, W. J. Junk & J. A. Davis, pp. 157–186. Leiden, the Netherlands: Backhuys.

Nekola, J. C. (1994) The environment and vascular flora of northeastern Iowa fen communities. *Rhodora* **96**: 121–169.

Nelson, W. G. (1993) Beach restoration in the southeastern US: environmental effects and biological monitoring. *Ocean* and Coastal Management 19: 157–182.

Neves, C. F. & Muehe, D. (1995) Potential impacts of sea-level rise on the metropolitan region of Recife, Brazil. *Tournal of Coastal Research* Special Issue 14: 116–131.

New, M. (2002) Trends in freshwater and marine production systems. In: *Production Systems in Fishery Management*, Fisheries Centre Research Report No. 10(8), ed. D. Pauly & M. L. Palomares, pp. 21–27. Vancouver, BC, Canada: University of British Columbia.

Newell, R. C., Lucas, M. I., Velimirov, B. & Seiderer, L. J. (1980) The quantitative significance of dissolved organic losses following fragmentation of kelp (*Ecklonia maxima* and *Laminaria pallida*). *Marine Ecology Progress Series* 2: 45–59.

Newell, R. C., Moloney, C. L., Field, J. G., Lucas, M. I. & Probyn, T. A. (1988) Nitrogen models at the community level: plant-animal-microbe interactions. In: *Nitrogen Cycling in Coastal Marine Environments*, ed. T. H. Blackburn & J. Sorensen, pp. 379–414. New York, USA: John Wiley.

Newey, S. & Seed, R. (1995) The effects of the Braer oil spill on rocky intertidal communities in South Shetland, Scotland. *Marine Pollution Bulletin* **30**: 274–280.

Newton, L. C., Parkes, E. V. H. & Thompson, R. C. (1993) The effects of shell collecting on the abundance of gastropods on Tanzanian shores. *Biological Conservation* 63: 241–245.

Nicholls, R. J. & Branson, J. (1998) Coastal resilience and planning for an uncertain future: an introduction. *Geographical Journal* 164: 255–258.

Nicholls, R. J. & Hoozemans, F. M. J. (1996) The Mediterranean: vulnerability to coastal implications of climate change. Ocean and Coastal Management 31: 105–132.

Nicholls, R. J. & Leatherman, S. P. (1996) Adapting to sea-level rise: relative sea-level trends to 2100 for the United States. *Coastal Management* 24: 301–324.

Nicholls, R. J. & Small, C. (2002) Improved estimates of coastal populations and exposure to hazards released. *Eos* 83: 301–307.

Nicholls, R. J., Hoozemans, F. M. J. & Marchand, M. (1999) Increasing flood risk and wetland losses due to global sea-level rise: regional and global analyses. *Global Environmental Change* 9: 69–87. Nichols, F. H., Cloern, J. E., Luoma, S. N. & Peterson, D. H. (1986) The modification of an estuary. *Science* 231: 567–572.

Nickson, R., McArther, J., Burgees, W., et al. (1998) Arsenic poisoning of Bangladesh groundwater. Nature 395: 338.

Nilsson, C. & Berggren, K. (2000) Alterations of riparian ecosystems caused by river regulation. *BioScience* 50: 783–792.

Nilsson, C., Pizzuto, J. E., Moglen, G. E., et al. (2003) Ecological forecasting and the urbanization of stream ecosystems: challenges for economists, hydrologists, geomorphologists, and ecologists. *Ecosystems* 6: 659–674.

Nilsson, P.Å. (1999) Finjasjön, summer 1999: a summary of monitoring data. Hässleholm, Sweden (in Swedish) [www document]. URL http://www.hassleholmsvatten.se/

Ninio, R., Meekan, M., Done, T. & Sweatman, H. (2000) Temporal patterns in coral assemblages on the Great Barrier Reef from local to large spatial scales. *Marine Ecology Progress Series* 194: 65–74.

Nixon, S. W. (1980) Between coastal marshes and coastal waters: a review of twenty years of speculation and research on the role of salt marshes in estuarine productivity and water chemistry. In: *Estuarine and Wetlands Processes with Emphasis on Modeling*, ed. P. Hamilton & K. B. Macdonald, pp. 437–525. New York, USA: Plenum Press.

Nixon, S. W. (1995) Coastal marine eutrophication: a definition, social causes, and future concerns. *Ophelia* **41**: 199–219.

Nixon, S. W. & Pilson, M. E. (1983) Nitrogen in estuarine and coastal marine ecosystems. In: *Nitrogen in the Marine Environment*, ed. E. J. Carpenter & D. G. Capone, pp. 565–648. New York, USA: Academic Press.

NOAA (1998) Draft Technical Guidance to NMFS for Implementing the Essential Fish Habitat Requirements for the Magnuson-Stevens Act [www document]. URL http:// www.nmfs.noaa.gov/habitat/habitatprotection/ essentialfishhabitat.htm

NOAA (2006) Florida Keys National Marine Sanctuary [www document]. URL http://www.fknms.nos.noaa.gov

Noaksson, E., Linderoth, M., Bosveld, A. T. C., et al. (2003) Endocrine disruption in brook trout (Salvelinus fontinalis) exposed to leachate from a public refuse dump. Science of the Total Environment 305: 87–103.

NODC (2003) World Ocean Circulation Experiment [www document]. URL http://woce.nodc.noaa.gov/wdice

Noe, G.B. & Zedler, J.B. (2001) Spatial and temporal variation of salt marsh seedling establishment in relation to the abiotic and biotic environment. *Journal of Vegetation Science* 12: 61–74.

- Nogueira, F. & Junk, W. J. (2000) Mercury from goldmining in Amazon wetlands: contamination sites, intoxication levels and dispersion pathways. In: *The Central Amazon Floodplain: Actual Use and Options for a Sustainable Management*, ed. W. J. Junk, J. Ohly, M. T. F. Piedade & M. G. M. Soares, pp. 477–503. Leiden, the Netherlands: Backhuvs.
- Norby, R. J. & Cotrufo, M. F. (1998) A question of litter quality. *Nature* **396**: 17–18.
- Nordstrom, K. F. & Mauriello, M. N. (2001) Restoring and maintaining naturally functioning landforms and biota on intensively developed barrier islands under a no-retreat scenario. *Shore and Beach* **69**(3): 19–28.
- Nordstrom, K. F. (1992) *Estuarine Beaches*. London, UK: Elsevier.
- Nordstrom, K. F. (2000) *Beaches and Dunes of Developed Coasts.* Cambridge, UK: Cambridge University Press.
- Nordstrom, K. F. (2003) Restoring naturally functioning beaches and dunes on developed coasts using compromise management solutions: an agenda for action. In: *Values at Sea: Ethics for the Marine Environment*, ed. D. Dallmeyer, pp. 204–229. Athens, GA, USA: University of Georgia Press.
- Nordstrom, K. F. & Roman, C. T., eds. (1996) *Estuarine Shores: Evolution, Environments, and Human Alterations.* New York, USA: John Wiley.
- Nordstrom, K. F., Jackson, N. L., Bruno, M. S. & de Batts, H. A. (2002) Municipal initiative for managing dunes in coastal residential areas: a case study of Avalon, New Jersey, USA. *Geormorphology* 47: 137–152.
- Norkko, A., Thrush, S. F., Hewitt, J. E., et al. (2002) Smothering of estuarine sandflats by terrigenous clay: the role of wind–wave disturbance and bioturbation in sitedependent macrofaunal recovery. *Marine Ecology Progress Series* 234: 23–41.
- Norse, E. A. & Crowder, L. B. (2005) *Marine Conservation Biology*. Washington DC, USA: Island Press.
- North, W. J. (1994) Review of *Macrocystis* biology. In: *Biology* of *Economic Algae*, ed. I. Akatsuka, pp. 447–527. The Hague, the Netherlands: SBP Academic Publishing.
- Northeast Fisheries Science Center (2002) Large marine ecosystems of the world: regional summaries [www document]. URL http://www.edc.uri.edu/lme/ clickable-map.htm
- Norton, T. A. & Benson, M. R. (1983) Ecological interactions between the brown seaweed *Sargassum muticum* and its associated fauna. *Marine Biology* 75: 169–177.
- Notenboom, J. (2001) Managing ecological risks of groundwater pollution. In: *Groundwater Ecology: A Tool for Management of Water Resources*, ed. C. Griebler, D. L.

Danielopol, J. Gibert, H. P. Nachtnebel & J. Notenboom, pp. 248–262. Luxembourg: Office for Official Publications of the European Communities.

- Notenboom, J., Plénet, S. & Turquin, M. J. (1994)
 Groundwater contamination and its impact on groundwater animals and ecosystems. In: *Groundwater Ecology*, ed. J. Gibert, D. L. Danielopol & J. A. Stanford, pp. 477–504.
 San Diego, CA, USA: Academic Press.
- NRC (1993) Managing Wastewater in Coastal Urban Areas. Washington, DC, USA: National Academy Press.
- NRC (1995) Beach Nourishment and Protection. Washington, DC, USA: National Academy Press.
- NRC (1999) Sustaining Marine Fisheries. Washington, DC, USA: National Academy Press.
- NRC (2000a) Clean Coastal Waters: Understanding and Reducing the Effects of Nutrient Pollution. Washington, DC, USA: National Academy Press.
- NRC (2000b) 50 Years of Ocean Discovery. Washington, DC, USA: National Academy Press.
- NRC (2001) Compensating for Wetland Losses under the Clean Water Act. Washington, DC, USA: National Academy Press.
- NRC (2002) *Riparian Areas: Functions and Strategies for Management.* Washington, DC, USA: National Academy Press.
- Nuttle, W. K., Brinson, M. M., Cahoon, D., et al. (1997) Conserving coastal wetlands despite sea level rise. Eos 78: 260–261.
- Nyambe, N. & Breen, C. (2002) Environmental flows, power relations and the use of river system resources, ENVIRO FLOWS, Proceedings of the International Conference on Environmental Flows for River Systems, incorporating the 4th International Ecohydraulics Symposium. 3–8 March, unpublished proceedings, Cape Town, South Africa.
- Nybakken, J. W. (1988) Marine Biology: An Ecological Approach, 2nd edn. New York, USA: Harper & Row.
- Nyman, J. A., DeLaune, R. D., Roberts, H. H. & Patrick, W. H. (1993) Relationships between vegetation and soil formation in a rapidly submerging coastal marsh. *Marine Ecology Progress Series* 96: 269–279.
- Nystrom, M., Folke, C. & Moberg, F. (2000) Coral reef disturbance and resilience in a human-dominated environment. *Trends in Ecology and Evolution* 15: 413–417.
- Ochieng, C. A. & Erftemeijer, P. L. A. (2003) The seagrasses of Kenya and Tanzania. In: *World Atlas of Seagrasses*, ed.
 E. P. Green & F. T. Short, pp. 92–104. Berkeley, CA, USA: University of California Press.

O'Connor, T. P. & Beliaeff, B. (1995) Recent Trends in Coastal Environmental Quality: Results from the Mussel Watch Project, NOAA Technical Report. Rockville, MD, USA: National Oceanic and Atmospheric Administration.

O'Connor, T. P. & Paul, J. F. (2000) Misfit between sediment toxicity and chemistry. *Marine Pollution Bulletin* 40: 59–64.

Odén, S. (1967) Nederbördens försurning. *Dagens Nyheter* (*Stockholm*) 24 October 1967 (in Swedish).

Odum, E. P. (1969) The strategy of ecosystem development. *Science* **164**: 262–270.

Odum, E. P., Finn, J. T. & Franz, E. H. (1979) Perturbation theory and the subsidy-stress gradient. *BioScience* 29: 349-352.

Odum, W. E. & Heald, E. J. (1972) Trophic analysis of an estuarine mangrove community. *Bulletin of Marine Science* 22: 671–738.

Oebius, H. U., Becker, H. J., Rolinski, S. & Jankowski, J. (2001) Parametrization and evaluation of marine environmental impacts produced by deep-sea manganese nodule mining. *Deep-Sea Research II* 48: 3453–3467.

OECD Working Committees (2000) Analytical report on sustainable development: managing natural resources. Unpublished report. Paris, France: OECD General Secretariat.

Oechel, W. C., Cowles, S., Grulke, N., *et al.* (1994) Transient nature of CO₂ fertilization in Arctic tundra. *Nature* **371**: 500–503.

Oechel, W. C., Hastings, S. J., Vourlitis, G., et al. (1993) Recent change of Arctic tundra ecosystems from a net carbon dioxide sink to a source. *Nature* 361: 520–523.

Oechel, W. C., Vourlitis, G., Hastings, S. J. & Bocharev, S. A. (1995) Change in arctic CO₂ flux over 2 decades: effects of climate-change at Barrow, Alaska. *Ecological Applications* 5: 846–855.

Office of Technology Assessment (1986) *Pollutant Discharges* to Surface Waters in Coastal Regions. Washington, DC, USA: Office of Technology Assessment, USA Congress.

Ohly, J. J. & Hund, M. (2000) Floodplain animal husbandry in central Amazonia. In: *The Central Amazon Floodplain: Actual Use and Options for a Sustainable Management*, ed.
W. J. Junk, J. J. Ohly & M. T. F. Piedade, pp. 313–343. Leiden, the Netherlands: Backhuys.

Ohmi, H. (1951) Studies on Isoyake or decrease of seaweeds along the coast of Northern Japan. *Bulletin of the Faculty of Fisheries Hakodate, Hokkaido University* 2: 109–117.

Ojeda, F. & Santelices, B. (1984) Ecological dominance of *Lessonia nigrescens* (Phaeophyta) in central Chile. *Marine Ecology Progress Series* 19: 83–91.

Oleksyn, J. & Reich, P. B. (1994) Pollution, habitat destruction, and biodiversity in Poland. *Conservation Biology* 8: 943–960.

Olivero, A. M. (2001) Classification and Mapping of New York's Calcareous Fen Communities. Albany, NY, USA: New York Natural Heritage Program.

Olsgard, F. & Gray, J. S. (1995) A comprehensive analysis of the effects of offshore oil and gas exploration and production on the benthic communities of the Norwegian continental shelf. *Marine Ecology Progress Series* 122: 277–306.

Olson, D. M. & Dinerstein, E. (1998) The global 200: a representation approach to conserving the earth's most biologically valuable ecosystems. *Conservation Biology* **12**: 502–515.

O'Neill, K. P. (2000) Role of bryophyte-dominated ecosystems in the global carbon budget. In: *Bryophyte Biology*, ed.A. J. Shaw & B. Goffinet, pp. 344–368. Cambridge, UK: Cambridge University Press.

Ong, J. E. (1982) Aquaculture, forestry and conservation of Malaysian mangroves. *Ambio* 11: 252–257.

Ong, J. E. (1993) Mangroves: a carbon source and sink. Chemosphere 27: 1097–1107.

Ong, J. E. (1995) The ecology of mangrove management and conservation. *Hydrobiologia* 295: 343–351.

Ong, J. E., Gong, W. K. & Chan, H. C. (2001) Governments of developing countries grossly undervalue their mangroves? In: Proceedings of the International Symposium on Protection and Management of Coastal Marine Ecosystems, pp. 179–184. Bangkok, Thailand: United Nations Environment Programme.

Ong, J. E., Gong, W. K. & Wong, C. H. (1980) Ecological survey of the Sungai Merbok estuarine mangrove ecosystem. Unpublished report. Penang, Malaysia: Malaysian Fisheries Development Authority (MAJUIKAN).

Onuf, C. P., Phillips, R. C., Moncreiff, C. A., Raz-Guzman, A. & Herrera-Silveira, J. A. (2003) The seagrasses of the Gulf of Mexico. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 244–253. Berkeley, CA, USA: University of California Press.

Öquist, M. G., Svensson, B. H., Groffman, P. & Taylor, M. (1996) Non-tidal wetlands. In: Climate Change 1995: Impacts, Adaptations and Mitigation of Climate Change – Contribution of Working Group II to the Second Assessment Report of the Intergovernmental Panel on Climate Change, ed. R. T. Watson, M. C. Zinoyowera & R. H. Moss, pp. 215–239. Cambridge, UK: Cambridge University Press. O'Reilly, C., Alin, S., Plisnier, P. -D., Cohen, A. & McKee, B. (2003) Climate change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature* 424: 766–768.

Oren, A., ed. (2002) Cellular Origin and Life in Extreme Habitats, vol. 5, Halophilic Microorganisms and Their Environments. Boston, MA, USA: Kluwer.

O'Riordan, T. & Cameron, J., eds. (1994) *Interpreting the Precautionary Principle*. London, UK: Earthscan Publications.

Ormond, R., Bradbury, R., Bainbridge, S., et al. (1988) Test of a model of regulation of crown-of-thorns starfish by fish predators. In: Acanthaster and the Coral Reef: a Theoretical Perspective, ed. R. H. Bradbury, pp. 190–207. Townsville, Queensland, Australia: Springer-Verlag.

Orsi, J. J. & Mecum, W. (1986) Zooplankton distribution and abundance in the Sacramento–San Joaquin Delta in relation to certain environmental factors. *Estuaries* 9: 326–335.

Orth, R. J. & Moore, K. A. (1983) Chesapeake Bay: an unprecedented decline in submerged aquatic vegetation. *Science* 222: 53–57.

Orth, R. J., Fishman, J. R., Wilcox, D. J. & Moore, K. A. (2002) Identification and management of fishing gear impacts in a recovering seagrass system in the coastal bays of the Delmarva Peninsula, USA. *Journal of Coastal Research* Special Issue 37: 111–129.

OSPAR (2000a) Quality Status Report 2000: Region II – Greater North Sea. London, UK: OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic.

OSPAR (2000b) Quality Status Report 2000: Region I – Arctic Waters. London, UK: OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic.

OSPAR (2002) OSPAR Convention for the Protection of the Marine Environment of the North-East Atlantic, Report No. 02/21/1-E. London, UK: OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic.

OSPAR (2003) Declaration of the Joint Ministerial Meeting of the Helsinki and OSPAR Commissions, Report No. JMM 2003/3-E, London, UK: OSPAR Commission for the Protection of the Marine Environment of the North-East Atlantic.

Ostrander, G. K., Armstrong, K. M., Knobbe, E. T., Gerace, D. & Scully, E. P. (2000) Rapid transition in the structure of a coral reef community: the effects of coral bleaching and physical disturbance. *Ecology* **97**: 5297–5302. Osvald, H. (1923) Die Vegetation des Hochmoores Komosse, Svenska Växtsociologiska Sällskapets Handlingar 1. Uppsala, Sweden: Almqvist & Wicksell.

Oudot, J. & Dutrieux, E. (1989) Hydrocarbon weathering and biodegradation in a tropical estuarine ecosystem. *Marine Environmental Research* 27: 195–213.

Ozturk, M., Ozdemir, F. & Yucel, E. (1997) An overview of the environmental issues in the Black Sea region. In: *Scientific, Environmental, and Political Issues in the Circum-Caspian Region*, ed. M. H. Glantz & I. S. Zonn, pp. 213–226. Dordrecht, the Netherlands: Kluwer.

Packham, J. R. & Liddle, M. J. (1970) The Cefni saltmarsh and its recent development. *Field Studies* 3: 331–356.

Paerl, H. W. (1985) Enrichment of marine primary production by nitrogen-enriched acid rain. *Nature* 316: 747–749.

Paerl, H. W. (1995) Coastal eutrophication in relation to atmospheric nitrogen deposition: current perspectives. *Ophelia* 41: 237–259.

Paerl, H. W. & Whitall, D. R. (1999) Anthropogenicallyderived atmospheric nitrogen deposition, marine eutrophication and harmful algal bloom expansion: is there a link? *Ambio* 28: 307–311.

Pagnan, J. L. (2000) Arctic marine protection. *Arctic* 53: 469–473.

Pahl-Wostl, C. (2002) Towards sustainability in the water sector: the importance of human actors and processes of social learning. *Aquatic Sciences* 64: 394–411.

Paijmans, K. (1990) Wooded swamps in New Guinea. In: *Ecosystems of the World*, vol. 15, *Forested Wetlands*, ed. A. E. Lugo, M. Brinson & S. Brown, pp. 335–355. Amsterdam, the Netherlands: Elsevier.

Paine, R. T. (1969) The *Pisaster-Tegula* interaction: prey patches, predator food preference, and intertidal community structure. *Ecology* 50: 950–961.

Paine, R. T. (1980) Food webs: linkages, interaction strength and community infrastructure. *Journal of Animal Ecology* 49: 667–685.

Paine, R. T. (1994) Marine Rocky Shores and Community Ecology: An Experimentalist's Perspective. Luhe, Germany: Ecology Institute.

Paine, R. T., Ruesink, J. L., Sun, A., et al. (1996) Trouble on oiled waters: lessons from the Exxon Valdez oil spill. Annual Review of Ecology and Systematics 27: 197–235.

Paine, R. T., Tenger, M. J. & Johnson, E. A. (1998) Compounded perturbations yield ecological surprises. *Ecosystems* 1: 535–545.

Painting, S. J., Moloney, C.L, Probyn, T. A. & Tibbles, B. (1992) Microhetrotrophic pathways in the southern Benguela upwelling system. *South African Journal of Marine Sciences* **12**: 527–543.

Paling, E. I., van Keulen, M., Wheeler, K., Phillips, J. & Dryberg, R. (2001) Mechanical seagrass transplantation in Western Australia. *Ecological Engineering* 16: 331–339.

PAME (2003) *Arctic Marine Strategic Plan.* Akureyri, Iceland: Protection of the Arctic Marine Environment (PAME) International Secretariat.

Pandolfi, J. M. (1996) Limited membership in Pleistocene reef coral assemblages from the Huon Peninsula, Papua New Guinea: constancy during global changes. *Paleobiology* 22: 152–176.

Pandolfi, J. M. (1999) Response of Pleistocene coral reefs to environmental change over long temporal scales. *American Zoology* 39: 113–130.

Pandolfi, J. M., Bradbury, R. H., Sala, E., *et al.* (2003) Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301: 955–958.

Pang, L. & Pauly, D. (2001) Part 1. Chinese marine capture fisheries from 1950 to the late 1990s: the hopes, the plans and the data. In: *The Marine Fisheries of China: Development and Reported Catches*, Fisheries Centre Research Report No. 9(2), ed. R. Watson, L. Pang & D. Pauly, pp. 1–27. Vancouver, BC, Canada: University of British Columbia.

Pantulu, V. R. (1986) The Mekong river system. In: *The Ecology of River Systems*, ed. B. R. Davies & K. F. Walker, pp. 695–719. The Hague, the Netherlands: Dr W. Junk.

Parker, D., Folland, C. K. & Jackson, M. (1995) Marine surface temperature: observed variations and data requirements. *Climatic Change* 31: 559–600.

Parkinson, C. L. (1992) Spatial patterns of increases and decreases in the length of the sea ice season in the north polar region, 1979–1986. *Journal of Geophysical Research* 97: 14 337–14 388.

Parkinson, C. L. (1994) Spatial patterns in the length of the sea ice season in the Southern Ocean. *Journal of Geophysical Research* 99: 16 327–16 339.

Parkinson, C. L. (1998) Length of the sea ice season in the southern ocean, 1987–1994, in Antarctic sea ice: physical processes, interactions and variability. *Antarctic Research Series* 74: 173–186.

Parkinson, C. L. (2000) Variability of Arctic sea ice: the view from space, an 18-year record. *Arctic* 53: 341–358.

Parkinson, C. L., Cavalieri, D. J., Gloersen, P., Zwally, H. J. & Comiso, J. C. (1999) Arctic sea ice extents, areas and trends, 1978–96. *Journal of Geophysical Research* 104: 20 837–20 856. Parolin, P. (2000) Growth, productivity, and use of trees in white water floodplains. In: *The Central Amazon Floodplain:* Actual Use and Options for a Sustainable Management, ed. W. J. Junk, J. J. Ohly & M. T. F. Piedade, pp. 375–391. Leiden, the Netherlands: Backhuys.

Parrish, R. H. (1995) Lanternfish heaven: the future of world fisheries? *Naga* 18(3): 7–9.

Parrish, R. H. (1998) Life history strategies for marine fishes in the late Holocene. In: *Global versus Local Change in Upwelling Areas*, ed. M. H. Durand, P. Cury, R. Mendelssohn, A. Bakun, C. Roy & D. Pauly, pp. 524–535. Séries Colloques et Séminaires ORSTOM. Paris, France: IRD.

Parsons, J. J. & Bowen, W. A. (1966) Ancient ridged fields of San Jorge River floodplain, Colombia. *Geographical Review* 56: 317–343.

Pasqualini, V., Clabaut, P., Pergent, G., Benyoussef, L. & Pergent-Martini, C. (2000) Contribution of side scan sonar to the management of Mediterranean littoral ecosystems. *International Journal of Remote Sensing* 21: 367–378.

Patrick, R. (1995) Rivers of the United States, vol. 2, Chemical and Physical Characteristics. New York, USA: John Wiley.

Patriquin, D. G. (1975) Migration of blowouts in seagrass beds at Barbados and Carriacou West Indies and its ecological and geological implications. *Aquatic Botany* 1: 163–189.

Paul, A. J., Leavitt, P. R., Schindler, D. W. & Hardie, A. K. (1995) Direct and indirect effects of predation by a calanoid copepod (subgenus: *Hesperodiaptomus*) and of nutrients in a fishless alpine lake. *Canadian Journal of Fisheries and Aquatic Sciences* 52: 2628–2638.

Paul, M. J. & Meyer, J. L. (2001) Streams in the urban landscape. Annual Review of Ecology and Systematics 32: 333–365.

Pauly, D. (1980) On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal du Conseil Internationale pour l'Exploration de la Mer* 39: 175–192.

Pauly, D. (1997) Small scale fisheries in the tropics; marginality, marginalization, and some implications for fisheries management. In: *Global Trends in Fisheries Management*, ed. E. K. Pikitch, P. D. Huppert & M. P. Sissenweiwe, pp. 40–49. Bethesda, MD, USA: American Fisheries Society.

Pauly, D. (2000) Global change, fisheries and the integrity of marine ecosystems: the future has already begun. In: *Ecological Integrity: Integrating Environment, Conservation, and Health*, ed. D. Pimentel, L. Westra & R. F. Ross, pp. 227–239. Washington, DC, USA: Island Press. Pauly, D. (2002) Review of The Skeptical Environmentalist: Measuring the Real State of the World by B. Lomborg. Fish and Fisheries 3: 3–4.

Pauly, D. & Christensen, V. (1995) Primary production required to sustain global fisheries. *Nature* 374: 255–257.

Pauly, D. & Chuenpagdee, R. (2003) Development of fisheries in the Gulf of Thailand Large Marine Ecosystem: analysis of an unplanned experiment. In: Large Marine Ecosystems of the World: Trends in Exploitation, Protection, and Research, ed. G. Hempel & K. Sherman, pp. 337–354. Amsterdam, the Netherlands: Elsevier.

Pauly, D. & Maclean, J. (2003) In a Perfect Ocean: The State of Fisheries and Ecosystems in the North Atlantic Ocean. Washington DC, USA: Island Press.

Pauly, D., Alder, J., Bennett, E., *et al.* (2003) The future for fisheries. *Science* 302: 1359–1361.

Pauly, D., Alder, J., & Watson, R. (2005). Global trends in world fisheries: impacts on marine ecosystems and food security. *Philosophical Transactions of the Royal Society of London B* 360: 5–12.

Pauly, D., Christensen, V., Dalsgaard, J., Froese, R. & Torres Jr, F. (1998) Fishing down marine food webs. *Science* 279: 860–863.

Pauly, D., Christensen, V., Guénette, S., Pitcher, T.J., Sumaila, U. R., Walters, C. J., Watson, R. & Zeller, D. (2002) Toward sustainability in world fisheries. *Nature* 418: 689–695.

Pauly, D., Palomares, M. L., Froese, R., et al. (2001) Fishing down Canadian aquatic food webs. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 51–62.

Payne, A. I. L., Brink, K. H., Mann, K. L. & Hilborn, R., eds. (1992) Benguela trophic functioning. *South African Journal* of Marine Science 12.

Pearce, C. M. & Smith, D. G. (2003) Saltcedar: distribution, abundance, and dispersal mechanisms, northern Montana, USA. *Wetlands* 23: 215–228.

Pearce, F. (2003) Mangrove plantations pose threat to coral reefs. *New Scientist* **178**: 11.

Peck, A. J. & Williamson, D. H. (1987) Effects of forest clearing on groundwater. *Journal of Hydrology* 94: 47–65.

Peltier, W. R. & Tushingham, A. M. (1989) Global sea level rise and the greenhouse effect: might they be connected? *Science* 244: 806–810.

Penang State Economic Planning Unit & DANCED (1999) The Status of and Conservation Plans for the Mangroves of Penang, Integrated Coastal Zone Management Project, Penang Component, Work Group 1: Mangroves, Final Report. Penang, Malaysia: State Economic Planning Unit and Danish Cooperation for Environment and Development.

Penland, S. & Ramsey, K. E. (1990) Relative sea level rise in Louisiana and the Gulf of Mexico: 1908–1988. *Journal of Coastal Research* 6: 323–342.

Percy, J. A. (1999) Whither the waters? Tidal and riverine restrictions in the Bay of Fundy. *Fundy Issues* 11: 1–10.

Percy, J. A. (2000) Salt marsh saga: conserving Fundy's marine meadows. *Fundy Issues* 16: 1–9.

Perelman, S. B., Leon, J. C. & Oesterheld, M. (2001) Cross-scale vegetation patterns of flooding pampa grasslands. *Journal of Ecology* 89: 562–577.

Pergent, G., Mendez, S. & Pergent-Martini, C. (1999) Preliminary data on the impact of fish farming facilities on *Posidonia oceanica* meadows in the Mediterranean. *Oceanologia Acta* 22: 95–107.

Persson, A. (1997) Phosphorus release by fish in relation to external and internal load in a eutrophic lake. *Limnology and Oceanography* 43: 577–583.

Peters, A. F. & Breeman, A. M. (1993) Temperature tolerances and latitudinal range of brown algae from temperate Pacific South America. *Marine Biology* 115: 143–150.

Peters, E. C. (1997) Diseases of coral-reef organisms. In: Life and Death of Coral Reefs, ed. C. Birkeland, pp. 114–139. New York, USA: Chapman & Hall.

Petersen, I., Masters, Z., Hildrew, A. G. & Ormerod, S. J. (2004) Dispersal of adult aquatic insects in catchments of differing land use. *Journal of Applied Ecology* 41: 934–950.

Petersen, I., Winterbottom, J., Orton, S., et al. (1999) Emergence and lateral dispersal of adult Plecoptera and Trichoptera from Broadstone Stream, UK. Freshwater Biology 42: 401–416.

Peterson, C. H. (1985) Patterns of lagoonal bivalve mortality after heavy sedimentation and their paleoecological significance. *Paleobiology* 11: 139–153.

Peterson, C. H. & Estes, J. A. (2001) Conservation and management of marine communities. In: *Marine Community Ecology*, ed. M. D. Bertness, S. D. Gaines & M. E. Hay, pp. 469–508. Sunderland, MA, USA: Sinauer Associates.

Peterson, C. H., Hickerson, D. H. M. & Johnson, G. G. (2000a) Short-term consequences of nourishment and bulldozing on the dominant large invertebrates of a sandy beach. *Journal of Coastal Research* 16: 368–378.

Peterson, C. H., McDonald, L. L., Green, R. H. & Erickson, W. P. (2000b) Sampling design begets conclusions: the statistical basis for detection of injury to and recovery of shoreline communities after the 'Exxon Valdez' oil spill. *Marine Ecology Progress Series* 210: 255–283. Peterson, C. H., Summerson, H. C. & Fegley, S. R. (1983) Relative efficiency of two clam rakes and their contrasting impacts on seagrass biomass. *Fisheries Bulletin* 81: 429–434.

Peterson, C. H., Summerson, H. C. & Fegley, S. R. (1987) Ecological consequences of mechanical harvesting of clams. *Fisheries Bulletin* 85: 281–298.

Pethick, J. (2002) Estuarine and tidal wetland restoration in the United Kingdom: policy versus practice. *Restoration Ecology* 10: 431–437.

Petts, G. E. (1984) Impounded Rivers: Perspectives for Ecological Management. Chichester, UK: John Wiley.

Petts, G. E. (1990) Forested river corridors: a lost resource. In: Water, Engineering, and Landscapes: Water Control and Landscape Transformation in the Modern Period, ed.
D. Cosgrove & G. E. Petts, pp. 12–34. London, UK: Bel-

haven Press.

Pfadenhauer, J. & Kloetzli, F. (1996) Restoration experiments in middle European wet terrestrial ecosystems: an overview. *Vegetatio* **126**: 101–115.

Pfeiffer, W. J. & Wiegert, R. G. (1981) Grazers on Spartina and their predators. In: *The Ecology of a Salt Marsh*, ed. L. R. Pomeroy & R. G. Wiegert, pp. 87–112. New York, USA: Springer-Verlag.

Phillips, R. C. (2003) The seagrasses of the Arabian Gulf and Arabian Region. In: *World Atlas of Seagrasses*, ed.
E. P. Green & F. T. Short, pp. 84–91. Berkeley, CA, USA: University of California Press.

Phillips, V. D. (1998) Peatswamp ecology and sustainable development in Borneo. *Biodiversity and Conservation* 7: 651–671.

Phinn, S. R., Stow, D. A. & Zedler, J. B. (1996) Monitoring wetland habitat restoration in southern California using airborne digital multispectral video data. *Restoration Ecology* 4: 412–422.

Piazzi, L., Balestri, E. & Cinelli, F. (1994) Presence of *Caulerpa racemosa* in the north-western Mediterranean. *Cryptogamie et Algologie* 15: 183–189.

Piechota, T. C., Dracup, J. A. & Fovell, R. G. (1997) Western US streamflow and atmospheric circulation patterns during El Niño–Southern Oscillation. *Journal of Hydrology* 201: 249–271.

Pimbert, M. P. & Gujja, B. (1997) Village voice challenging wetland management policies: experiences in participatory rural appraisal from India and Pakistan. *Nature and Resources* 33: 34–42.

Pimentel, D., Lach, L., Zuniga, R. & Morrison, D. (2000) Environmental and economic costs of nonindigenous species in the United States. *BioScience* 50: 53–65. Pinet, P. R. (2000) Invitation to Oceanography, 2 edn. Boston, MA, USA: Jones & Bartlett.

Pinnegar, J.K., Jennings, S., O'Brien, C. M. & Polunin, N. V. C. (2002) Long-term changes in the trophic level of the Celtic Sea fish community and fish market price distribution. *Journal of Applied Ecology* 39: 377–390.

Pinnegar, J. K., Polunin, N. V. C., Francour, P., et al. (2000) Trophic cascades in benthic marine ecosystems: lessons for fisheries and protected-area management. *Environmental Conservation* 27: 179–200.

Pinnegar, J. K., Viner, D., Hadley, D., *et al.* (2006)
Alternative future scenarios for marine ecosystems: technical report. Unpublished report. Lowestoft, UK: Centre for Environment, Fisheries and Aquaculture Science.

Pitcher, T. (1995) Species changes and fisheries in African lakes: outline of the issues. In: *The Impact of Species Changes in African Lakes*, ed. T. J. Pitcher & P. J. B. Hart, pp. 1–16. London, UK: Chapman & Hall.

Pitcher, T. (2001) Fisheries managed to rebuild ecosystems? Reconstructing the past to salvage the future. *Ecological Applications* 11: 601–617.

Plaziat, J.-C., Cavagnetto, C., Koeniguer, J.-C. & Baltzer, F. (2001) History and biogeography of the mangrove ecosystem, based on a critical reassessment of the paleontological record. *Wetlands Ecology and Management* 9: 161–179.

Plotnikov, I. S., Aladin, N. V. & Philippov, A. A. (1991) The past and present of the Aral Sea Fauna. *Zoologichesky Zhurna* 70(4): 5–15 (in Russian).

PMSEIC (1998) Dryland Salinity and Its Impact on Rural Industries and the Landscape, Prime Minister's Science, Engineering and Innovation Council Occasional Paper No. 1. Canberra, ACT, Australia: Department of Industry, Science and Resources.

Poff, N. L. (1992) Regional hydrologic response to climate change: an ecological perspective. In: *Global Climate Change* and Freshmater Ecosystems, ed. P. Firth & S. G. Fisher, pp. 88–115. Berlin, Germany: Springer-Verlag.

Poff, N. L., Brinson, M. M. & Day Jr, J. W. (2002) Aquatic Ecosystems and Global Climate Change: Potential Impacts on Inland Freshwater and Coastal Wetland Ecosystems in the United States, Report No. 7. Arlington, VA, USA. Pew Center on Global Climate Change.

Poiani, K. A., Richter, B. D., Anderson, M. G. & Richter, H. E. (2000) Biodiversity conservation at multiple scales: functional sites, landscapes, and networks. *BioScience* 50: 133–146.

Poiner, I., Walker, D. I. & Coles, R. B. (1989) Regional studies: seagrasses of tropical Australia. In: *Biology of*

Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region, ed. A.W.D. Larkum, A.J. McComb & S.A. Shepherd, pp. 279–303. Amsterdam, the Netherlands: Elsevier.

Polis, G. A. & Hurd, S. D. (1996) Linking marine and terrestrial food webs: allochthonous inputs from the ocean supports high secondary productivity on small islands and coastal land communities. *American Naturalist* 147: 396–423.

Pollard, D. A. (1984) A review of ecological studies on seagrass–fish communities with particular reference to recent studies in Australia. *Aquatic Botany* 18: 3–42.

Polley, H. W., Wilsey, B. J. & Derner, J. D. (2003) Do species evenness and plant density influence the magnitude of selection and complementarity effects in annual plant species mixtures? *Ecology Letters* 6: 248–256.

Polunin, N., Morales-Nin, B., Pawsey, W., et al. (2001) Feeding relationships in Mediterranean bathyal assemblages elucidated by stable nitrogen and carbon isotope data. *Marine Ecology Progress Series* 220: 13–23.

Polyakov, I. V., Alekseev, G. V., Bekryaev, R. V., et al. (2003a) Long-term ice variability in Arctic marginal seas. *Journal of Climate* 16: 2078–2085.

Polyakov, I. V., Bekryaev, R. V., Alekseev, G. V., et al. (2003b) Variability and trends of air temperature and pressure in the maritime Arctic, 1875–2000. *Journal of Climate* 16: 2067–2077.

Ponce, V. M., ed. (1995) Hydrological and Environmental Impact of the Paraná Paraguay Waterway on the Pantanal of Mato Grosso (Brazil). San Diego, CA, USA: San Diego State University.

Poole, G. C. (2002) Fluvial landscape ecology: addressing uniqueness within the river continuum. *Freshwater Biology* 47: 641–660.

Pope, J. G., Shepherd, J. G. & Webb, J. (1994) Successful surf-riding on size spectra: the secret of survival in the sea. *Philosophical Transactions of the Royal Society of London B* 343: 41–49.

Popp, M. (1995) Salt resistance in herbaceous halophytes and mangroves. *Progress in Botany* 56: 416–429.

Popper, K. R. (1972) Objective Knowledge. Oxford, UK: Oxford University Press.

Porter, J. W., Dustan, P., Jaap, W., et al. (2001) Patterns of spread of coral disease in the Florida Keys. *Hydrobiologia* 460: 1–24.

Postel, S. L. (2000) Entering an era of water scarcity: the challenges ahead. *Ecological Applications* 10: 941–948.

Postel, S. (2001) Growing more food with less water. *Scientific American* 284: 34–37.

Postel, S. L., Daily, G. C. & Ehrlich, P. R. (1996) Human appropriation of renewable fresh water. *Science* 271: 785–788.

Potts, M. (2000) The unmet need for family planning. Scientific American 282: 75–77.

Pounds, J. A. & Crump, M. L. (1994) Amphibian declines and climate disturbance: the case of the golden toad and the harlequin frog. *Conservation Biology* 8: 72–85.

Preen, A. R., Long, W. J. L. & Coles, R. G. (1995) Flood and cyclone related loss, and partial recovery, of more than 1000 km² of seagrass in Hervey Bay, Queensland, Australia. *Aquatic Botany* 52: 3–17.

Preston, C. D., Pearman, D. A. & Dines, T. D. (2002) New Atlas of the British and Irish Flora. Oxford, UK: Oxford University Press.

Pretty, J. N. & Guijt, I. (1992) Primary environmental care: an alternative paradigm for development assistance. *Environment and Urbanization* 4: 22–36.

Primavera, J. H. (1994) Shrimp farming in the Asia-Pacific: environmental and trade issues and regional cooperation [www document]. URL http://www.nautilus.org/papers/ enviro.html/trade/shrimp.html

Pringle, C. M. (2001) Hydrologic connectivity and the management of biological reserves: a global perspective. *Ecological Applications* 11: 981–998.

Pringle, C. M. (2003) What is hydrologic connectivity and why is it ecologically important. *Hydrological Processes* 17: 2685–2689.

Pringle, J. D., Sharp, G. J. & Caddy, J. F. (1980) Proceedings of the workshop on the relationship between sea urchin grazing and commercial plant/animal harvesting. *Canadian Technical Report of Fisheries and Aquatic* Sciences 954.

Prinn, R. G. & Hartley, D. E. (1992) Atmosphere, ocean and land: critical gaps in earth system models. In: *Modeling the Earth System*, ed. D. Ojima, pp. 9–38. Boulder, CO, USA: University Corporation for Atmospheric Research.

Prinn, R. G., Jacoby, H., Sokolov, A., et al. (1999) Integrated global system model for climate policy assessment: feedbacks and sensitivity studies. *Climatic Change* 41: 469–546.

Priskin, J. (2003) Physical impact of four-wheel drive related tourism and recreation in a semi-arid, natural coastal environment. Ocean and Coastal Management 46: 127–155.

Probert, P. K., McKnight, D. G. & Grove, S. L. (1997) Benthic invertebrate bycatch from a deep-water trawl fishery, Chatham Rise, New Zealand. *Aquatic Conservation: Marine and Freshwater Ecosystems* 7: 27–40. Procaccini, G., Buia, M. C., *et al.* (2003) The seagrasses of the Western Mediterranean. In: *World Atlas of Seagrasses*, ed.
E. P. Green & F. T. Short, pp. 56–66. Berkeley, CA, USA: University of California Press.

Proctor, M. C. F. (1997) Aspects of water chemistry in relation to surface degradation on ombrotrophic mires. In: *Blanket Mire Degradation: Causes, Consequences and Challenges*, ed. J. H. Tallis, R. Meade & P. D. Hulme, pp. 140–152. Aberdeen, UK: Macaulay Land Use Research Institute.

Puckridge, J. T., Sheldon, F., Walker, K. F. & Boulton, A. J. (1998) Flow variability and the ecology of large rivers. *Marine and Freshwater Research* 49: 55–72.

Pulfrich, A., Parkins, C. A. & Branch, G. M. (2003a) The effects of shore-based diamond-diving on intertidal and subtidal biological communities and rock lobsters in southern Namibia. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13: 233–255.

Pulfrich, A., Parkins, C. A., Branch, G. M., Bustamante, R. H.
& Velasquez, C. R. (2003b) The effects of sediment deposits from Namibian diamond mines on intertidal and subtidal reefs and rock lobster populations. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13: 257–278.

Purcell, M. A., Malej, A. & Benovic, A. (1999) Potential links of jellyfish to eutrophication and fisheries. In: *Ecosystems at the Land–Sea Margin: Drainage Basin to Coastal Sea*, ed. T. C. Malone, A. Malej, L. W. Harding Jr, N. Smodlana & R. E. Turner, pp. 241–263. Washington, DC, USA: American Geophysical Union.

Pyle, J. A. (2000) Stratospheric ozone depletion: a discussion of our present understanding. In: *Causes and Environmental Implications of Increased UV-B Radiation*, ed. R. E. Hester & R. M. Harrison, pp. 1–16. Cambridge, UK: Royal Society of Chemistry.

Pysek, P. & Prach, K. (1993) Plant invasion and the role of riparian habitats: a comparison of four species alien to central Europe. *Journal of Biogeography* 20: 413–420.

Quinn, F. & Croley, T. (1999) Potential climate change impacts on Lake Erie. In: *State of Lake Erie: Past, Present and Future*, ed. M. Munawar, T. Edsall & I. F. Munawar, pp. 23–30. Leiden, the Netherlands: Backhuys.

Rabalais, N. N. (1992) An Updated Summary of Status and Trends in Indicators of Nutrient Enrichment in the Gulf of Mexico, Report to the Gulf of Mexico Program, Nutrient Enrichment Subcommittee, Publication No. EPA/800-R-92-004. Stennis Space Center, MS, USA: US Environmental Protection Agency, Office of Water, Gulf of Mexico Program. Rabalais, N. N. (2005) Eutrophication. In: *The Sea*, vol. 13, ed. A. R. Robinson & K. H. Brink, pp. 821–866. Cambridge, MA, USA: Harvard University Press.

Rabalais, N. N., Turner, R. E. & Wisemann, W. J. (2002) Hypoxia in the Gulf of Mexico, aka 'The Dead Zone'. Annual Review of Ecology and Systematics 33: 235–263.

Raffaelli, D. & Hawkins, S. (1996) *Intertidal Ecology*. London, UK: Chapman & Hall.

Raimondi, P. T., Barnett, A. M. & Krause, P. R. (1997) The effects of drilling muds on marine invertebrate larvae and adults. *Environmental Toxicology and Chemistry* 16: 1218–1228.

Rakocinski, C. F., Heard, R. W. & LeCroy, S. E. (1996) Responses by macrobenthic assemblages to extensive beach restoration at Perdido Key, Florida, USA. *Journal of Coastal Research* 12: 326–353.

Ramanathan, V., Crutzen, P. J., Kiehl, J. T. & Rosenfeld, D. (2001) Aerosols, climate, and the hydrological cycle. *Science* 294: 2119–2124.

Ramsar (2004) The Ramsar Convention on Wetlands [www document]. URL http://www.ramsar.org/

Ramsar (2006) Criteria for planning a wetland inventory [www document]. URL http://www.ramsar.org/res/lay_ res_viii_06_e.htm

Ramsar & IUCN (1999) Wetlands and global change [www document]. URL http://www.ramsar.org/key_ unfccc_bkgd.htm

Rangeley, R. W. & Kramer, D. L. (1995) Tidal effects on habitat selection and aggregation by juvenile pollock *Pollachius virens* in the rocky intertidal zone. *Marine Ecology Progress Series* 126: 19–29.

Ranwell, D. S. (1972) *Ecology of Salt Marshes and Sand Dunes*. London, UK: Chapman & Hall.

Ranwell, D. S. (1981) Introduced coastal plants and rare species in Britain. In: *Biological Aspects of Rare Plant Conservation*, ed. H. Synge, pp. 413–419. Chichester, UK: John Wiley.

Rasmussen, E. (1977) The wasting disease of eelgrass (*Zostera marina*) and its effects on environmental factors and fauna.
In: *Seagrass Ecosystems*, ed. C. P. McRoy & C. Helfferich, pp. 1–51. New York, USA: Marcel Dekker.

Raven, J. A. (1997) Inorganic carbon acquisition by marine autotrophs. Advances in Botanical Research 27: 185–209.

Raven, J. A., Walker, D. I., Johnston, A. M., Handley, L. L. & Kubler, J. E. (1995) Implications of ¹³C natural abundance measurements for photosynthetic performance by marine macrophytes in their natural environment. *Marine Ecology Progress Series* 123: 193–205. Rawson, D. S. (1946) Successful introduction of fish in a large saline lake. *Canadian Fish Culturalist* 1: 5–8.

Reaka-Kudla, M. L. (1997) The global biodiversity of coral reefs: a comparison with rain forests. In: *Biodiversity II: Understanding and Protecting our Biological Resources*, ed. M. L. Reaka-Kudla, D. E. Wilson & E. O. Wilson, pp. 83–108. Washington, DC, USA: Joseph Henry Press.

Reaka-Kudla, M. L., Feingold, J. S. & Glynn, P. W. (1996) Experimental studies of rapid bioerosion of coral reefs in the

Galapagos Islands. Coral Reefs 15: 101–109.
Reati, G. J., Florin, M., Fernandez, G. J. & Montes, C. (1997) The Laguna de Mar Chiquita (Cordoba, Argentina): a little known, secularly fluctuating, salt lake. International Journal of Salt Lake Research 5: 187–219.

Redclift, M. (1992) Sustainable Development: Exploring the Contradictions. London, UK: Routledge.

Redfield, A. C. (1972) Development of a New England salt marsh. *Ecological Monographs* 42: 201–237.

Reed, D. (2001) Economic Change, Governance and Natural Resource Wealth: The Political Economy of Change in Southern Africa. London, UK: Earthscan Publications.

Reed, D. (2002) Poverty Is Not a Number: The Environment Is Not a Butterfly. Washington, DC, USA: WWF Macroeconomics Program Office.

Reed, D. C. & Foster, M. S. (1984) The effects of canopy shading on algal recruitment and growth in a giant kelp forest. *Ecology* 65: 937–948.

Reeves Jr, C. C. (1978) Economic significance of playa lake deposits. Special Publications of the International Association for Sediments 2: 279–290.

Rehfeldt, G. E., Tchebakova, N. M., Parfenova, Y. I., et al. (2002) Intraspecific responses to climate in *Pinus sylvestris*. *Global Change Biology* 8: 912–929.

Rehfeldt, G. E., Ying, C. C., Spittlehouse, D. L. & Hamilton, D. A. (1999) Genetic responses to climate in *Pinus contorta*: niche breadth, climate change, and reforestation. *Ecological Monographs* 69: 375–407.

Reid, D. & Beeton, A. (1992) Large lakes of the world: a global science opportunity. *Geojournal* 28: 67–72.

Reid, D. & Orlova, M. (2002) Geological and evolutionary underpinnings for the success of Ponto-Caspian species invasions in the Baltic Sea and North American Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1144–1158.

Reid, K. & Croxall, J. P. (2001) Environmental response of upper trophic-level predators reveals a system change in an Antarctic marine ecosystem. *Proceedings of the Royal Society* of London B 268: 377–384. Reilly, F. J. & Bellis, V. J. (1983) The Ecological Impact of Beach Nourishment with Dredged Materials on the Intertidal Zone at Bogue Banks, North Carolina, Miscellaneous Report No. 83–3. Fort Belvoir, VA, USA: US Army Corps of Engineers, Coastal Engineering Research Center.

Reilly, J., Stone, P., Forest, C., *et al.* (2001) Uncertainty and climate change assessments. *Science* 291: 430–433.

Reise, K. (2002) More sand to the shorelines of the Wadden Sea: harmonizing coastal defense with habitat dynamics. In: *Marine Science Frontiers for Europe*, ed. G. Wefer, F. Lamy & F. Mantoura, pp. 203–216. Berlin, Germany: Springer-Verlag.

Rejmánková, E., Pope, K. O., Post, R. & Maltby, E. (1996) Herbaceous wetlands of the Yucatan Peninsula: communities at extreme ends of environmental gradients. *Internationale Revue der Gesamten Hydrobiologie* 81: 225–254.

Renn, C. E. (1936) The wasting disease of *Zostera marina*: a phytological investigation of the diseased plant. *Biological Bulletin* 70: 148–158.

Reschke, C. (1990) Ecological Communities of New York State. Latham, NY, USA: New York Natural Heritage Program, New York State Department of Environmental Conservation.

Retière, C. (1994) Tidal power and the aquatic environment of La Rance. *Biological Journal of the Linnean Society* 51: 25–36.

Revenga, C., Brunner, J., Henninger, N., Kassem, K. & Payne, R. (2000) *Pilot Analysis of Global Ecosystems. Freshwater Systems.* Washington, DC, USA: World Resources Institute.

Revenga, C., Murray, S., Abramovitz, J. & Hammond, A. (1998) Watersheds of the World. Washington, DC, USA: World Resources Institute.

Reyes, J., Sansón, M. & Afonso-Carrillo, J. (1995) Distribution and reproductive phenology of the seagrass *Cymodocea nodosa* (Ucria) Ascherson in the Canary Islands. *Aquatic Botany* **50**: 171–180.

Reynolds, C. S. (1993) Scales of disturbance and their role in plankton ecology. *Hydrobiologia* 249: 157–171.

Ricciardi, A. (2001) Facilitative interactions among invaders: is an 'invasional meltdown' occurring in the Great Lakes? *Canadian Journal of Fisheries and Aquatic Sciences* 58: 2513–2525.

Ricciardi, A. (2003) Predicting the impacts of an introduced species from its invasion history: an empirical approach applied to zebra mussel invasions. *Freshmater Biology* 48: 972–981. Ricciardi, A. (2006) Patterns of invasion in the Laurentian Great Lakes in relation to changes in vector activity. *Diversity and Distributions* 12: 425–433.

Ricciardi, A. & Bourget, E. (1999) Global patterns of macroinvertebrate biomass in marine intertidal communities. *Marine Ecology Progress Series* 185: 21–35.

Ricciardi, A. & MacIsaac, H. (2000) Recent mass invasion of the North American Great Lakes by Ponto-Caspian species. *Trends in Ecology and Evolution* 15: 62–65.

Ricciardi, A. & Rasmussen, J. B. (1999) Extinction rates of North American freshwater fauna. *Conservation Biology* 13: 1220–1222.

Ricciardi, A., Neves, R. J. & Rasmussen, J. B. (1998) Impending extinctions of North American freshwater mussels (Unionoida) following the zebra mussel (*Dreissena* polymorpha) invasion. Journal of Animal Ecology 67: 613–619.

Rice, A. L. & Owen, P. (1998) *Decommissioning the* Brent Spar. London, UK: E. and F.N. Spon.

Richardson, C. J., Reiss, P., Hussain, N. A., Alwash, A. J. & Pool, D. J. (2005) The restoration potential of the Mesopotamian marshes of Irag. *Science* **307**: 1307–1311.

Richardson, D. M., Pyšek, P., Rejmánek, M., et al. (2000) Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distributions* 6: 93–107.

Richardson, J. L. & Richardson, A. E. (1972) History of an African rift lake and its climatic implications. *Ecological Monographs* 42: 499–534.

Richardson, L. L. (1998) Coral diseases: what is really known? Trends in Ecology and Evolution 13: 438–443.

Ridd, P. V., Sandstrom, M. W. & Wolanski, E. (1988) Outwelling from tropical tidal salt flats. *Estuarine, Coastal* and Shelf Science 26: 243–253.

Ridout, P. S., Willcocks, A. D., Morris, R. J., White, S. L. & Rainbow, P. S. (1985) Concentrations of Mn, Fe, Cu, Zn and Cd in the mesopelagic decapod *Systellaspis cebilis* from the East Atlantic Ocean. *Marine Biology* 87: 285–288.

Riedel, R., Caskey, L. & Costa-Pierce, B. A. (2002) Fish biology and fisheries ecology of the Salton Sea, California. *Hydrobiologia* 473: 229–244.

Riemann, B. & Hoffmann, E. (1991) Ecological consequences of dredging and bottom trawling in the Limfjord, Denmark. *Marine Ecology Progress Series* 69: 171–178.

Rigor, I. G., Colony, R. L. & Martin, S. (2000) Variations in surface air temperature observations in the Arctic, 1979–97. *Journal of Climate* 13: 896–914.

Rigor, I. G., Wallace, J. M. & Colony, R. L. (2002) Response of the sea ice to the Arctic Oscillation. *Journal of Climate* 15: 2648–2668. Risager, M. (1998) Impacts of nitrogen on *Sphagnum* dominated bogs with emphasis on critical load assessment. Unpublished Ph.D. thesis, University of Copenhagen, Copenhagen, Denmark.

Risk, M. J. & Sammarco, P. W. (1982) Bioerosion of corals and the influence of damselfish territoriality: a preliminary study. *Oecologia* 52: 376–380.

Risk, M. J. & Sammarco, P. W. (1991) Cross-continental shelf trends in skeletal density of the massive coral *Porites lobata* from the Great Barrier Reef. *Marine Ecology Progress Series* 69: 195–200.

Risk, M. J., Sammarco, P. W. & Edinger, E. N. (1995) Bioerosion in *Acropora* across the continental shelf of the Great Barrier Reef. *Coral Reefs* 14: 79–86.

Ritter, C. & Montagna, P. A. (1999) Seasonal hypoxia and models of benthic response in a Texas Bay. *Estuaries* 22: 7–20.

RIZA (2000) Ecological Gradients in the Danube Delta Lakes, Report No. 2000.015. Lelystad, the Netherlands: RIZA (Institute for Inland Water Management and Waste Water Treatment).

Roberts, C. & Polunin, N. (1993) Marine reserves: simple solutions to managing complex fisheries. *Ambio* 22: 363–368.

Roberts, C. M. & Hawkins, J. P. (1999) Extinction risk in the sea. Trends in Ecology and Evolution 14: 241–246.

Roberts, C. M., McClean, C. J., Veron, J. E. N., *et al.* (2002) Marine biodiversity hotspots and conservation priorities for tropical reefs. *Science* 295: 280–284.

Robertson, A. (1991) Effects of a toxic bloom of *Chrysomulina* polylepis on the common dogwhelk, *Nucella lapillus*, on the Swedish west coast. *Journal of the Marine Biological* Association of the United Kingdom **71**: 569–578.

Robertson, A. I. & Mann, K. H. (1984) Disturbance by ice and life-history adaptations of the seagrass *Zostera marina*. *Marine Biology* 80: 131–141.

Robertson, A. L., Rundle, S. D. & Schmid-Araya, J. M. (2000) Putting the meio- into stream ecology: current findings and future directions for lotic meiofaunal research. *Freshwater Biology* 44: 177–183.

Robinson, C. T., Tockner, K. & Ward, J. V. (2002) The fauna of dynamic riverine landscapes. *Freshmater Biology* 47: 661–677.

Robinson, J. D. F. & Robb, G. A. (1995) Methods for the control and treatment of acid mine drainage. *Coal International* 243: 152–156.

Robinson, T. & Minton, C. (1990) The enigmatic banded stilt. Birds International 1990: 72–85. Rodbell, D. T., Seltzer, G. O., Anderson, D. M., et al. (1999) An ~15 000-year record of El Niño-driven alluviation in Southwestern Ecuador. Science 283: 516–520.

Rodgers, W. A. (1988) Domestic livestock and wildlife conservation: can they coexist? In: *International Rangeland Congress*, ed. P. Singh, V. Shankar & A. K. Srivastava, pp. 639–641. Jhansi, New Delhi, India: Range Management Society of India.

Rodhe, H., Grenfeldt, P., Wisniewski, J., et al. (1995) Acid reign '95? Conference summary statement. Water, Air and Soil Pollution 85: 1–14.

Rodhe, W. & Herrera, R. (1988) *Acidification in Tropical Countries.* New York, USA: John Wiley.

Rodríguez, S. R. (2003) Consumption of drift kelp by intertidal populations of the sea urchin *Tetrapygus niger* on the central Chilean coast: possible consequences at different ecological levels. *Marine Ecology Progress Series* 251: 141–151.

Rodríguez-Itrube, I. (2000) Ecohydrology: a hydrologic perspective of climate-soil-vegetation dynamics. *Water Resources Research* 36: 3-9.

Rodríguez-Itrube, I. (2003) Hydrologic dynamics and ecosystem structure. *Water Science and Technology* 47: 18–94.

Rodwell, M. J., Rowell, D. P. & Folland, C. K. (1999) Oceanic forcing of the wintertime North Atlantic Oscillation and European climate. *Nature* 398: 320–323.

Roelse, P. (1990) Beach and dune nourishment in the Netherlands. Coastal Engineering: Proceedings of the 22nd Coastal Engineering Conference, pp. 1984–1997. New York, USA: American Society of Civil Engineers.

Roemmich, D. (1992) Ocean warming and sea level rise along the southwest US coast. *Science* 257: 373–375.

Rogers, A. D. (1994) The biology of seamounts. Advances in Marine Biology 30: 306–350.

Rogers, A. D. (1999) The biology of Lophelia pertusa (Linnaeus 1758) and other deep-water reef-forming corals and impacts from human activities. International Review of Hydrobiology 84: 315–406.

Rogers, C. S. (1990) Responses of coral reefs and reef organisms to sedimentation. *Marine Ecology Progress Series* 62: 185–202.

Roman, J. & Palumbi, S. R. (2003) Whales before whaling in the North Atlantic. *Science* 301: 508–510.

Ronen, D., Magaritz, M. & Almon, E. (1988) Contaminated aquifers are a forgotten component of the global N₂O budget. *Nature* 335: 57–59. Roosevelt, A. C. (1999) Twelve thousand years of human-environment interaction in the Amazon Floodplain.
In: Várzea: Diversity, Development, and Conservation of Amazonia's Whitewater Floodplains, Advances in Economic Botany, vol. 13, ed. C. Padoch, J. M. Ayres,
M. Pinedo-Vasquez, A. Henderson, pp. 371–392. New York, USA: New York Botanical Garden Press.

Rose, C. S. & Risk, M. J. (1985) Increase in *Cliona delitrix* infestation of *Montastrea cavernosa* heads on an organically polluted portion of the Grand Cayman fringing reef. *Marine Ecology* 6: 345–363.

Rose, K. A. (2000) Why are quantitative relationships between environmental quality of fish populations so elusive? *Ecological Applications* 10: 367–385.

Rose, N., Appleby, P., Boyle, J., MacKay, A. & Flower, R. (1998) The spatial and temporal distribution of fossil-fuel derived pollutants in the sediment record of Lake Baikal. *Journal of Paleolimnology* 20: 151–162.

Rosegrant, M., Cai, X. & Cline, S. (2002) World Water and Food to 2025: Dealing with Scarcity. Washington, DC, USA: International Food Policy Research Institute. Available online at http://www.ifpri.org/pubs

Rosenzweig, M. L. (2001) The four questions: what does the introduction of exotic species do to diversity? *Evolutionary Ecology Research* 3: 361–367.

Rothgeb, J. M. (1996) Foreign Investment and Political Conflict in Developing Countries. Westport, CT, USA: Greenwood Press.

Rothrock, D. A., Yu, Y. & Maykut, G. A. (1999) Thinning of the Arctic sea ice cover. *Geophysical Research Letters* 26: 3469–3472.

Rothrock, D. A., Zhang, J. & Yu, Y. (2003) The Arctic ice thickness anomaly of the 1990s: a consistent view from observations and models. *Journal of Geophysical Research* 108: 3083.

Roughgarden, J., Gaines, S. & Possingham, H. (1988) Recruitment dynamics in complex life cycles. *Science* 241: 1460–1466.

Rowan, D. & Rasmussen, J. (1992) Why don't Great Lakes fish reflect environmental concentrations of organic contaminants? An analysis of between lake variability in the ecological partitioning of PCBs and DDT. *Journal of Great Lakes Research* 18: 724–741.

Rowe, G. T. (1971) Benthic biomass and surface productivity. In: *Fertility of the Sea*, ed. J. D. Costlow, pp. 441–454. New York, USA: Gordon & Breach.

RSPB (2003) *The RSPB's 'No Airport at Cliffe' Campaign*. Sandy, UK: Royal Society for the Protection of Birds. Ruck, K. R. & Cook, P. A. (1998) Sabellid infestations in the shells of South African molluscs: implications for mariculture. *Journal of Shellfish Research* 17: 693–699.

Ruesink, J. L. & Srivastava, D. S. (2001) Numerical and per capita responses to species loss: mechanisms maintaining ecosystem function in a community of stream insect detritivores. *Oikos* 93: 221–234.

Ruiz, G. M. & Carlton, J. T. (2003) Invasive Species: Vector and Management Strategies. Washington, DC, USA: Island Press.

Ruiz, G. M., Fofonoff, P., Hines, A. H. & Grosholz, E. D. (1999) Non-indigenous species as stressors in estuarine and marine communities: assessing invasion impacts and interactions. *Limnology and Oceanography* 44: 950–972.

Ruiz-Luna, A. & Berlanga-Robles, C. A. (2003) Land use, land cover changes and coastal lagoon surface reduction associated with urban growth in northwest Mexico. *Landscape Ecology* 18: 159–171.

Rumbold, D. G., Davis, P. W. & Perretta, C. (2001) Estimating the effect of beach nourishment on *Caretta caretta* (loggerhead sea turtle) nesting. *Restoration Ecology* 9: 304–310.

Rundle, S. D., Bilton, D. T. & Shiozawa, D. K. (2000) Global and regional patterns in lotic meiofauna. *Freshwater Biology* 44: 123–134.

Russ, G. R. (2004) Marine reserves: long-term protection is required for full recovery of predatory fish populations. *Oecologia* 138: 622–627.

Russ, G. R. & Zeller, D. (2003) From Mare Liberum to Mare Reservarum. Marine Policy 27: 75–78.

Ryan, P. (1988) The characteristics and distribution of plastic particles on the sea surface off the southwestern Cape Province, South Africa. *Marine Environmental Research* 25: 249–273.

Rydin, H. & Clymo, R. S. (1989) Transport of carbon and phosphorus compounds about *Sphagnum*. Proceedings of the Royal Society of London B 237: 63–84.

Rysgaard, S., Nielsen, T. G. & Hansen, B. W. (1999) Seasonal variation in nutrients, pelagic primary production and grazing in a high-Arctic coastal marine ecosystem, Young Sound, Northeast Greenland. *Marine Ecology Progress Series* 179: 13–25.

Ryther, J. (1969) Photosynthesis and fish production in the sea. *Science* **166**: 72–76.

Ryther, J. H. & Dunstan, W. N. (1971) Nitrogen, phosphorus and eutrophication in the coastal marine environment. *Science* 171: 1008–1013. Sabo, J. L., Sponseller, R., Dixon, M., et al. (2005) Riparian zones increase regional species diversity by harboring different, not more, species. *Ecology* 86: 56–62.

Sadovy, Y. & Cheung, W. L. (2003) Near extinction of a highly fecund fish: the one that nearly got away. *Fish and Fisheries* 4: 86–99.

Saenger, P. (1998) Mangrove vegetation: an evolutionary perspective. *Marine and Freshwater Research* 49: 277–286.

Saenger, P. (2002) Mangrove Ecology, Silviculture and Conservation. Dordrecht, the Netherlands: Kluwer.

Saenger, P., Hegerl, E. J. & Davie, J. D. S. (1983) Global status of mangrove ecosystems. *Environmentalist* 3(Suppl.): 1–88.

Sainsbury, K. J. (1988) The ecological basis of multispecies fisheries and management of a demersal fishery in tropical Australia. In: *Fish Population Dynamics*, ed. J. Gulland, pp. 349–382. Chichester, UK: John Wiley.

Saintilan, N. & Williams, R. J. (1999) Mangrove transgression into saltmarsh environments in south-east Australia. *Global Ecology and Biogeography* 8: 117–124.

Saji, N. H., Goswami, B. N., Vinayachandran, P. N. & Yamagata, T. (1999) A dipole mode in the tropical Indian Ocean. *Nature* 401: 360–363.

Sakaguchi, Y. (1986) *Rivers in Japan*. Tokyo, Japan: Iwanami Shoten (in Japanese).

Sala, E., Boudouresque, C. F. & Harmelin-Viven, M. (1998) Fishing, trophic cascades, and the structure of algal assemblages: evaluation of an old but untested paradigm. *Oikos* 82: 425–439.

Sala, O. E., Chapin III, F. S., Armesto, J. J., et al. (2000) Global biodiversity scenarios for the year 2100. Science 287: 1770–1774.

Salama, R. B., Farrington, P., Bartle, G. A. & Watson, G. D. (1993) Salinity trends in the wheatbelt of Western Australia: results of water and salt balance studies from Cubaling catchment. *Journal of Hydrology* 145: 41–63.

Salih, A., Larkum, A., Cox, G., Kuhl, M. & Hoegh-Guldberg, O. (2000) Fluorescent pigments in corals are photoprotective. *Nature* 408: 850–853.

Salm, R. V. & Coles, S. L. (2001) Coral Bleaching and Marine Protected Areas. Honolulu, HI, USA: The Nature Conservancy, Asia Pacific Coastal Marine Program.

Saltonstall, K. (2002) Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceedings of the National Academy of Sciences of the USA* 99: 2445–2449.

Saltonstall, K. (2003) Genetic variation among North American populations of *Phragmites australis*: implications for management. *Estuaries* **26**: 444–451. Sammarco, P. W. (1982) Echinoid grazing as a structuring force in coral communities: whole reef manipulations. *Journal of Experimental Marine Biology and Ecology* 61: 31–35.

Sammarco, P. W. (1996) Comments on coral reef regeneration, bioerosion, biogeography, and chemical ecology: future directions. *Journal of Experimental Marine Biology and Ecology* 200: 135–168.

Sammarco, P. W., Risk, M. J. & Rose, C. (1987) Effects of grazing and damselfish territoriality on internal bioerosion of dead corals: indirect effects. *Journal of Experimental Marine Biology and Ecology* 112: 185–199.

Sampat, P. (2000) Deep Trouble: The Hidden Threat of Groundwater Pollution, World Watch Paper No. 154. Washington, DC, USA: World Watch Institute.

San Francisco Estuary Project (1998) *State of the Estuary:* 1992–1997, Technical Report. Oakland, CA, USA: San Francisco Estuary Project.

Sánchez, R. (2002) The role of science in an unsteady, marketdriven fishery: the Patagonian case. In: Oceans 2020: Science for Future Needs, ed. J. G. Field, G. Hempel & C. P. Summerhayes, pp. 114–116. Washington, DC, USA: Island Press.

Sandalow, D. B. & Bowles, I. A. (2001) Fundamentals of treaty-making on climate change. *Science* 292: 1839–1840.

Sanderson, E. W., Jaitheh, M., Levy, M. A., et al. (2002) The human footprint and the last of the wild. *BioScience* 52: 891–904.

Santelices, B. & Ojeda, F. P. (1984a) Effects of canopy removal on the understory algal community structure of coastal forests of *Macrocystis pyrifera* from southern South America. *Marine Ecology Progress Series* 14: 165–173.

Santelices, B. & Ojeda, F. P. (1984b) Population dynamics of coastal forests of *Macrocystis pyrifera* in Puerto Toro, Isla Navarino, Southern Chile. *Marine Ecology Progress Series* 14: 175–183.

Sargeant, H. J. (2001) Vegetation Fires in Sumatra, Indonesia: Oil Palm Agriculture in the Wetlands of Sumatra – Destruction or Development?, European Union Forest Fire Prevention and Control Project with Dinas Kehutanan Propinsi Sumatera Selatan. Jakarta, Indonesia: European Union and Ministry of Forestry.

Sax, D. F. & Gaines, S. D. (2003) Species diversity: from general decrease to local increase. *Trends in Ecology and Evolution* 18: 561–566.

Sayce, K. (1988) Introduced cordgrass Spartina alterniflora Loisel. In: Salt Marshes and Tidelands of Willipa Bay, Washington, Final Report, US Fish and Wildlife Service, USFWS FWSI-87058 TS. Ilwaco, WA: National Wildlife Réfuge.

Scatolini, S. R. & Zedler, J. R. (1996) Epibenthic invertebrates of natural and constructed marshes of San Diego Bay. *Wetlands* 16: 24–37.

Scheffer, M., Carpenter, S., Foley, J. A., Folke, C. & Walter, B. (2001) Catastrophic shifts in ecosystems. *Nature* 413: 591–596.

Scheibling, R. E., Hennigar, A. W. & Balch, T. (1999) Destructive grazing, epiphytism, and disease: the dynamics of sea urchin-kelp interactions in Nova Scotia. *Canadian Journal of Fisheries and Aquatic Sciences* 56: 1–15.

Scherrer, P. & Mille, G. (1989) Biodegradation of crude oil in an experimentally polluted peaty mangrove soil. *Marine Pollution Bulletin* 20: 430–432.

Schiel, D. R. (1990) Macroalgal assemblages in New Zealand: structure, interactions and demography. *Hydrobiologia* 192: 59–76.

Schiel, D. R., Andrew, N. L. & Foster, M. S. (1995) The structure of subtidal algal and invertebrate assemblages at the Chatham Islands, New Zealand. *Marine Biology* 123: 355–367.

Schimel, D. S. (1995) Terrestrial ecosystems and the carbon cycle. *Global Change Biology* 1: 77–91.

Schimel, D. S. & Baker, D. (2002) The wildfire factor. *Nature* **420**: 29–30.

Schindler, D. W. (2001) The cumulative effects of climate warming and other human stresses on Canadian freshwaters in the new millennium. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 18–29.

Schindler, D. W., Curtis, P. J., Parker, B. R. & Stainton, M. P. (1996) Consequences of climate warming and lake acidification for UV-B penetration in North American boreal lakes. *Nature* 379: 705–708.

Schlesinger, W. H. (1997) *Biogeochemistry: An Analysis of Global Change*, 2nd edn. New York, USA: Academic Press.

Schleyer, M. & Celliers, L. (2002) A consideration of biodiversity and future of southern African coral reefs. In: *Coral Reef Degradation in the Indian Ocean: Status Report* 2002, ed. O. Linden, D. Souter, D. Wilhelmsson & D. Obura. pp. 83–90. Kalmar, Sweden: CORDIO.

Schmidt, T. C., Morgenroth, E., Schirmer, M., Effenberger, M. & Haderlein, S. B. (2001) Use and occurrence of fuel oxygenates in Europe. In: Oxygenates in Gasoline: Environmental Aspects, ed. A. F. Diaz & D. L. Drogos, pp. 58–79. Washington, DC, USA: American Chemical Society.

Schmitz, D. C. & Simberloff, D. (1997) Biological invasions: a growing threat. *Issues in Science and Technology* 13: 33–40.

- Schneider, G. & Van Dijk, D. (1997) Environmental management measures for preventing further deterioration of African freshwater bodies: the case of the Lake Victoria Basin. In: *African Inland Fisheries, Aquaculture and the Environment*, ed. K. Remane, pp. 296–304. Oxford, UK: Fishing News Books.
- Schneider, S. H. (1992) Introduction to climate modeling. In: *Climate System Modeling*, ed. K. Trenberth, pp. 3–26. Cambridge, UK: Cambridge University Press.
- Schnitter, J. J. (1994) A History of Dams: The Useful Pyramids. Rotterdam, the Netherlands: A.A. Balkema.
- Schofield, N. J., Stoneman, G. L. & Loh, I. C. (1989) Hydrology of the jarrah forest. In: *The Jarrah Forest: A Complex Mediterranean Ecosystem*, ed. B. Dell, J. J. Havel & N. Malajczuk, pp. 179–201. Dordrecht, the Netherlands: Kluwer.
- Schofield, P. J. & Chapman, L. J. (1999) Interactions between Nile perch, *Lates niloticus*, and other fishes in Lake Nabugabo, Uganda. *Environmental Biology of Fishes* 55: 343–358.
- Schonbeck, M. & Norton, T. (1978) Factors controlling the upper limits of fucoid algae. *Journal of Experimental Marine Biology and Ecology* 31: 303–313.
- Schramm, W. (1996) The Baltic Sea and its transition zones. In: Marine Benthic Vegetation: Recent Changes and the Effects of Eutrophication, ed. W. Schramm & P. Nienhuis, pp. 131–164. Berlin, Germany: Springer-Verlag.
- Schroeder, R. A., Orem, W. H. & Kharaka, Y. K. (2002) Chemical evolution of the Salton Sea, California: nutrient and selenium dynamics. *Hydrobiologia* 473: 23–45.
- Schroeder, R. A., Setmire, J. G. & Wolfe, J. C. (1988) Trace elements and pesticides in the Salton Sea area, California. In: *Proceedings of Conference on Planning Now for Irrigation and Drainage*, Lincoln, NE, 19–21 July, pp. 700–707. New York, USA: American Society of Civil Engineers.
- Schuijt, K. (2002) Land and Water Use of Wetlands in Africa: Economic Value of African Wetlands. Laxenburg, Austria: Institute for Applied System Analyses.
- Schulte-Wülwer-Leidig, A. (1995) Ecological master plan for the Rhine catchment. In: *The Ecological Basis for River Management*, ed. D. M. Harper & A. J. D. Fergusen, pp. 505–514. Chichester, UK: John Wiley.
- Schultze, E.-D. & Mooney, H. A., eds. (1993) Biodiversity and Ecosystem Function. Berlin, Germany: Springer-Verlag.
- Schwartzlose, R. A., Alheit, J., Bakun, A., et al. (1999) Worldwide large-scale fluctuations of sardine and anchovy populations. South African Journal of Marine Science 21: 289–347.

- Scott, D. A. (1992) Asia and the Middle East. In: *Wetlands*, ed.M. Finlayson & M. Moser, pp. 149–178. Oxford, UK: Facts on File.
- Scott, D. A. (1993) Wetlands of West Asia: a regional overview. In: Wetland and Waterfowl Conservation in South and West Asia, Proceedings of an International Symposium, ed. M. Moser & J. van Vessem, pp. 9–22. Karachi, Pakistan: International Waterfowl and Wetlands Research Bureau.
- Scott, D. A., ed. (1995) A Directory of Wetlands in the Middle East. Gland, Switzerland: IUCN, and Slimbridge, UK: IWRB.
- Scott, D. A. & Jones, T. A. (1995) Classification and inventory of wetlands: a global overview. *Vegetatio* 118: 3–16.
- Scott, M. L. & Auble, G. T. (2002) Conservation and restoration of semiarid riparian forests: a case study from the Upper Missouri River, Montana. In: *Flood Pulsing* and Wetlands: Restoring the Natural Balance, ed.
- B. A. Middleton, pp. 145–190. New York, USA: John Wiley. Sculthorpe, C. D. (1968) *Biology of Aquatic Vascular Plants*.
- London, UK: Edward Arnold.
- Seabloom, E. W. & Van der Valk, A. G. (2003) Plant diversity, composition, and invasion of restored and natural prairie pothole wetlands: implications for restoration. *Wetlands* 23: 1–12.
- Seckler, D., Upali, A., Molden, D., de Silva, R. & Barker, R. (1998) World Water Demand and Supply, 1990 to 2025: Scenarios and Issues, Research Report No. 19. Colombo, Sri Lanka: International Water Management Institute.
- Sedell, J. R., Steedman, R. J., Regier, H. A. & Gregory, S. V. (1991) Restoration of human impacted land-water ecotones. In: Ecotones: The Role of Landscape Boundaries in the Management and Restoration of Changing Environments, ed. M. M. Holland, P. G. Risser & R. J. Naiman, pp. 105–129. New York, USA: Chapman & Hall.
- Seehausen, O., Van Aphen, J. & White, F. (1997) Cichlid fish diversity threatened by eutrophication that curbs sexual selection. *Science* 277: 1808–1811.
- Seidenschwarz, I. A., ed. (1986) Pioniervegetation von Flussufer und Kulturland im vorandinen Amazonasgebiet Perus: Ein pflanzensoziologischer Vergleich von vorandinem Flussufer und Kulturland. Langen, Germany: TRIOPS Verlag.
- Seigal, P. R. & Wenner, A. M. (1984) Abnormal reproduction of the sand crab *Emerita analoga* in the vicinity of a nuclear generating station in Southern California. *Marine Biology* 80: 341–345.
- Seitzinger, S. P., Kroeze, C., Bouwman, A. F., et al. (2002) Global patterns of dissolved inorganic and particulate nitrogen inputs to coastal systems: recent conditions and future projections. *Estuaries* 25: 640–655.

Seki, M. (1994) *Rivers of the Earth*. Tokyo, Japan: Soshisya (in Japanese).

Serreze, M. C., Maslanik, J. A., Key, J. R., Kokaly, R. F. & Robinson, D. A. (1995) Diagnosis of the record minimum in Arctic sea ice area during 1990 and associated snow cover extremes. *Geophysical Research Letters* 22: 2183–2186.

Serreze, M. C., Walsh, J. E., Chapin, F. S., et al. (2000) Observational evidence of recent change in the northern high-latitude environment. *Climatic Change* 46: 159–207.

Serruya, C. & Pollingher, U. (1983) *Lakes of the Warm Belt*. Cambridge, UK: Cambridge University Press.

Servant-Vildary, S., Servant, M. & Jimenez, O. (2001) Holocene hydrological and climatic changes in the southern Bolivian Altiplano according to diatom assemblages in paleowetlands. *Hydrobiologia* 466: 267–277.

Seymour, E. A. & Bergheim, A. (1991) Towards a reduction of pollution from intensive aquaculture with reference to the farming of salmonids in Norway. *Aquacultural Engineering* 10: 73–88.

Seymour, R. J., Tegner, M. J., Dayton, P. K. & Parnell, P. E. (1989) Storm wave induced mortality of giant kelp, *Macrocystis pyrifera*, in Southern California. *Estuarine*, *Coastal and Shelf Science* 28: 277–292.

Sægrov, H., Hobæk, A. & Lábe-Lund, H. H. (1996) Vulnerability of melanic *Daphnia* to brown trout predation. *Journal of Plankton Research* 18: 2113–2118.

Shah, T., Molden, D., Sakthvadivel, R. & Seckler, D. (2000) The Global Groundwater Situation: Overview of Opportunities and Challenges. Colombo, Sri Lanka: International Water Management Institute. Available online at www.cgiar.org/ iwmi/pubs/WWVisn/GrWater.htm

Shank, T. M., Fornari, D. J., Von Damm, K. L., et al. (1998) Temporal and spatial patterns of biological community development at nascent deep-sea hydrothermal vents (9° 50' N, East Pacific Rise). Deep-Sea Research II 45: 465–515.

Shannon, L. V., Lutjeharms, J. R. E. & Nelson, G. (1990) Causative mechanisms for intra- and interannual variability in the marine environment around southern Africa. *South African Journal of Science* 86: 356–373.

Sharma, R., Nagender Nath, B., Parthiban, S. & Jai Sankar, S. (2001) Sediment redistribution during simulated benthic disturbance and its implications on deep seabed mining. *Deep-Sea Research I* 48: 3363–3380.

Shatalov, V. (2001) The floodplain forests in the European part of the Russian federation. In: *The Floodplain Forests in Europe*, ed. E. Klimo & H. Hager, pp. 185–201. Leiden, the Netherlands: E. J. Brill. Shaw, A. J. (2000) Phylogeny of the Sphagnopsida based on chloroplast and nuclear DNA sequences. *Bryologist* 103: 277–306.

Shaw, D. G. (1992) The Exxon Valdez oil-spill: ecological and social consequences. *Environmental Conservation* 19: 253–258.

Shaw, D. G. & Day, R. H. (1994) Color- and form-dependent loss of plastic micro-debris from the North Pacific Ocean. *Marine Pollution Bulletin* 28: 39–45.

Sheaffer, J. R., Mullan, J. D. & Hinch, N. B. (2002) Encouraging wise use of floodplains with market-based incentives. *Environment* 44: 33–43.

Shears, N. I. & Babcock, R. I. (2002) Marine reserves demonstrate top-down control of community structure on temperate reefs. *Oecologia* 132: 131–142.

Sheppard, C. R. C. (2000) The Chagos Archipelago. In: Coral Reefs of the Indian Ocean: Their Ecology and Conservation, ed. T. R. McClanahan, C. R. C. Sheppard & D. O. Obura, pp. 445–470. New York, USA: Oxford University Press.

Sherman, D. J., Barron, K. M. & Ellis, J. T. (2002) Retention of beach sands by dams and debris basins in Southern California. *Journal of Coastal Research* Special Issue 36: 662–674.

Sherman, D. J., Nordstrom, K. F., Jackson, N. L. & Allen, J. R. (1994) Sediment mixing depths on a low-energy reflective beach. *Journal of Coastal Research* 10: 297–305.

Sherman, K. & Alexander, L. M., eds. (1986) Variability and Management of Large Marine Ecosystems. Boulder, CO, USA: Westview Press.

Sherman, K., Sissenwine, M., Christensen, V., et al. (2005) A global movement toward an ecosystem approach to management of marine resources. *Marine Ecology Progress* Series 300: 275–279.

Shick, J. M., Lesser, M. P. & Jokiel, P. L. (1996) Effects of ultraviolet radiation on corals and other coral reef organisms. *Global Change Biology* 2: 527–545.

Shick, J. M., Romaine-Lioud, S., Ferrier-Pages, C. & Gattuso, J. P. (1999) Ultraviolet-B radiation stimulates shikimate pathway-dependent accumulation of mycosporine-like amino acids in the coral *Stylophora pistillata* despite decreases in its population of symbiotic dinoflagellates. *Limnology and Oceanography* 44: 1667–1682.

Shiklomanov, I. A. (1990) Global water resources. Natural Resources 26: 34–43.

Shiklomanov, I. A. (1998) World Water Resources: A New Appraisal and Assessment for the Twenty-First Century. Paris, France: UNESCO. Shiklomanov, I. A., ed. (1999) World Water Resources: Modern Assessment and Outlook for the Twenty-First Century (summary of the monograph World Water Resources at the Beginning of the Twenty-First Century (Shiklomanov & Rodda (2003), prepared in the framework of IHP-UNESCO). St Petersburg, Russia: Federal Service of Russia for Hydrometeorology and Environmental Monitoring, State Hydrological Institute.

Shiklomanov, I. A. (2000) World water resources and water use: present assessment and outlook for 2025. In: *Water Scenarios*, ed. F. R. Rijsberman, pp. 160–203. London, UK: Earthscan Publications.

Shiklomanov, I. A. & Rodda, J. C., eds. (2003) World Water Resources at the Beginning of the Twenty-First Century. Cambridge, UK: Cambridge University Press.

Shindell, D. T., Miller, R. L., Schmidt, G. A. & Pandolfo, L. (1999) Simulation of recent northern winter climate trends by greenhouse-gas forcing. *Nature* 399: 452–455.

Shine, C. & de Klemm, C. (1999) Wetlands, Water and the Law: Using Law to Advance Wetland Conservation and Wise Use. Gland, Switzerland: IUCN.

Shinn, E. A., Reich, C. D., Hickey, T. D. & Lidz, B. H. (2003) Staghorn tempestites in the Florida Keys. *Coral Reefs* 22: 91–97.

Shinn, E. A., Smith, G. W., Prospero, J. M., et al. (2000) African dust and the demise of Caribbean coral reefs. *Geophysical Research Letters* 27: 3029–3032.

Shipley, F. S. & Kiesling, R. W., eds. (1994) The State of the Bay: A Characterization of the Galveston Bay Ecosystem, Publication No. GBNEP-44. Webster, TX, USA: Galveston Bay National Estuary Program.

Shireman, J. V. & Cichra, C. E. (1994) Evaluation of aquaculture effluents. *Aquaculture* 123: 55–68.

Short, A. D. (1996) The role of wave height, period, slope, tide range and embaymentisation in beach classification: a review. *Revista Chilena de Historia Natural* 69: 589–604.

Short, A. D. & Hesp, P. A. (1982) Wave, beach and dune interactions in southeastern Australia. *Marine Geology* 48: 259–284.

Short, F. T. & Burdick, D. M. (1996) Quantifying eelgrass habitat loss in relation to housing development and nitrogen loading in Waquoit Bay, Massachusetts. *Estuaries* 19: 730–739.

Short, F. T. & Neckles, H. (1999) The effects of global climate change on seagrasses. *Aquatic Botany* 63: 169–196.

Short, F. T. & Short, C. A. (1984) The seagrass filter: purification of coastal water. In: *The Estuary as a Filter*, ed. V. S. Kennedy, pp. 395–413. New York, USA: Academic Press. Short, F. T. & Short, C. A. (2003) Seagrasses of the western North Atlantic. In: *World Atlas of Seagrasses*, ed.
E. P. Green & F. T. Short, pp. 225–233. Berkeley, CA, USA: University of California Press.

Short, F. T. & Wyllie-Echeverria, S. (1996) Natural and human-induced disturbances of seagrass. *Environmental Conservation* 23: 17–27.

Short, F. T. & Wyllie-Echeverria, S. (2000) Global seagrass declines and effect of climate change. In: Seas at the Millennium: An Environmental Evaluation, vol. 3, ed. C. R. C. Sheppard, pp. 10–11. Amsterdam, the Netherlands: Elsevier.

Short, F. T., Burdick, D. M., Granger, S. & Nixon, S. W. (1996) Long-term decline in eelgrass, *Zostera marina* L., linked to increased housing development. In: *Seagrass Biology: Proceedings of an International Workshop*, Rottnest Island, Western Australia, 25–29 January 1996, ed. J. Kuo, R. C. Phillips, D. I. Walker & H. Kirkman, pp. 291–298. Nedlands, WA, Australia: Sciences UWA.

Short, F. T., Burdick, D. M., Short, C. A., Davis, R. C. & Morgan, P. A. (2000) Developing success criteria for restored eelgrass, salt marsh and mud flat habitats. *Ecological Engineering* 15: 239–252.

Short, F. T., Davis, R. C., Kopp, B. S., Short, C. A. & Burdick, D. M. (2002a) Site selection model for optimal restoration of eelgrass, *Zostera marina* L. *Marine Ecology Progress Series* 227: 253–267.

Short, F. T., Ibelings, B. W. & den Hartog, C. (1988) Comparison of a current eelgrass disease to the wasting disease of the 1930s. *Aquatic Botany* 30: 295–304.

Short, F. T., Kopp, B. S., Gaeckle, J. & Tamaki, H. (2002b) Seagrass ecology and estuarine mitigation: a low-cost method for eelgrass restoration. *Japan Fisheries Science* 68: 1759–1762.

Short, F. T., Mathieson, A. C. & Nelson, J. I. (1986) Recurrence of the eelgrass wasting disease at the border of New Hampshire and Maine, USA. *Marine Ecology Progress* Series 29: 89–92.

Short, F. T., McKenzie, L. J., Coles, R. G. & Vidler, K. P. (2002c) SeagrassNet Manual for Scientific Monitoring of Seagrass Habitat. Cairns, Queensland, Australia: Queensland Department of Primary Industries. Available online at www.SeagrassNet.org

Shpigel, M. & Fishelson, L. (1991) Experimental removal of piscivorous groupers of the genus *Cephalopholis* (Serranidae) from coral habitats in the Gulf of Aqaba (Red Sea). *Environmental Biology of Fishes* 31: 131–138. Shuford, W. D., Warnock, N., Molina, K. C. & Sturm, K. K. (2002) The Salton Sea as critical habitat to migratory and resident waterbirds. *Hydrobiologia* 473: 255–274.

Shuisky, Y. D. & Schwartz, M. L. (1988) Human impact and rates of shoreline retreat along the Black Sea coast. *Journal* of Coastal Research 4: 405–416.

Shukla, J. B. & Dubey, B. (1996) Effect of changing habitat on species: application to Keoladeo National Park, India. *Ecological Modelling* 86: 91–99.

Shy, T. L. & Walsh, J. E. (1996) North Pole ice thickness and association with ice motion history. *Geophysical Research Letters* 23: 2975–2978.

Sidaway, R. (1991) A Review of Marina Developments in Southern England. Sandy, UK: Royal Society for the Protection of Birds.

Siebeck, O. (1988) Experimental investigations of UV tolerance in hermatypic corals (Scleractinia). *Marine Ecology Progress Series* **43**: 95–103.

Sienkiewicz, J., Kloss, M. & Grzyb, M. (2001) The floodplain forest ecosystem in Poland. In: *The Floodplain Forests in Europe*, ed. E. Klimo & H. Hager, pp. 249–297. Leiden, the Netherlands: E. J. Brill.

Silliman, B. R. & Bertness, M. D. (2002) A trophic cascade regulates salt marsh primary production. *Proceedings* of the National Academy of Sciences of the USA 99: 10 500–10 505.

Simas, T., Nunes, J. P. & Ferreira, J. G. (2001) Effects of global climate change on coastal salt marshes. *Ecological Modelling* 139: 1–15.

Simberloff, D. (2003) How much information on population biology is needed to manage introduced species? *Conservation Biology* 17: 83–92.

Simberloff, D. & von Holle, B. (1999) Positive interactions of non-indigenous species: invasional meltdown? *Biological Invasions* 1: 21–32.

Simenstad, C. A. & Fresh, K. L. (1995) Influence of intertidal aquaculture on benthic communities in Pacific northwest estuaries: scales of disturbance. *Estuaries* 18: 43–70.

Simenstad, C. A., Estes, J. A. & Kenyon, K. W. (1978) Aleuts, sea otters, and alternate stable-state communities. *Science* **200**: 403–411.

Simmonds, I., Jones, D. A. & Walland, D. J. (1998) Multi-decadal climate variability in the Antarctic region and global change. *Annals of Glaciology* 27: 617–622.

Simon, K. & Townsend, C. R. (2003) The impacts of freshwater invaders at different levels of ecological organization, with emphasis on ecosystem consequences. *Freshwater Biology* 48: 982–994. Simonovic, S. P. & Carson, R. W. (2003) Flooding in the Red River basin: lessons from post flood activities. *Natural Hazards* 28: 345–365.

Sims, N. C. & Thoms, M. C. (2002) What happens when floodplains wet themselves: vegetation response to inundation on the Lower Balonne Floodplain. *International* Association of Hydrological Sciences 276: 226–236.

Sippel, S. J., Hamilton, S. K., Melack, J. M. & Novo, E. M. M. (1998) Passive microwave observations of inundation area and the area/stage relation in the Amazon River floodplain. *International Journal of Remote Sensing* 19: 3055–3074.

Sissenwine, M. P. & Rosenberg, A. A. (1996) Marine fisheries at a critical juncture. In: *Oceanography: Contemporary Readings in Ocean Sciences*, 3rd edn, ed. R. G. Pirie, pp. 293–302. New York, USA: Oxford University Press.

Sissenwine, M. P., Cohen, E. B. & Grosslein, M. D. (1984) Structure of the Georges Bank ecosystem, Rapport et procès-verbaux des réunions. *Journal du Conseil International pour l'Exploration de la Mer* 183: 243–254.

Sival, F. P., Grootjans, A. P., Stuyfzand, P. J. & Verschoore de la Houssaye, T. (1997) Variation in groundwater composition and decalcification depth in a dune-slack: effects on basiphilous vegetation. *Journal of Coastal Conservation* 3: 79–86.

Sivertsen, K. (1997) Geographic and environmental factors affecting the distribution of kelp beds and barren grounds and changes in biota associated with kelp reduction at sites along the Norwegian coast. *Canadian Journal of Fisheries* and Aquatic Sciences 54: 2872–2887.

Sket, B. (1977) Gegenseitige Beeinflussung der Wasserpollution und des Hoehlenmilieus. *Proceedings* of the 6th International Congress of Speleology, Olomouc, Czechoslovakia, 1973, 5: 253–262.

Sket, B. (1999) The nature of biodiversity in hypogean waters and how it is endangered. *Biodiversity and Conservation* 8: 1319–1338.

Sloan, C. E. (1972) Ground-Water Hydrology of Prairie Potholes in North Dakota, US Geological Survey Professional Paper No. 585-C. Washington, DC, USA: US Government Printing Office.

Slootweg, R. & van Schooten, M. L. F. (1995) Partial restoration of floodplain functions at the village level: the experience of Gounougou, Benue Valley, Cameroon. In: *Tropical Freshwater Wetlands*, ed. H. Roggeri, pp. 159–166. Dordrecht, the Netherlands: Kluwer.

Small, M. P. & Gosling, E. M. (2001) Population genetics of a snail species complex in the British Isles: *Littorina saxatilis*

452 References

(Olivi), *L. neglecta* (Bean) and *L. tenebrosa* (Montagu), using SSCP analysis of cytochrome-B gene fragments. *Journal of Molluscan Studies* 67: 69–80.

Smayda, T. J. (1997) Harmful algal blooms: their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnology and Oceanography* 42: 1137–1153.

Smil, V. (1997) Global population and the nitrogen cycle. Scientific American 277: 76–81.

Smith, A.B. (1992) Pastoralism in Africa: Origins and Development Ecology. London, UK: C. Hurst & Co.; and Athens, OH, USA: Ohio University Press.

Smith, A. H., Lopipero, P. A., Bates, M. N. & Steinmaus, C. M. (2002) Arsenic epidemiology and drinking water standards. *Science* 296: 2145–2146.

Smith, C. R. (1994) Tempo and mode in deep-sea benthic ecology: punctuated equilibrium revisited. *Palaios* 9: 3–13.

Smith, C. R. (1999) The biological environment in the nodule provinces of the deep sea. In: Deep-Seabed Polymetallic Nodule Exploration: Development of Environmental Guidelines, pp. 41–68. Kingston, Jamaica: International Seabed Authority.

Smith, C. R. & Baco, A. R. (2003) Ecology of whale falls at the deep-sea floor. Oceanography and Marine Biology Annual Review 41: 311–354.

Smith, C. R. & Demopoulos, A. (2003) Ecology of the deep Pacific Ocean floor. In: *Ecosystems of the World*, vol. 28, *Ecosystems of the Deep Ocean*, ed. P. A. Tyler, pp. 179–218. Amsterdam, the Netherlands: Elsevier.

Smith, C. R. & Rabouille, C. (2002) What controls the mixedlayer depth in deep-sea sediments? The importance of POC flux. *Limnology and Oceanography* 47: 418–426.

Smith, C. R., Berelson, W., Demaster, D. J., et al. (1997) Latitudinal variations in benthic processes in the abyssal equatorial Pacific: control by biogenic particle flux. Deep-Sea Research II 44: 2295–2317.

Smith, C. R., Hoover, D. J., Doan, S. E., et al. (1996) Phytodetritus at the abyssal seafloor across 10° of latitude in the central equatorial Pacific. *Deep-Sea Research II* 43: 1309–1338.

Smith, C. R., Levin, L. A., Hoover, D. J., McMurtry, G. & Gage, J. D. (2000) Variations in bioturbation across the oxygen minimum zone in the northwest Arabian Sea. *Deep-Sea Research II* 47: 227–257.

Smith, C. R., Present, T. M. C. & Jumars, P. A. (1988) Development of Benthic Biological Monitoring Criteria for Disposal of Low-Level Radioactive Waste in the Abyssal Deep Sea, Final Report for EPA Contract No. 68-02-4303. Washington, DC, USA: US Environmental Protection Agency.

Smith, D. C. & Stewart, B. D. (1994) Development of Methods to Age Commercially Important Dories and Oreos, Fisheries Research and Development Corporation Final Report. Queenscliff, Victoria, Australia: Victorian Fisheries Research Institute.

Smith, D. K. (1991) Seamount abundances and size distributions, and their geographic variations. *Reviews in Aquatic Sciences* 5: 197–210.

Smith, J. E. (1968) Torrey Canyon Pollution and Marine Life. Cambridge, UK: Cambridge University Press.

Smith, K. L. & Druffel, E. R. M. (1998) Long time-series monitoring of an abyssal site in the NE Pacific: an introduction. *Deep-Sea Reseach II* 45: 573.

Smith, K. L. & Kaufmann, R. S. (1999) Long-term discrepancy between food supply and demand in the deep eastern north Pacific. *Science* 284: 1174–1177.

Smith, K. L., Kaufmann, R. S., Baldwin, R. J. & Carlucci, A. F. (2001) Pelagic–benthic coupling in the abyssal eastern North Pacific: an 8-year time-series study of food supply and demand. *Limnology and Oceanography* 46: 543–556.

Smith, M. & Mettler, P. (2002) The role of the flood pulse in maintaining *Boltonia decurrens*, a fugitive plant species of the Illinois River floodplain: a case history of a threatened species. In: *Flood Pulsing in Wetlands: Restoring the Natural Balance*, ed. B. A. Middleton, pp. 109–144. New York, USA: John Wiley.

Smith, R. (1872) *Air and Rain*. London, UK: Longman, Green.

Smith, R. C., Aniley, D. G., Baker, K., et al. (1999) Marine ecosystem sensitivity to climate change. *BioScience* 49: 393–404.

Smith, R. C., Baker, K. S., Byers, M. & Stammerjohn, S. E. (1998) Primary productivity of the Palmer Long Term Ecological Research area and the Southern Ocean. *Journal* of Marine Systems 17: 245–259.

Smith, R. C., Fraser, W. R. & Stammerjohn, S. E. (2003a)
Climate variability and ecological response of the marine ecosystem in the Western Antarctic Peninsula (WAP)
region. In: *Climate Variability and Ecological Response*, ed. D. Greenland, D. Gooding & R. C. Smith, pp. 158–173. Oxford, UK: Oxford University Press.

Smith, R. C., Prezelin, B. B., Baker, K. S., et al. (1992) Ozone depletion: ultraviolet radiation and phytoplankton biology in Antarctic waters. *Science* 255: 952–959.

Smith, R. C., Yuan, X., Liu, J., Martinson, D. G. & Stammerjohn, S. E. (2003*b*) The quasi-quintennial time scale climate variability and ecological response. In: *Climate Variability and Ecological Response*, ed. D. Greenland,
D. Gooding & R. C. Smith, pp. 196–206. Oxford, UK: Oxford University Press.

Smith, R. D., Ammann, A., Bartoldus, C. & Brinson, M. M. (1995) An Approach for Assessing Wetland Functions Using Hydrogeomorphic Classification, Reference Wetlands, and Functional Indices, Technical Report No. TR-WRP-DE-9. Vicksburg, MS, USA: Waterways Experiment Station, Army Corps of Engineers. Available online at http://el.erdc.usace.army.mil/wetlands/pdfs/ wrpde9.pdf

Smith, S. (1972) Factors of ecologic succession in oligotrophic fish communities of the Laurentian Great Lakes. *Journal of* the Fisheries Research Board of Canada 29: 717–730.

Smith, S. V. (1984) Phosphorus versus nitrogen limitation in the marine environment. *Limnology and Oceanography* 29: 1149–1160.

Smith, T. J. & Odum, W. E. (1981) The effects of grazing by snow geese on coastal salt marshes. *Ecology* 62: 98–106.

Smith III, T. J., Boto, K. G., Frusher, S. D. & Giddins, R. L. (1991) Keystone species and mangrove forest dynamics: the influence of burrowing by crabs on soil nutrient status and forest productivity. *Estuarine Coastal and Shelf Science* 33: 419–432.

Smith, V. H., Tilman, G. D. & Nekola, J. C. (1999) Eutrophication: impacts of excess nutrient inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100: 179–196.

Smock, L. A., Smith, L. C., Jones, J. B. & Hooper, S. M. (1994) Effects of drought and a hurricane on a coastal headwater stream. *Archiv für Hydrobiologie* 131: 25–38.

Snedaker, S. C. & Araújo, R. J. (1998) Stomatal conductance and gas exchange in four species of Caribbean mangroves exposed to ambient and increased CO₂. *Marine and Freshwater Research* 49: 325–327.

Snedaker, S. C., Meeder, J. F., Ross, M. S. & Ford, R. G. (1994) Discussion of Ellison, J. C. and Stoddart, D. R. (1991) Mangrove ecosystem collapse during predicted sealevel rise: holocene analogues and implications. *Journal of Coastal Research* 10: 497–498.

Snelgrove, P. V. R. & Smith, C. R. (2002) A riot of species in an environmental calm: the paradox of the species-rich deep sea. Oceanography and Marine Biology Annual Review 40: 311–342.

Snow, A. A. & Vince, S. W. (1984) Plant zonation in an Alaskan salt marsh. II. An experimental study of the role of edaphic conditions. *Journal of Ecology* 72: 669–684. Soares, A. G., Schlacher, T. A. & McLachlan, A. (1997) Carbon and nitrogen exchange between sandy beach clams (*Donax serra*) and kelp beds in the Benguela coastal upwelling region. *Marine Biology* 127: 657–664.

Somerville, R. C. J. & Remer, L. A. (1984). Cloud optical thickness feedbacks in the CO₂ climate problem. *Journal of Geophysical Research* 89: 9668–9672.

Sommaruga-Wögrath, S., Koinig, K. A., Schmidt, R., et al. (1997) Temperature effects on the acidity of remote alpine lakes. *Nature* 387: 64–67.

Song, M. & Dong, M. (2002) Clonal plants and plant species diversity in wetland ecosystems of China. *Journal of Vegetation Science* 13: 237–244.

Sophocleous, M. A., ed. (1998) Perspectives on sustainable development of water resources in Kansas. *Kansas Geological Survey Bulletin* 239.

Sophocleous, M. A. (2002) Interactions between groundwater and surface water: the state of the science. *Hydrogeology Journal* 10: 52–67.

Sophocleous, M. A. (2003) Environmental implications of intensive groundwater use with special regard to streams and wetlands. In: *Groundwater Intensive Use: Challenges and Opportunities*, ed. E. Custodio & R. Llamas, pp. 93–112. Lisse, the Netherlands: A.A. Balkema Publishers.

Soto, R. (1985) Efectos del fenómeno El Niño 1982–83. Ecosistemas de la I Región: Investigación Pesquera (Chile) 32: 199–206.

Soulsby, C., Turnbull, D., Langan, S. J., Owen, R. & Hirst, D. (1995) Long-term trends in stream chemistry and biology in Northeast Scotland: evidence for recovery. *Water, Air and Soil Pollution* 85: 689–694.

South Africa, Government of (1998) National Water Act 36. Available online at www.elaw.org/resources/text.asp? ID=1153

Southgate, T., Wilson, K., Myers, A. A. et al. (1984) Recolonization of a rocky shore in S. W. Ireland following a toxic bloom of the dinoflagellate Gyrodinium aureolum. Journal of the Marine Biological Association of the United Kingdom 64: 485–492.

Southward, A. J. (1967) Recent changes in abundance of intertidal barnacles in southwest England: a possible effect of climatic deterioration. *Journal of the Marine Biological* Association of the United Kingdom 47: 81–85.

Southward, A. J. (1980) Western English Channel: an inconstant ecosystem. *Nature* 285: 361–366.

Southward, A. J. (1982) An ecologist's view of the implications of the observed physiological and biochemical effects of petroleum compounds on marine organisms and ecosystems. *Philosophical Transactions of the Royal Society of London B* 297: 241–255.

Southward, A. J. (1991) Forty years of changes in species composition and population density of barnacles on a rocky shore near Plymouth. *Journal of the Marine Biological* Association of the United Kingdom 71: 495–513.

Southward, A. J. & Crisp, D. J. (1954) The distribution of certain intertidal animals around the Irish coast. *Proceedings* of the Royal Irish Academy 57B: 1–29.

Southward, A. J. & Orton, J. H. (1954) The effects of wave-action on the distribution and numbers of the commoner plants and animals living on the Plymouth breakwater. *Journal of the Marine Biological Association of* the United Kingdom 33: 1–19.

Southward, A. J. & Southward, E. C. (1978) Recolonization of rocky shores in Cornwall after use of toxic dispersants to clean up the *Torrey Canyon* spill. *Journal of the Fisheries Research Board of Canada* 35: 682–706.

Southward, A. J., Hawkins, S. J. & Burrows, M. T. (1995) Seventy years of changes in distribution and abundance of zooplankton and intertidal organisms in the western English channel in relation to rising sea temperature. *Journal of Thermal Biology* 20: 127–155.

Southward, A. J., Langmead, O., Hardman-Mountford, N. J., et al. (2005) Long-term oceanographic and ecological research in the western English Channel. Advances in Marine Biology 47: 1–105.

Spalding, M., Taylor, M., Ravilious, C., Short, F. & Green, E. (2003) Global overview: the distribution and status of seagrasses. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 5–26. Berkeley, CA, USA: University of California Press.

Spalding, M. D., Blasco, F. & Field, C. D. (1997) World Mangrove Atlas. Okinawa, Japan: International Society for Mangrove Ecosystems.

Spalding, V. & Jackson, N. L. (2001) Effect of bulkheads on meiofaunal abundance in the foreshore of an estuarine sand beach. *Journal of Coastal Research* 17: 363–370.

Spangler, D. P. (1984) Geologic variability among six cypress domes in North-Central Florida. In: *Cypress Swamps*, ed. K. C. Ewel & H. T. Odum, pp. 60–66. Gainesville, FL, USA: University of Florida Press.

Spanjol, Z., Tikvic, I. & Baricevic, D. (1999) Protected sites of floodplain forest ecosystems in the Republic of Croatia. *Ekológia (Bratislava)* 18: 82–90.

Sparks, R. E., Nelson, J. C. & Yin, Y. (1998) Naturalization of the flood regime in regulated rivers. *BioScience* 48: 706–720. Spence, S. K., Bryan, G. W., Gibbs, P. E., et al. (1990) Effects of TBT contamination on Nucella populations. Functional Ecology 4: 425–432.

Spencer, P. D. & Collie, J. S. (1997) Patterns of population variability in marine fish stocks. *Fisheries Oceanography* 6: 188–204.

Squires, D., Grafton, R. Q., Alam, M. F. & Omar, M. H. (1998) Where the Land Meets the Sea: Intergrated Sustainable Fisheries Development and Artisanal Fishing, Department of Economics Discussion Paper No. 98–26. San Diego, CA, USA: University of California.

Stachowicz, J. J., Whitlatch, R. B. & Osman, R. W. (1999) Species diversity and invasion resistance in a marine ecosystem. *Science* 286: 1577–1579.

Stammerjohn, S. E. & Smith, R. C. (1996) Spatial and temporal variability of western Antarctic Peninsula sea ice coverage. *Antarctic Research Series* 70: 81–104.

Stammerjohn, S. E. & Smith, R. C. (1997) Opposing southern ocean climate patterns as revealed by trends in regional sea ice coverage. *Climatic Change* 37: 617–639.

Stanford, J. A. & Ward, J. V. (1993) An ecosystem perspective of alluvial rivers: connectivity and the hyporheic corridor. *Journal of the North American Benthological Society* 12: 48–60.

Stanley, E. H. & Doyle, M. W. (2003) Trading off: the ecological effects of dam removal. *Frontiers in Ecology and* the Environment 1: 15–22.

Statistics Bureau of Japan (2003) State of the land [www document]. URL http://www.stat.go.jp/data/psi/3.htm (in Japanese).

Stearns, S. C. (1976) Life-history tactics: a review of the ideas. *Quarterly Review of Biology* 51: 3–47.

Steele, J. H. (1974) *The Structure of Marine Ecosystems*. Oxford, UK: Blackwell Scientific Publications.

Steele, J. H. (1978) Spatial Pattern in Plankton Communities. New York, USA: Plenum Press.

Steele, J. H. (1991) Can ecological theory cross the land-sea boundary? *Journal of Theoretical Biology* 153: 425–436.

Steele, J. H. (1995) Can ecological concepts span the land and ocean domains? In: *Ecological Time Series*, ed. T. M. Powell & J. H. Steele, pp. 5–19. New York, USA: Chapman & Hall.

Steele, J. H. (1998) Regime shifts in marine ecosystems. *Ecological Applications* 8: S33–S36.

Steele, J. H. (2004) Regime shifts: reconciling observation and theory. *Progress in Oceanography* 60: 135–141.

Steele, J. H. & Collie, J. S. (2005) Functional diversity and stability of coastal ecosystems. In: *The Sea*, vol. 13, The Global Coastal Ocean: Interdisciplinary Regional Studies and Syntheses, ed. A. R. Robinson & K. Brink, pp. 785–819. Cambridge, MA, USA: Harvard University Press.

Steele, J. H. & Henderson, E. W. (1984) Modeling long-term fluctuations in fish stocks. *Science* 224: 985–987.

Steele, J. H. & Hoagland, P. (2003) Are fisheries sustainable? Fisheries Research 64: 1–3.

Steffani, C. N. & Branch, G. M. (2003a) Growth rate, condition, and shell shape of *Mytilus galloprovincialis*: responses to wave action. *Marine Ecology Progress Series* 246: 197–209.

Steffani, C. N. & Branch, G. M. (2003b) Spatial comparisons of populations of an indigenous limpet *Scutellastra* argenvillei and the alien mussel *Mytilus galloprovincialis* along a gradient of wave energy. *South African Journal of Marine Science* 25: 195–212.

Steffani, C. N. & Branch, G. M. (2003c) Temporal changes in an interaction between an indigenous limpet *Scutellastra* argenvillei and an alien mussel *Mytilus galloprovincialis*: effects of wave exposure. *South African Journal of Marine Science* 25: 213–229.

Steidinger, K. A., Landsberg, J. H., Tomas, C. R. & Vargo, G. A., eds. (2004) *Harmful Algae 2002*. St Petersburg, FL, USA: Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of UNESCO.

Stein, B. A. (2001) A fragile cornucopia assessing the status of US biodiversity. *Environment* 43: 11–22.

Steneck, R. S. (1997) Fisheries-induced biological changes to the structure and function of the Gulf of Maine ecosystem. In: Proceedings of the Gulf of Maine Ecosystem Dynamics Scientific Symposium and Workshop, RARGOM Report No. 91–1, pp. 151–165. Hanover, NH, USA: Regional Association for Research in the Gulf of Maine.

Steneck, R. S. (1998) Human influences on coastal ecosystems: does overfishing create trophic cascades? *Trends in Ecology* and Evolution 13: 429–430.

Steneck, R. S. & Carlton, J. T. (2001) Human alterations of marine communities: students beware! In: *Marine Community Ecology*, ed. M. Bertness, S. Gaines & M. Hay, pp. 445–468. Sunderland, MA, USA: Sinauer Associates.

Steneck, R. S. & Dethier, M. N. (1994) A functional group approach to the structure of algal-dominated communities. *Oikos* 69: 476–498.

Steneck, R. S. & Watling, L. E. (1982) Feeding capabilities and limitations of herbivorous molluscs: a functional group approach. *Marine Biology* 68: 299–319. Steneck, R. S., Graham, M. H., Bourque, B. J., et al. (2002) Kelp forest ecosystem: biodiversity, stability, resilience and their future. *Environmental Conservation* 29: 436–459.

Steneck, R. S., Vavrinec, J. & Leland, A. V. (2004) Accelerating trophic level dysfunction in kelp forest ecosystems of the western North Atlantic. *Ecosystems* 7: 323–331.

Stenson, J. A. E., Svensson, J.-E. & Cronberg, G. (1993) Changes and interactions in the pelagic community in acidified lakes in Sweden. *Ambio* 22: 277–282.

Stephenson, T. A. & Stephenson, A. (1972) Life between the Tidemarks on Rocky Shores. San Francisco, CA, USA: W.H. Freeman.

Stevens, N. E. (1936) Environmental conditions and the wasting disease. *Science* 84: 87–89.

Stevenson, J. C., Rooth, J. E., Kearney, M. S. & Sundberg, K. L. (2000) The health and long-term stability of natural and restored marshes in Chesapeake Bay. In: *Concepts and Controversies in Tidal Marsh Ecology*, ed. M. P. Weinstein & D. A. Kreeger, pp. 709–736. Dordrecht, the Netherlands: Kluwer.

Stevenson, J. C., Ward, L. G. & Kearney, M. S. (1986) Vertical accretion in marshes with varying rates of sea level rise. In: *Estuarine Variability*, ed. D. Wolfe, pp. 2412–2459. San Diego, CA, USA: Academic Press.

Stewart, K. & Diggins, T. (2002) Heavy metals in the sediments of Buffalo River at Buffalo, NY, USA. Verhandlungen Internationale Vereinigung für Limnologie 28: 1262–1266.

Stiassny, M. L. J. (1996) An overview of freshwater biodiversity: with some lessons from African fishes. *Fisheries* 21(9): 7–13.

Stirling, I., Lunn, N. J. & Iacozza, J. (1999) Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climatic change. *Arctic* 52: 294–306.

Stoddard, J. L., Jeffries, D. S., Lükewille, A., et al. (1999) Regional trends in aquatic recovery from acidification in North America and Europe. *Nature* 401: 575–578.

Stokstad, E. (2003) Can well-timed jolts keep out unwanted exotic fish? *Science* 301: 157–158.

Stone, L., Huppert, A., Rajagopalan, B., Bhasin, H. & Loya, Y. (1999) Mass coral reef bleaching: a recent outcome of increased El Niño activity? *Ecological Letters* 2: 325–330.

Storrs, M. J. & Finlayson, M. (1997) Overview of the Conservation Status of Wetlands of the Northern Territory, Report No. 116. Darwin, NT, Australia: Supervising Scientist.

Stouffer, R. J., Manabe, S. & Bryan, K. (1989) Interhemispheric asymmetry in climate response to a gradual increase in atmospheric CO₂. *Nature* 342: 660–662. Strand, J. A. & Weisner, S. B. (2001) Dynamics of submerged macrophyte populations in response to biomanipulation. *Freshmater Biology* 46: 1397–1408.

Stromberg, J. C. & Chew, M. K. (2002) Flood pulses and the restoration of riparian vegetation in the American Southwest. In: *Flood Pulsing and Wetlands: Restoring the Natural Balance*, ed. B. A. Middleton, pp. 11–50. New York, USA: John Wiley.

Sturgess, P. (1992) Clear-felling dune plantations: studies in vegetation recovery. In: *Coastal Dunes: Geomorphology*, *Ecology and Management for Conservation*, ed. R. W. G. Carter, T. G. F. Curtis & M. J. Sheehy-Skeffington,

pp. 339–349. Rotterdam, the Netherlands: A.A. Balkema.

Succow, M. & Lange, E. (1984) The mire types of the German Democratic Republic. In: *European Mires*, ed. P. D. Moore, pp. 149–175. London, UK: Academic Press.

Sugunan, V. V. & Gopal, B. eds. (2006) Wetlands, fisheries and livelihoods. *International Journal of Ecology and Environmental Sciences* Special Issue 32.

Sullivan, T. (2000) *Aquatic Effects of Acid Deposition*. Boca Raton, FL, USA: Lewis Publishers.

Supanawanid, C. & Lewmanomont, K. (2003) The seagrasses of Thailand. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 158–165. Berkeley, CA, USA: University of California Press.

Supanawanid, C., Albertsen, J. O. & Mukai, H. (2001) Methods for assessing the grazing effects of large herbivores on seagrasses. In: *Global Seagrass Research Methods*, ed. F. T. Short & R. G. Coles, pp. 293–312. Amsterdam, the Netherlands: Elsevier.

Suzuki, T., Kuma, K., Kudo, I. & Matsunaga, K. (1995) Iron requirement of the brown macroalgae *Laminaria japonica*, *Undaria pinnatifida* and the crustose coralline algae, and their competition in the northern Japan Sea. *Phycologia* 34: 201–205.

Svensson, B. H., Christensen, T. R., Johansson, E. & Öquist, M. (1999) Interdecadal changes in CO₂ and CH₄ fluxes of a subarctic mire: Stordalen revisited after 20 years. *Oikos* 85: 22–30.

Svensson, G. (1988) Bog development and environmental conditions as shown by the stratigraphy of Store Mosse mire in southern Sweden. *Boreas* 17: 89–111.

Swales, S., Storey, A. W., Roderick, I. D. & Figa, B. S. (1999) Fishes of floodplain habitats of the Fly River System, Papua New Guinea, and changes associated with El Niño droughts and algal blooms. *Environmental Biology of Fishes* 54: 389–404. Swanson, T. (1996) The underlying causes of biodiversity decline: and economic analysis [www document]. URL http://biodiversityeconomics.org/pdf/960401-08.pdf

Sweatman, H., Osborne, K., Smith, L., et al. (2002) Status of coral reefs of Australasia: Australia and Papua New Guinea. In: Status of Coral Reefs of the World: 2002, ed. C. Wilkinson, pp. 163–180. Townsville, Queensland, Australia: Australian Institute of Marine Science.

Sweeney, B. W., Bott, T. L., Jackson, J. K., et al. (2004) Riparian deforestation, stream narrowing, and loss of stream ecosystem services. Proceedings of the National Academy of Sciences of the USA 101: 14132–14137.

Swiss Re (1998) *Floods: An Insurable Risk?* Zurich, Switzerland: Swiss Reinsurance Company.

Syms, C. & Jones, G. P. (1999) Scale of disturbance and the structure of a temperate fish guild. *Ecology* 80: 921–940.

Symstad, A. J., Chapin III, F. S., Wall, D. H., et al. (2003) Long-term and large-scale perspectives on the relationship between biodiversity and ecosystem functioning. *BioScience* 53: 89–98.

Szmant, A. M. (2001) Why are coral reefs world-wide becoming overgrown by algae? 'Algae, algae everywhere, and nowhere a bite to eat!'. *Coral Reefs* 19: 299–302.

Taastrom, H. M. & Jacobsen, L. (1999) The diet of otters (*Lutra lutra* L.) in Danish freshwater habitats: comparisons of prey fish populations. *Journal of Zoology* (*London*) 248: 1–13.

Tabacchi, E. & Planty-Tabacchi, A.-M. (2000) Riparian plant community composition and the surrounding landscape: functional significance of incomers. In: *Riparian Ecology* and Management in Multi-Land Use Watersheds, ed.
P. J. Wigington Jr & R. L. Beschta, pp. 11–16. Corvallis, OR, USA: American Water Resources Association.

TAC/CGIAR (1989) Sustainable Agricultural Production: Implications of an International Agricultural Research, Research and Development Paper No. 4. Rome, Italy: Technical Advisory Committee/ Consultative Group on International Agricultural Research.

Takizawa, M., Straube, R. T. & Colwell, R. R. (1993) Near-bottom pelagic bacteria at a deep-water sewage sludge disposal site. *Applied and Environmental Microbiology* 59: 3406–3410.

Talley, T. S., Crooks, J. A. & Levin, L. A. (2001) Habitat utilization and alteration by the invasive burrowing isopod, *Sphaeroma quoyanum* in California salt marshes. *Marine Biology* 138: 561–573.

Tallis, J. H. (1985) Mass movement and erosion of a southern Pennine blanket peat. *Journal of Ecology* **73**: 283–315. Tallis, J. H. (1987) Fire and flood at Holme Moss: erosion processes in an upland blanket mire. *Journal of Ecology* 75: 1099–1129.

Tallis, J. H. (1998) Growth and degradation of British and Irish blanket mires. *Environmental Reviews* 6: 81–122.

Tam, N. F. Y. & Wong, Y. S. (1997) Accumulation and distribution of heavy metals in a simulated mangrove system treated with sewage. *Hydrobiologia* 352: 67–75.

Tamatamah, R., Duthie, H. & Hecky, R. (2004) The importance of atmospheric deposition to the phosphorus loading of Lake Victoria (East Africa). *Biogeochemistry* 73: 1–20.

Tanabe, S. & Tatsukawa, R. (1983) Vertical transport and residence time of chlorinated hydrocarbons in the open ocean water column. *Journal of the Oceanographical Society* of Japan 39: 53–62.

Tanner, J. E. (1995) Competition between scleractinian corals and macroalgae: an experimental investigation of coral growth, survival and reproduction. *Journal of Experimental Marine Biology and Ecology* 190: 151–168.

Tao, F., Yokozava, M., Hayashi, Y. & Lin, E. (2003) Terrestrial water cycle and the impact of climate change. *Ambio* 32: 295–301.

Tao, W. & Wei, W. (1997) Water resources and agricultural environment in arid regions of China. In: *Freshwater Resources in Arid Lands*, ed. J. Uitto & J. Schneider, pp. 44–60. New York, USA: United Nations Press.

Tapp, J. F., Shillabeer, N. & Ashman, C. M. (1993) Continued observations of the benthic fauna of the industrialized Tees estuary, 1979–1990. *Journal of Experi*mental Marine Biology and Ecology 172: 67–82.

Tarazona, J. & Parendes, C. (1992) Impacto de les eventos El-Niño sobre las comunidades bentonicas de playa arenosa durante 1976–1986. In: *Paleo ENSO Records International Symposium*, Lima, Extended Abstracts, pp. 299–303. Available online at www.crid.or.cr/crid/CD_El_Ni%F1o/ pdf/spa/doc9263/doc9263.htm

Tarazona, J., Arntz, W., Canahuire, E., Ayala, Z. & Robles, A. (1985) Modificaciones producidas durante 'El-Niño' en la infauna bentonica de areas someras del ecosistema de afloriamento peruano. In: *El-Jernomeno El-Niño y su Impacto en la Fauna Marina*, ed. W. Arntz, A. Landa & J. Tarazona, pp. 55–63. Callao, Peru: Bolletín Instituto del Mardel, Special Issue 1985.

Tarazona, J., Salzwedel, H. & Arntz, W. (1988) Oscillations of marcobenthos in shallow waters of the Peruvian central coast induced by the El-Niño 1982–83. *Journal of Marine Research* 46: 593–611. Taylor, A. H. & Stephens, J. A. (1980) Latitudinal displacements of the Gulf Stream (1966 to 1977) and their relation to changes in temperature and zooplankton abundance in the NE Atlantic. *Oceanologica Acta* 3: 145–149.

Taylor, P. (1993) The state of the marine environment: a critique of the work and role of the Joint Group of Experts on Scientific Aspects of Marine Pollution (GESAMP). *Marine Pollution Bulletin* **26**: 120–127.

Taylor, P. R. & Littler, M. M. (1982) The roles of compensatory mortality, disturbance and substrate retention in development and organization of a sand-influenced, rocky-intertidal community. *Ecology* 63: 135–146.

Tegner, M. J. & Dayton, P. K. (1987) El Niño effects on southern California kelp forest communities. *Advances in Ecological Research* 17: 243–279.

Tegner, M. J. & Dayton, P. K. (1991) Sea urchins, El Niños, and the long-term stability of Southern California kelp forest communities. *Marine Ecology Progress Series* 77: 49–63.

Tegner, M. J. & Levin, L. A. (1983) Spiny lobsters and sea urchins: analysis of a predator-prey interaction. *Journal of Experimental Marine Biology and Ecology* 73: 125–150.

Tegner, M. J., Basch, L. V. & Dayton, P. K. (1996a) Near extinction of an exploited marine invertebrate. *Trends in Ecology and Evolution* 11: 278–279.

Tegner, M. J., Dayton, P. K., Edwards, P. B. & Riser, K. L. (1996b) Is there evidence for long-term climatic change in southern California kelp forests? *California Cooperative Oceanic Fisheries Investigations Reports* 37: 111–126.

Tegner, M. J., Dayton, P. K., Edwards, P. B. & Riser, K. L. (1997) Large-scale, low-frequency oceanographic effects on kelp forest succession: a tale of two cohorts. *Marine Ecology Progress Series* 146: 117–134.

Terrados, J., Duarte, C. M., Fortes, M. D., et al. (1998) Changes in community structure and biomass of seagrass communities along gradients of siltation in SE Asia. *Estuarine, Coastal and Shelf Science* 46: 757–768.

Thayer, G. W., Adams, S. M. & LaCroix, M. W. (1975) Structural and functional aspects of a recently established *Zostera marina* community. In: *Estuarine Research*, vol. 1, *Chemistry, Biology, and the Estuarine System*, ed. L. E. Cronin, pp. 518–540. New York, USA: Academic Press.

The Nature Conservancy (2003) *Emiquon: Restoring the Illinois River*. Arlington, VA, USA: The Nature Conservancy. Available online at http://nature.org/wherewework/ northamerica/ states/illinois/preserves/art1112.html

The Times Atlas of the World (1999) London, UK: Collins Bartholomew. Thesiger, W. (1964) *The Marsh Arabs.* London, UK: Penguin Books.

Thiel, H. (2003) Anthropogenic impacts on the deep sea. In: *Ecosystems of the World*, vol. 28, *Ecosystems of the Deep Oceans*, ed. P. A. Tyler, pp. 427–471. Amsterdam, the Netherlands: Elsevier.

Thiel, H., Angel, M. V., Foell, E. J., Rice, A. L. & Schriever,
G. (1998) *Environmental Risks from Large-Scale Ecological Research in the Deep Sea*. Bremerhaven, Germany:
Commission of the European Communities Directorate-General for Science, Research and Development.

Thiel, H., Pfannkuche, O., Schriever, G., *et al.* (1989) Phytodetritus on the deep-sea floor in a central oceanic region of the north-east Atlantic. *Biological Oceanography* 6: 203–239.

Thiel, H., Schriever, G., Ahnert, A., *et al.* (2001) The largescale environmental impact experiment DISCOL: reflection and foresight. *Deep-Sea Research II* 48: 3869–3882.

Thiele, D. & Gill, P. C. (1999) Cetacean observations during a winter voyage into Antarctic sea ice south of Australia. *Antarctic Science* 11: 48–53.

Thom, B. G. (1967) Mangrove ecology and deltaic geomorphology, Tabasco, Mexico. *Journal of Ecology* 55: 301–343.

Thom, B. G. (1982) Mangrove ecology: a geomorphological perspective. In: *Mangrove Ecosystems in Australia*, ed.
B. F. Clough, pp. 3–17. Canberra, ACT, Australia: Australian National University Press.

Thom, R. M., Shrefler, D. K. & MacDonald, K. B. (1994) Shoreline Armoring Effects on Coastal Ecology and Biological Resources in Puget Sound, Washington. Olympia, WA, USA: Shorelands and Coastal Zone Management Program, Washington Department of Ecology.

Thomas, C. A. & Bendell-Young, L. I. (1999) The significance of diagenesis versus riverine input in contributions to the sediment geochemical matrix of iron and manganese in an intertidal region. *Estuarine, Coastal and Shelf Science* 48: 635–647.

Thomas, D. N. & Dieckmann, G. S. (2003) Sea-Ice: An Introduction to its Physics, Chemistry, Biology and Geology. Oxford, UK: Blackwell Science.

Thompson, B., Anderson, B., Hunt, J., Taberski, K. & Phillips, B. (1999) Relationships between sediment contamination and toxicity in San Francisco Bay. *Marine Environmental Research* 48: 285–309.

Thompson, D. W. J. & Solomon, S. (2002) Interpretation of recent Southern Hemisphere climate change. *Science* 296: 895–899. Thompson, D. W. J. & Wallace, J. M. (1998) The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. *Geophysical Research Letters* 25: 1297–1300.

Thompson, D. W. J. & Wallace, J. M. (2000) Annular modes in the extratropical circulation. I. Month-to-month variability. *Journal of Climate* 13: 1000–1016.

Thompson, D. W. J. & Wallace, J. M. (2001) Regional climate impacts of the Northern Hemisphere annular mode. *Science* 293: 85–89.

Thompson, D. W. J., Wallace, J. M. & Hegerl, G. C. (2000) Annular modes in the extratropical circulation. II. Trends. *Journal of Climate* 13: 1018–1036.

Thompson, G. B. & Drake, B. G. (1994) Insect and fungi on a C_3 sedge and a C_4 grass exposed to elevated atmospheric CO_2 concentrations in open-top chambers in the field. *Plant, Cell and Environment* 17: 1161–1167.

Thompson, J. D. (1991) The biology of an invasive plant. BioScience 41: 393-401.

Thompson, J. R. & Polet, G. (2000) Hydrology and land use in a Sahelian floodplain wetland. *Wetlands* **20**: 639–659.

Thompson, J. R. (1996) Africa's floodplains: a hydrologic overview. In: Water Management and Wetlands in Sub-Saharan Africa, ed. M. C. Acreman & G. E. Hollis, pp. 5–20. Gland, Switzerland: IUCN.

Thompson, R. C., Crowe, T. P. & Hawkins, S. J. (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environmental Conservation* 29: 168–191.

Thompson, R. C., Norton, T. A. & Hawkins, S. J. (2004) Physical stress and biological control regulate the balance between producers and consumers in marine intertidal biofilms. *Ecology* 85: 1372–1382.

Thompson, R. C., Roberts, M. F., Norton, T. A. & Hawkins, S. J. (2000) Feast or famine for intertidal grazing molluscs: a mis-match between seasonal variations in grazing intensity and the abundance of microbial resources. *Hydrobiologia* 440: 357–367.

Thormann, M. N. & Bayley, S. E. (1997) Aboveground net primary production along a bog-fen-marsh gradient in southern boreal Alberta, Canada. *Ecoscience* 4: 374–384.

Thresher, R., Proctor, C., Ruiz, G., et al. (2003) Invasion dynamics of the European shore crab, *Carcinus maenas*, in Australia. *Marine Biology* 142: 867–876.

Thrush, S. F. & Dayton, P. K. (2002) Disturbance to marine benthic habitats by trawling and dredging: implications for marine biodiversity. *Annual Review of Ecology and Systematics* 33: 449–473. Thrush, S. F., Hewitt, J. E., Cummings, V. J., *et al.* (1998) Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications* 8: 866–879.

Thrush, S. F., Hewitt, J. E., Norkko, A., Cummings, V. J. & Funnell, G. A. (2003) Catastrophic sedimentation on estuarine sandflats: recovery of macrobenthic communities is influenced by a variety of environmental factors. *Ecological Applications* 13: 1433–1455.

Tibbetts, J. (2002) Coastal cities: living on the edge. Environmental Health Perspectives 110: 674–681.

Tiega, A. (2001) Priorities for wetland biodiversity conservation in Africa. In: Wetland Inventory, Assessment and Monitoring: Practical Techniques and Identification of Major Issues, ed. C. M. Finlayson & N. C. Davidson, pp. 112–120. Darwin, NT, Australia: Supervising Scientist.

Tilman, D., Fargione, J., Wolff, B., et al. (2001) Forecasting agriculturally driven global environmental change. *Science* 292: 281–284.

Tilman, D., Wedin, D. & Knops, J. (1996) Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* **379**: 718–720.

Timchenko, V., Oksijuk, O. & Gore, J. (2000) A model for ecosystem state and water quality management in the Dnieper River delta. *Ecological Engineering* 16: 119–125.

Timms, B. V. (1976) A comparative study of the limnology of three maar lakes in western Victoria. I. Physiography and physicochemical features. *Australian Journal of Marine and Freshwater Research* 27: 35–60.

Tinker, P. B. (1997) The environmental implications of intensified land use in developing countries. *Philosophical Transactions of the Royal Society of London B* 352: 1023–1033.

Titus, J. G. (1990) Greenhouse effect, sea level rise, and barrier islands: case study of Long Beach Island, New Jersey. *Coastal Management* 18: 65–90.

Titus, J. G. & Narayanan, V. K. (1995) The Probability of Sea Level Rise, EPA Report No. 230-R-95-008. Washington, DC, USA: US Environmental Protection Agency.

Tockner, K. & Stanford, J. A. (2002) Riverine flood plains: present state and future trends. *Environmental Conservation* 29: 308–330.

Tockner, K., Malard, F. & Ward, J. V. (2000) An extension of the flood pulse concept. *Hydrological Processes* 14: 2861–2883.

Tockner, K., Pennetzdorfer, D., Reiner, N., Schiemer, F. & Ward, J. V. (1999) Hydrologic connectivity and the exchange of organic matter and nutrients in a dynamic river-floodplain system (Danube, Austria). *Freshmater Biology* 41: 521–535. Tockner, K., Ward, J. V., Arscott, B. A., *et al.* (2003) The Tagliamento River: a model ecosystem of European importance. *Aquatic Sciences* **65**: 239–253.

Todd, C. D. (1998) Larval supply and recruitment of benthic invertebrates: do larvae always disperse as much as we believe? *Hydrobiologia* **376**: 1–21.

Todd, C. D. & Lewis, J. R. (1984) Effects of low air temperature on *Laminaria digitata* L. in southwestern Scotland. *Marine Ecology Progress Series* 16: 199–201.

Todd, C. D., Lambert, W. J. & Thorpe, J. P. (1998) The genetic structure of intertidal populations of two species of nudibranch molluscs with planktotrophic and pelagic lecithotrophic larval stages: are pelagic larvae 'for' dispersal? *Journal of Experimental Marine Biology and Ecology* 228: 1–28.

Todhunter, P. E. & Rundquist, B. C. (2004) Terminal lake flooding and wetland expansion in Nelson County, North Dakota. *Physical Geography* **25**: 68–85.

Tolba, M. K. & El-Kholy, O. A., eds. (1992) The World Environment 1972–1992: Two Decades of Challenge. London, UK: Chapman & Hall and United Nations Environment Programme.

Tolonen, K. (1967) Über die Entwicklung der Moore im finnischen Nordkarelien. Annales Botanici Fennici 4: 219–416.

Tomalin, B. J. & Kyle, R. (1998) Subsistence and recreational mussel (*Perna perna*) collecting in KwaZulu–Natal, South Africa: fishing mortality and precautionary management. *South African Journal of Science* 33: 12–22.

Tomascik, T., Mah, A.J., Nontji, A. & Moosa, M.K. (1997) *The Ecology of the Indonesian Seas*, Part 2. Hong Kong: Periplus.

Tomasko, D. A., Dawes, C. J. & Hall, M. O. (1996) The effects of anthropogenic nutrient enrichment on turtle grass (*Thalassia testudinum*) in Sarasota Bay, Florida. *Estuaries* 19: 448–456.

Tomicic, J. (1985) Efectos del fenómeno El Niño 1982–83 en las comunidades litorales de la península de Mejillones. *Investigación Pesquera (Chile)* 32: 209–213.

Tóth, J. (1963) A theoretical analysis of groundwater flow in small drainage basins. *Journal of Geophysical Research* 68: 4795–4812.

Tóth, J. (1999) Groundwater as a geologic agent: an overview of the causes, processes, and manifestations. *Hydrogeology Journal* 7: 1–14.

Towns, D. R. & Ballantine, W. J. (1993) Conservation and restoration of New Zealand island ecosystems. *Trends in Ecology and Evolution* 8: 452–457. Townsend, C. R. (1996) Concepts in river ecology: pattern and process in the catchment hierarchy. *Archiv für Hydrobiologie* 113 (Suppl.): 3–21.

Townsend, C. R. (2003) Individual, population, community, and ecosystem consequences of a fish invader in New Zealand streams. *Conservation Biology* 17: 38–47.

Townsend, C. R., Dolédec, S., Norris, R., Peacock, K. & Arbuckle, C. J. (2003) The influence of scale and geography on relationships between stream community composition and landscape variables: description and prediction. *Freshwater Biology* 48: 768–785.

Tracey, D. M. & Horn, P. L. (1999) Background and review of ageing orange roughy (*Hoplostethus atlanticus*, Trachichthyidae) from New Zealand and elsewhere. *New Zealand Journal of Marine and Freshwater Research* 33: 67–86.

Travis, J. M. J. (2003) Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proceedings of the Royal Society of London B* 270: 467–473.

Trupin, A. & Wahr, J. (1990) Spectroscopic analysis of global tide gauge sea level data. *Geophysical Journal International* 100: 441–453.

Truscott, A. (1984) Control of Spartina anglica on the amenity beaches of Southport. In: Spartina anglica in Great Britain, ed. P. Doody, pp. 64–69. Attingham Park, UK: Nature Conservation Council.

Tucker, W. B., Weatherly, J. W., Eppler, D. T., Farmer, L. D.
& Bentley, D. L. (2001) Evidence for rapid thinning of sea ice in the western Arctic Ocean at the end of the 1980s. *Geophysical Research Letters* 28: 2851–2854.

Tudhope, A. W., Chilcott, C. P., McColluch, M. T., et al. (2001) Variability in the El Niño–Southern Oscillation through a glacial–interglacial cycle. *Science* 291: 1511–1517.

Tunnicliffe, V., Embley, R. W., Holden, J. F., et al. (1997) Biological colonization of new hydrothermal vents following an eruption on Juan de Fuca Ridge. *Deep-Sea Research I* 44: 1627–1644.

Turetsky, M., Wieder, K., Halsey, L. & Vitt, D. H. (2002) Current disturbance and the diminishing peatland carbon sink. *Geophysical Research Letters* 29: 1400.

Turner, A. (2000) Trace metal contamination in sediments from UK estuaries: an empirical evaluation of the role of hydrous iron and manganese oxides. *Estuarine, Coastal and Shelf Science* 50: 355–371.

Turner, G. (1994) Fishing and the conservation of the endemic fishes of Lake Malawi. Archiv für Hydrobiologie 44: 481–494. Turner, R. E. & Lewis, R. R. (1997) Hydrologic restoration of coastal wetlands. Wetlands Ecology and Management 4: 65–72.

Turner, R. E. & Warren, R. S. (2003) Valuation of continuous and intermittent *Phragmites* control. *Estuaries* 26: 618–623.

Turner, R. K. (2000) Integrating natural and socio-economic science in coastal management. *Journal of Marine Systems* 25: 447–460.

Turner, R. K., Lorenzoni, I., Beaumont, N., et al. (1998) Coastal management for sustainable development: analysing environmental and socio-economic changes on the UK coast. *Geographical Journal* 164: 269–281.

Tweddle, D. (1992) Conservation and threats to the resources of Lake Malawi. *Mitteilungen Internationale Vereinigung für Limnologie* 24: 17–24.

Twenhöven, F. L. (1992a) Untersuchungen zur Wirkung stickstoffhaltiger Niederschläge auf die Vegetation von Hochmooren. Mitteilungen der Arbeitsgemeinschaft Geobotanik in Schleswig-Holstein und Hamburg 44: 1–172.

Twenhöven, F. L. (1992b) Competition between two Sphagnum species under different deposition levels. Journal of Bryology 17: 71–80.

Twilley, R. R., Chen, R. H. & Hargis, T. (1992) Carbon sinks in mangroves and their implications to carbon budget of tropical coastal ecosystems. *Water, Air, and Soil Pollution* 64: 265–288.

Twilley, R. R., Gottfried, R. R., Rivera-Monroy, V. H., et al. (1998a) An approach and preliminary model of integrating ecological and economic constraints of environmental quality in the Guayas River estuary, Ecuador. Environmental Science and Policy 1: 271–288.

Twilley, R. R., Rivera-Monroy, V. H., Chen, R. & Botero, L. (1998b) Adapting an ecological mangrove model to simulate trajectories in restoration ecology. *Marine Pollution Bulletin* 37: 404–419.

Tyler, P. A. (1988) Seasonality in the deep sea. Oceanography and Marine Biology Annual Review 26: 227–258.

Tyler, P. A. (2003a) Disposal in the deep sea: analogue of nature or *faux ami? Environmental Conservation* 30: 26–39.

Tyler, P. A. (2003b) The peripheral deep seas. In: Ecosystems of the World, vol. 28, Ecosystems of the Deep Ocean, ed. P. A. Tyler, pp. 261–293. Amsterdam, the Netherlands: Elsevier.

Tyler, P. A. & Young, C. M. (1999) Reproduction and dispersal at vents and cold seeps. *Journal of the Marine Biological Association of the United Kingdom* 79: 193–208.

Tyler, P. A., Grant, A., Pain, S. L. & Gage, J. D. (1982) Is annual reproduction in deep-sea echinoderms a response to variability in their environment? *Nature* 300: 747–750. Tynan, C. T. & DeMaster, D. (1997) Observations and predictions of Arctic climate change: potential effects on marine mammals. *Arctic* 50: 308–322.

Uchiyama, Y., Nadaoka, K., Rolke, P., Adachi, K. & Yagi, H. (2000) Submarine groundwater discharge into the sea and associated nutrient transport in a sandy beach. *Water Resources Research* 36: 1467–1479.

UKOOA (2002) UKOOA Drill Cuttings Initiative Final Report. United Kingdom Offshore Operators Association [www.document]. URL http://www.oilandgas.org.uk/ issues/drillcuttings/pdfs/finalreport.pdf

UN (1994) Report of the International Conference on Population and Development, UN Document No. A/CONF.171/13, New York, USA: UN Population Division.

UN (1997) World Urbanization Prospects: The 1996 Revision. New York, USA: UN Population Division.

UN (1998) Report of the conference of the parties on its third session, held at Kyoto from 1 to 11 December 1997.
United Nations Framework Convention on Climate Change [www document]. URL http://unfccc.int/resource/docs/cop3/07ao1.pdf

UN (1999) World Population Prospects: The 1998 Revision. New York, USA: UN Population Division.

UN (2002) United Nations human rights [www document]. URL http://www.un.org/rights/

UN (2003) World Population Prospects: The 2002 Revision. New York, USA: UN Population Division.

UNCED (1992) Chapter 17. Agenda 21: Earth Summit – The United Nations Programme of Action from Rio, United Nations Conference on Environment and Development. New York, USA: UN Department of Public Information.

UN Country Team (2004) Kazakhstan: achievements, issues, and prospects [www document]. URL http://www.undp.kz/library_of_publications/files/ 1871-31465.pdf

Underwood, A. J. (1993) Exploitation of species on the rocky coast of New South Wales (Australia) and options for its management. *Ocean and Coastal Management* 20: 41–62.

Underwood, A. J. (1999) History and recruitment in structure of intertidal assemblages on rocky shores: an introduction to problems for interpretation of natural change. In: *Aquatic Life Cycle Strategies Survival in a Variable Environment*, ed. M. Whitfield, J. Matthews & C. Reynolds, pp. 79–96. Plymouth, UK: Marine Biological Association.

Underwood, A. J. & Fairweather, P. G. (1989) Supply-side ecology and benthic marine assemblages. *Trends in Ecology* and Evolution 4: 16–20. Underwood, A. J., Denley, E. J. & Moran, M. J. (1983) Experimental analyses of the structure and dynamics of mid-shore rocky intertidal communities in New South Wales. *Oecologia* 56: 202–219.

UNDP (1995) *Human Development Report*. New York, USA: Oxford University Press.

UNEP (1999*a*) Strategic Action Plan for the South China Sea [www document]. URL http://www.unep.org/dewa/ giwa/areas/area54.asp

UNEP (1999b) Global Environment Outlook 2000: UNEP's Millennium Report on the Environment. London, UK: Earthscan Publications.

UNEP (2000a) Regional-Integrated Sustainable Management of Trans-Boundary Environmental Resources in Southwestern Djibouti and Northeastern Ethiopia, UNEP Division of GEF Coordination, Nairobi [www document]. URL http://www.iwlearn.net/docs/smter/smter10e.pdf

UNEP (2000*b*) Action Plan for the Protection of the Marine Environment and the Sustainable Development of the Coastal Areas of the Mediterranean [www document]. URL http://www.unep.org/regionalseas/Programmes/UN-EP_Administered_ Programmes/Mediterranean_Region/ default2.asp

UNEP (2000c) Regional Seas: A Survival Strategy for Our Oceans and Coasts. Geneva, Switzerland: United Nations Environment Programme.

UNEP (2001a) GLOBIO: Global methodology for mapping human impacts on the biosphere, UNEP/DEWA/TR No. 01-3, Nairobi, Kenya: United Nations Environment Programme.

UNEP (2001*b*) The Mesopotamian marshlands: demise of an ecosystem. Unpublished report. Nairobi, Kenya: United Nations Environment Programme.

UNEP (2002) GEO-3: Global Environment Outlook 3 – Past, Present and Future Perspectives. Nairobi, Kenya: United Nations Environment Programme.

UNEP (2003) *Water for People, Water for Life*, UN World Water Development Report. Paris, France: UNESCO.

UNEP-WCMC (2004) Biodiversity and Climate Change [www document]. URL http://www.unep-wcmc.org/ climate/impacts.htm

UNESCO (2000) Vision for the Aral Sea Basin. Paris, France: UNESCO.

UNESCO-IHE (2003) Partner in Capacity Building [www document]. URL: http://www.unesco-ihe.org/education/ intro.htm

UNEP (2004) Desk Study on the Environment in Liberia. Nairobi, Kenva: United Nations Environment Programme.

UNPD (2002) International Migration Report 2002. New York, USA: UN Population Division.

- UNPD (2006) World Population Prospects: The 2006 Revision. New York, USA: UN Population Division.
- UN World Water Action Programme (2003) Water for People, Water for Life. Paris, France: UNESCO.

Urban, F. E., Cole, J. E. & Overpeck, J. T. (2000) Influence of mean climate change on climate variability from a 155-year tropical Pacific coral record. *Nature* 407: 989–993.

US Army Corps of Engineers (1989) *Benefits Reevaluation Study: Brigantine Island, New Jersey.* Philadelphia, PA, USA: US Army Corps of Engineers.

US Army Corps of Engineers (2001) The New York District's Biological Monitoring Program for the Atlantic Coast of New Jersey, Asbury Park to Manasquan Section Beach Erosion Control Project. Vicksburg, MS, USA: Engineer Research and Development Center, Waterways Experiment Station.

US Army Corps of Engineers (2003) 1988 National List of Plants that Occur in Wetlands. Washington, DC, USA: US Army Corps of Engineers Available online at http://www. fws.gov/nwi/bha/list88.html

US Census Bureau (2003) International Data Base (IDB) [www.document]. URL http://www.census.gov/ipc/ www/idbnew.html

US EPA (1992) Framework for ecological risk assessment forum. Unpublished report EPA/630/R92/001. Washington, DC, USA: US Environmental Protection Agency.

US EPA (2000) Report to Congress Groundwater Chapters, Report No. EPA 816-R-00-013, EPA Office of Water, Washington, DC, USA: US Environmental Protection Agency.

US EPA (2001) *National Coastal Condition Report*, Report No. EPA-620/R-01/005. Washington, DC, USA: US Environmental Protection Agency.

US Fish and Wildlife Service (2003*a*) Wildlife Institute of India. Washington, DC, USA: US Fish and Wildlife Service, Office of International Affairs. Available online at http://international.fws.gov/laws/wii.html (accessed August 12, 2003)

US Fish and Wildlife Service (2003b) US/South Africa Binational Commission. Washington, DC, USA: US Fish and Wildlife Service, Office of International Affairs. Available online at http://international.fws.gov/laws/ ussabc.html

US Geological Survey (1996) National Water Summary of Wetland Resources, US Geological Survey Water-Supply Paper No. 2425. Washington, DC, USA: US Government Printing Office. Uthicke, S. & Benzie, J. A. H. (2001) Effect of beche-de-mer fishing on densities and size structure of *Holothuria nobilis* (Echinodermata: Holothuroidea) populations on the Great Barrier Reef. *Coral Reefs* 3: 271–276.

Vadas, R. L. & Steneck, R. S. (1988) Zonation of deep water benthic algae in the Gulf of Maine. *Journal of Phycology* 24: 338–346.

Valentine, J. P. & Johnson, C. R. (2003) Establishment of the introduced kelp Undaria pinnatifida in Tasmania depends on disturbance to native algal assemblages. Journal of Experimental Marine Biology and Ecology 295: 63–90.

Valette-Silver, N. J. (1993) The use of sediment cores to reconstruct historical trends in contamination of estuarine and coastal sediments. *Estuaries* 16: 577–608.

Valiela, I. (1995) Marine Ecological Processes, 2nd edn. New York, USA: Springer-Verlag.

Valiela, I., Cole, M. L., McClelland, J., et al. (2000) Role of salt marshes as part of coastal landscapes. In: Concepts and Controversies in Tidal Marsh Ecology, ed. M. P. Weinstein & D. P. Kreeger, pp. 23–38. Dordrecht, the Netherlands: Kluwer.

Valiela, I., Foreman, K., LaMontagne, M., et al. (1992) Couplings of watersheds and coastal waters: sources and consequences of nutrient enrichment in Waquoit Bay, Massachusetts. *Estuaries* 15: 443–457.

Valiela, I., Kremer, J., Lajtha, K., et al. (1997) Nitrogen loading from coastal watersheds to receiving estuaries: new method and application. *Ecological Applications* 7: 358–380.

Vallentyne, J. & Beeton, A. (1988) The ecosystem approach to managing human uses and abuses of natural resources in the Great Lakes Basin. *Environmental Conservation* 15: 59–62.

Valtýsson, H. (2001) The sea around Icelanders: catch history and discards in Icelandic waters. In: *Fisheries Impacts on North Atlantic Ecosystems: Catch, Effort and National/ Regional Data Sets*, Fisheries Centre Research Report No. 9 (3), ed. D. Zeller, R. Watson & D. Pauly, pp. 52–87.
Vancouver, BC, Canada: University of British Columbia.

Valtýsson, H. & Pauly, D. (2003) Fishing down the food web: an Icelandic case study. In: Competitiveness within the Global Fisheries, Proceedings of a Conference, Akureyri, Iceland, 6–7 April 2000, ed. E. Guðmundsson & H. Valtýsson. Akureyri, Iceland: University of Akureyri.

Van Boxel, J. H., Jungerius, P. D., Kieffer, N. & Hampele, N. (1997) Ecological effects of reactivation of artificially stabilized blowouts in coastal dunes. *Journal of Coastal Conservation* 3: 57–62. Van Buynder, P., Sam, G., Russell, R., *et al.* (1995) Barmah Forest Virus epidemic on the south coast of New South Wales. *Communicable Diseases Intelligence* **19**: 188–191.

Van Dam, R., Gitay, H., Finlayson, C. M., Davidson, N. & Orlando, B. (2002) Climate Change and Wetlands: Impacts, Adaptation and Mitigation. Information Paper. Ramsar COP8, Doc.11 + annex [www document]. URL http:// www.ramsar.org/cop8/cop8_doc_11_e.htm.

Van Dam, R. A., Finlayson, C. M. & Watkins, D., eds. (1999) Vulnerability Assessment of Major Wetlands in the Asia-Pacific Region to Climate Change and Sea Level Rise, Supervising Scientist Report No. 149, Canberra, ACT, Australia: Supervising Scientist.

Vandelannoote, A., Robberecht, H., Deelstra, H., Vyumvohore, F., Bitetera, L. & Ollevier, F. (1996) The impact of the River Ntahangwa, the most polluted Burundian affluent of Lake Tanganyika, on the water quality of the lake. *Hydrobiologia* 328: 161–171.

Van den Broek, W. L. F. & Tracey, D. M. (1981) Concentration and distribution of mercury in flesh of orange roughy (*Hoplostethus atlanticus*). New Zealand Journal of Marine and Freshwater Research 15: 255–260.

Van der Nat, D., Schmidt, A., Edwards, P. J., Tockner, K. & Ward, J. V. (2002) Inundation dynamics in braided floodplains (Fiume Tagliamento). *Ecosystems* 5: 636–647.

Vanderploeg, H., Johengen, T., Strickler, J., Liehig, J. & Nalepa, T. (2001) Zebra mussels (*Dreissena polymorpha*) selective filtration promoted *Microcystis* blooms in Saginaw Bay (Lake Huron) and Lake Erie. *Canadian Journal of Fisheries and Aquatic Sciences* 58: 1208–1221.

Vanderploeg, H., Nalepa, T., Jude, D., et al. (2002) Dispersal and emerging ecological impacts of Ponto-Caspian species in the Laurentian Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 59: 1209–1228.

Van der Wal, D. (1998) The impact of the grain-size distribution of nourishment sand on aeolian sand transport. *Journal of Coastal Research* 14: 620–631.

Van der Wal, D. (2000) Grain-size-selective aeolian sand transport on a nourished beach. *Journal of Coastal Research* 16: 896–908.

Van Dijk, G. M., Van Liere, L., Bannik, B. A. & Cappon, J. J. (1994) Present state of the water quality of European rivers and implications for management. *Science of the Total Environment* 145: 187–195.

Van Dover, C. L. (2000) The Ecology of Deep-Sea Hydrothermal Vents. Princeton, NJ, USA: Princeton University Press.

Van Dover, C. L., German, C. R., Speer, K. G., Parson, L. M. & Vrijenhoek, R. C. (2002) Evolution and biogeography of deep-sea vent and seep invertebrates. *Science* 295: 1253–1257.

Van Dover, C. L., Grassle, J. F., Fry, B., Garritt, R. H. & Staczak, V. R. (1992) Stable isotopic evidence for entry of sewage-derived organic material into a deep-sea food web. *Nature* 360: 153–156.

Van Duzer, C. (2004) Floating Islands: A Global Bibliography, with an Edition and Translation of G. C. Munz' Exercitatio academico de insulis natantibus (1711). Los Altos Hills, CA, USA: Cantor Press.

- Van Lent, F., Nienhuis, P. H. & Verschuure, J. M. (1991) Production and biomass of the seagrass Zostera noltii Hornem. and Cymodocea nodosa (Ucria) Ascherson at the Banc d'Arguin (Mauritania, NW Africa): a preliminary approach. Aquatic Botany 41: 353–367.
- Van Noordwiijk, J. M. & Peerbolte, E. B. (2000) Optimal sand nourishment decisions. *Journal of Waterway, Port, Coastal* and Ocean Engineering 126: 30–38.
- Van Soosten, C., Schmidt, H. & Westheide, W. (1998) Genetic variability and relationships among geographically widely separated populations of *Petitia amphophthalma* (Polychaeta: Syllidae): results from RAPD-PCR investigations. *Marine Biology* 131: 659–669.
- Van Wilgen, B. W., Richardson, D. M., Le Maitre, D. C., Marais, C. & Magadlela, D. (2001) The economic consequences of alien plant invasions: examples of impacts and approaches to sustainable management in South Africa. *Environment, Development and Sustainability* 3: 145–168.
- Vasquez, J. A. (1993) Abundance, distributional patterns and diets of main herbivorous and carnivorous species associated with *Lessonia trabeculata* kelp beds in northern Chile. Serie Ocasional de Universidad Católica del Norte 2: 213–229.
- Vasquez, J. A. & Buschmann, A. H. (1997) Herbivore–kelp interactions in Chilean subtidal communities: a review. *Revista Chilena de Historia Natural* 70: 41–52.

Vasquez, J. A., Castilla, J. C. & Santelices, B. (1984) Distributional patterns and diets of four species of sea urchins in giant kelp forest (*Macrocystis pyrifera*) of Puerto Toro, Navarino Island, Chile. *Marine Ecology Progress Series* 19: 55–63.

- Vaughan, D. G., Marshall, G. J., Connolley, W. M., King, J. C. & Mulvaney, R. (2001) Devil in the detail. *Science* 293: 1777–1779.
- Velimirov, B., Field, F. G., Griffiths, C. L. & Zoutendyk, P. (1977) The ecology of kelp bed communities in the Benguela upwelling system. *Helgoländer Wissenschaftliche Meeresuntersuchungen* 30: 495–518.

Venema, H. D., Schiller, E. J., Adamowski, K. & Thizy, J.-M. (1997) A water resources planning response to climate change in the Senegal River basin. *Journal of Environmental Management* 49: 125–155.

Verberg, P., Hecky, R. & Kling, H. (2003) Ecological consequences of a century of warming in Lake Tanganyika. *Science* 301: 505–507.

Verhoeven, J. T. A. & Bobbink, R. (2001) Plant diversity of fen landscapes in the Netherlands. In: *Biodiversity in Wetlands: Assessment, Function and Conservation*, ed. B. Gopal, pp. 65–87. Leiden, the Netherlands: Backhuys.

Verissimo, J. (1895) *A Pesca na Amazonia*. Rio de Janeiro, Brazil: Livraria Alves.

Verity, P. G. & Smetacek, V. (1996) Organism life cycles, predation, and the structure of marine pelagic ecosystems. *Marine Ecology Progress Series* 130: 277–293.

Verity, P. G., Smetacek, V. & Smayda, T. J. (2002) Status, trends, and the future of the marine pelagic ecosystem. *Environmental Conservation* 29: 207–237.

Vermaat, J. E., Beijer, J. A., Gijlstra, J., et al. (1993) Leaf dynamics and standing stocks of intertidal Zostera noltii Hornem. and Cymodocea nodosa (Ucria) Ascherson on the Banc d'Arguin (Mauritania). Hydrobiologia 258: 59–72.

Vermeer, J. G. & Joosten, J. H. J. (1992) Conservation and management of bog and fen reserves in the Netherlands. In: *Fens and Bogs in the Netherlands: Vegetation, History, Nutrient Dynamics and Conservation*, ed. J. T. A. Verhoeven, pp. 433–478. Dordrecht, the Netherlands: Kluwer.

Vermeij, G. J. (1986) Survival during biotic crises: the properties and evolutionary significance of refuges. In: *Dynamics* of *Extinction*, ed. D. K. Elliott, pp. 231–246. New York, USA: John Wiley.

Veron, J. E. N., ed. (1995) Corals in Space and Time: The Biogeography and Evolution of the Scleractinia. Sydney, NSW, Australia: University of New South Wales Press.

Veron, J. E. N. & Minchin, P. R. (1992) Correlations between sea surface temperature, circulation patterns and the distribution of hermatypic corals of Japan. *Continental Shelf Research* 12: 835–857.

Vetter, E. W. (1995) Detritus-based patches of high secondary production in the nearshore benthos. *Marine Ecology Progress Series* 120: 251–262.

Vetter, E. W. & Dayton, P. K. (1998) Macrofaunal communities within and adjacent to a detritus-rich submarine canyon system. *Deep-Sea Research II* 45: 25–54.

Vezina, A. R. & Platt, T. (1988) Food web dynamics in the ocean. I. Best estimates of flow networks using inverse methods. *Marine Ecology Progress Series* 42: 269–287. Viles, H. & Spencer, T. (1995) Coastal Problems: Geomorphology, Ecology and Society of the Coast. London, UK: Edward Arnold.

Vinnikov, K. Y., Robock, A., Stouffer, R. J., et al. (1999) Global warming and northern hemisphere sea ice extent. *Science* 286: 1934–1937.

Vinueza, L., Branch, G. M., Branch, M. L. & Bustamante, R. H. (2006) Top-down herbivory and bottom-up El Niño effects on Galapagos rocky-shore communities. *Ecological Monographs* 76: 111–131.

Vitousek, P. M., Aber, J. D., Howarth, R. W., et al. (1997a) Human alteration of the global nitrogen cycle: sources and consequences. *Ecological Applications* 7: 737–750.

Vitousek, P. M., D'Antonio, C. M., Loope, L. L. & Westbrooks, R. (1996) Biological invasions as global environmental change. *American Scientist* 84: 468–478.

Vitousek, P. M., Mooney, H. A., Lubchenco, J. & Melillo, J. M. (1997b) Human domination of earth's ecosystems. *Science* 277: 494–499.

Vitt, D. H. & Chee, W. (1990) The relationships of vegetation to surface water chemistry and peat chemistry in fens of Alberta, Canada. *Vegetatio* 89: 87–106.

Vitt, D. H., Halsey, L. A., Bauer, I. E. & Campbell, C. (2000) Spatial and temporal trends in carbon storage of peatlands of continental western Canada through the Holocene. *Canadian Journal of Earth Sciences* 37: 683–693.

Vitt, D. H., Halsey, L. A., Wieder, K. & Turetsky, M. (2003) Response of *Sphagnum fuscum* to nitrogen deposition: a case study of ombrogenous peatlands in Alberta, Canada. *Bryologist* 106: 235–245.

Vitt, D. H., Yenhung, L. & Belland, R. J. (1995) Patterns of bryophyte diversity in peatlands of continental western Canada. *Bryologist* 98: 218–227.

Vollmer, M., Weiss, R. & Bootsma, H. (2002) Ventilation of Lake Malawi/Nyasa. In: *The East African Great Lakes: Limnology, Palaeolimnology, and Biodiversity*, ed. E. Odada & D. Olago, pp. 209–234. Dordrecht, the Netherlands: Kluwer.

Volonte, C. R. & Nicholls, R. J. (1995) Uruguay and sea-level rise: potential impacts and responses. *Journal of Coastal Research* Special Issue 14: 262–285.

Vörösmarty, C. J., Askew, A., Barry, R., *et al.* (2001) Global water data: a newly endangered species. *Eos* 82(5): 54, 56, 58.

Vörösmarty, C. J., Green, P., Salisbury, J. & Lammers, R. B. (2000) Global water resources: vulnerability from climate change and population growth. *Science* 289: 284–288. Wadhams, P. & Davis, N. R. (2000) Further evidence of ice thinning in the Arctic Ocean. *Geophysical Research Letters* 27: 3973–3975.

Wagner, C. S., Brahmakulam, I., Jackson, B., Wong, A. & Yoda, T. (2001) Science and Technology Collaboration: Building Capacity in Developing Countries?, Report prepared for the World Bank. Santa Monica, CA, USA: RAND Science & Technology. Available online at: www.rand.org/ pubs/monograph_reports/MR1357.0/MR1357.0.pdf

Wagner, T. (1994) Remedial Action Plans: looking back and moving forward. *Focus on International Joint Commission Activities* 19: 16–17.

Wagner, W., Gawel, J., Furumai, H., et al. (2002) Sustainable watershed management: an international multi-watershed case study. *Ambio* 31: 2–13.

Wake, D. B. (1991) Declining amphibian populations. *Science* 253: 860.

Walden, D., van Dam, R., Finlayson, M., et al. (2004) A risk assessment of the tropical wetland weed Mimosa pigra in northern Australia. In: Research and Management of Mimosa pigra, ed. M. Julien, G. Flanagan, T. Heard, et al., pp. 11–21. Canberra, ACT, Australia: CSIRO.

Walker, D. I. (1989) Regional studies: seagrass in Shark Bay, the foundations of an ecosystem. In: *Biology of Seagrasses: A Treatise on the Biology of Seagrasses with Special Reference to the Australian Region*, ed. A. W. D. Larkum, A. J. McComb & S. A. Shepherd, pp. 182–210. Amsterdam, the Netherlands: Elsevier.

Walker, D. I. (1991) The effect of sea temperature on seagrasses and algae on the Western Australian coastline. *Journal of the Royal Society of Western Australia* 74: 71–77.

Walker, D. I. (2003) The seagrasses of Western Australia. In: World Atlas of Seagrasses, ed. E. P. Green & F. T. Short, pp. 121–130. Berkeley, CA, USA: University of California Press.

Walker, D. I. & McComb, A. J. (1992) Seagrass degradation in Australian coastal waters. *Marine Pollution Bulletin* 25: 191–195.

Walker, D. I. & Woelkerling, W. J. (1988) A quantitative study of sediment contribution by epiphytic coralline red algae in seagrass meadows in Shark Bay, Western Australia. *Marine Ecology Progress Series* 43: 71–77.

Walker, D. I., Hillman, K. A., Kendrick, G. A. & Lavery, P. (2001) Ecological significance of seagrasses: assessment for management of environmental impact in Western Australia. *Ecological Engineering* 16: 323–330.

Walker, D. I., Kendrick, G. A. & McComb, A. J. (1988) The distribution of seagrasses in Shark Bay, Western Australia, with notes on their ecology. *Aquatic Botany* 30: 305–317. Walker, D. I., Lukatelich, R. J., Bastyan, G. & McComb, A. J. (1989) Effect of boat moorings on seagrass beds around Perth, Western Australia. *Aquatic Botany* 36: 69–77.

Walker, J. & Peet, R. K. (1984) Composition and species-diversity of pine-wiregrass savannas of the Green Swamp, North Carolina. *Vegetatio* 55: 163–179.

Walker, K. F. & Thoms, M. C. (1993) Environmental effects of flow regulation on the lower River Murray, Australia. *Regulated Rivers: Research and Management* 8: 103–119.

Wallace, J. B. & Webster, J. R. (1996) The role of macroinvertebrates in stream ecosystem function. *Annual Review of Entomology* 41: 115–139.

Wallén, B. (1986) Above and below ground dry mass of the three main vascular plants on hummocks on a sub-arctic peat bog. *Oikos* 46: 51–56.

Wallén, B. (1992) Methods for studying below-ground production in mire ecosystems. Suo 43: 155–162.

Walters, B. B. (2000) Local mangrove planting in the Philippines: are fisherfolk and fishpond owners effective restorationists? *Restoration Ecology* 8: 237–246.

Walters, M. J. (1992) A Shadow and a Song: The Struggle to Save an Endangered Species. Post Mills, VT, USA: Chelsea Green Publishing.

Walther, G.-R., Post, E., Convey, P., et al. (2002) Ecological responses to recent climate change. Nature 416: 391–395.

Wang, C., Prinn, R. & Sokolov, A. (1998) A global interactive chemistry and climate model: formulation and testing. *Journal of Geophysical Research* 103: 3399–3417.

Wang, X. & Key, J. R. (2003) Recent trends in Arctic surface, cloud, and radiation properties from space. *Science* 299: 1725–1728.

Wania, F., Axelman, J. & Broman, D. (1998) A review of processes involved in the exchange of persistent organic pollutants across the air-sea interface. *Environmental Pollution* **102**: 3–23.

Wantzen, K. M. (1998) Siltation effects on benthic communities in first order streams in Mato Grosso, Brazil. *Verhandlungen Internationale Vereinigung für Limnologie* 26: 1155–1159.

Ward, J. V., Tockner, K. & Schiemer, F. (1999) Biodiversity of floodplain river ecosystems: ecotones and connectivity. *Regulated Rivers: Research and Management* 15: 125–139.

Ward, J. V., Tockner, K., Arscott, D. B. & Claret, C. (2002) Riverine landscape diversity. *Freshwater Biology* 47: 517–539.

Wardle, D. (2002) Communities and Ecosystems: Linking the Aboveground and Belowground Components. Princeton, NJ, USA: Princeton University Press. Wares, J. P., Goldwater, D. S., Kong, B. Y. & Cunningham, C. W. (2002) Refuting a controversial case of a human-mediated marine species introduction. *Ecology Letters* 5: 577–584.

Warne, A. G., Toth, L. A. & White, W. A. (2000) Drainagebasin-scale geomorphic analysis to determine reference conditions for ecologic restoration: Kissimmee River, Florida. *Geographical Society of America Bulletin* 112: 884–899.

Warren, R. S. & Niering, W. A. (1993) Vegetation change on a north east tidal marsh: interaction of sea-level rise and marsh accretion. *Ecology* 74: 96–103.

Warrick, R. A., Le Provost, C., Meier, M. F., Oerlemans, J. & Woodworth, P. L. (1996) Changes in sea level. In: *Climate Change 1996: The Science of Climate Change*, ed.

J. T. Houghton, L. G. Meira Filho, B. A. Callander, *et al.*, pp. 359–405. Cambridge, UK: Cambridge University Press.

Wassmann, P. (1998) Retention versus export food chains: processes controlling sinking loss from marine pelagic systems. *Hydrobiologia* 63: 29–57.

Watkins, A. B. & Simmonds, J. (2000) Current trends in Antarctic sea ice: the 1990s impact on a short climatology. *Journal of Climate* 13: 4441–4451.

Watling, L. & Norse, E. A. (1998) Disturbance of the seabed by mobile fishing gear: a comparison to forest clearcutting. *Conservation Biology* 12: 1180–1197.

Watson, R. & Pauly, D. (2001) Systematic distortions in world fisheries catch trends. *Nature* 414: 534–536.

Watson, J. G. (1928) *Mangrove Forests of the Malay Peninsula*, Malayan Forest Records No. 6. Singapore, Malaysia: Fraser & Neave.

WCD (2000) Dams and Development: A New Framework for Decision-Making. London, UK: Earthscan Publications for World Commission on Dams.

 WCMC (1998) Freshwater Biodiversity: a preliminary global assessment. Biodiversity Series No. 8. By B. Groombridge & M. Jenkin. World Conservation Monitoring Centre, Cambridge, UK: World Conservation Press.

Weber, P. (1994) It comes down to the coasts. World Watch-March/April: 20–29.

Webster, M. D., Babiker, M., Mayer, M., et al. (2002) Uncertainty in emissions projections for climate models. *Atmospheric Environment* 36: 3659–3670.

Webster, M. D., Forest, C., Reilly, J., et al. (2003) Uncertainty analysis of climate change and policy response. *Climatic Change* 61: 295–320.

Wehrmeyer, W. & Mulugetta, Y., eds. (1999) Growing Pains: Environmental Management in Developing Countries.Sheffield, UK: Greenleaf Publishing. Weier, J. (2000) Bright lights, big city [www document]. URL http://earthobservatory.nasa.gov/Study/Lights/

Weinstein, M. & Kreeger, D. A., eds. (2000) Concepts and Controversies in Tidal Marsh Ecology. Dordrecht, the Netherlands: Kluwer.

Weis, P., Weis, J. S. & Proctor, T. (1993) Copper, chromium, and arsenic in estuarine sediments adjacent to wood treated with chromated-copper-arsenate (CCA). *Estuarine, Coastal* and Shelf Science 36: 71–79.

Welch, K. A., Lyons, W. B., McKnight, D. M., et al. (2003) Climate and hydrologic variations and implications for lake and stream ecological response in the McMurdo Dry Valleys, Antarctica. In: *Climate Variability and Ecological Response*, ed. D. Greenland, D. Goodin & R. C. Smith, pp. 174–195. Oxford, UK: Oxford University Press.

Welcomme, R. L. (1975) The Fisheries Ecology of African Floodplains, CIFA Technical Paper No. 3. Rome, Italy: Food and Agriculture Organization of the United Nations.

Welcomme, R. L. (1979) *Fisheries Ecology of Floodplain Rivers*. Harlow, UK: Longman.

Welcomme, R. L., ed. (1985) *River Fisheries*, Fisheries Technical Paper No. 262. Rome, Italy: Food and Agriculture Organization of the United Nations.

Welcomme, R. L. (2006) Providing for the water requirements of river fish and fisheries. *International Journal of Ecology* and Environmental Sciences 32: 85–97.

Wellington, G. M. & Glynn, P. W. (2007) Coral reef responses to El Niño–Southern Oscillation sea warming events. In: Geological Approaches to Coral Reef Ecology: Placing the Current Crisis in the Historical Context, ed. R. Aronson, pp. 342–385. New York, USA: Springer-Verlag.

Wells, L. & McLain, A. (1973) Lake Michigan: Man's Effects on Native Fish Stocks and Other Biota, Technical Report No. 20. Ann Arbor, MI, USA: Great Lakes Fishery Commission.

Wenger, E. L., Zinke, A. & Gutzweiler, K.-A. (1990) Present situation of European floodplain forests. *Forest Ecology and Management* 33/34: 5–12.

West, J. M. & Salm, R. V. (2003) Resistance and resilience to coral bleaching: implications for coral reef conservation and management. *Conservation Biology* 17: 956–967.

West, J. M. & Zedler, J. B. (2000) Marsh–creek connectivity: fish use of a tidal salt marsh in southern California. *Estuaries* 23: 699–710.

Western Australia, Government of (2002) Policy for the Implementation of Ecologically Sustainable Development for Fisheries and Aquaculture within Western Australia, Report No. 157. Perth, WA, Australia: Department of Fisheries. Weston, D. P. (1990) Quantitative examination of macrobenthic community changes along an organic and organic-enrichment gradient. *Marine Ecology Progress Series* 61: 233–247.

Wetlands International (2002) A Directory of Wetlands of International Importance, 7th edn [www document]. URL http://www.wetlands.org/RDB/Directory.html

Wheatcroft, R. A. (2000) Oceanic flood sedimentation: a new perspective. *Continental Shelf Research* 20: 2059–2066.

Wheatcroft, R. A., Borgeld, J. C., Born, R. S., et al. (1996) The anatomy of an oceanic flood deposit. Oceanography 9: 158–162.

Wheeler, B. D. (1984) British fens: a review. In: European Mires, ed. P. Moore, pp. 237–282. New York, USA: Academic Press.

Wheeler, B. D. & Proctor, M. C. F. (2000) Ecological gradients, subdivisions and terminology of north-west European mires. *Journal of Ecology* 88: 187–203.

Wheeler, B. D., Shaw, S. C., Fojt, W. J. & Robertson, R. A. (1995) *Restoration of Temperate Wetlands*. New York, USA: John Wiley.

White, W. B. & Peterson, R. G. (1996) An Antarctic circumpolar wave in surface pressure, wind, temperature and sea-ice extent. *Nature* **380**: 699–702.

White, W. B., Chen, S. C. & Peterson, R. G. (1998) The Antarctic circumpolar wave: a beta effect in ocean– atmosphere coupling over the Southern Ocean. *Journal of Physical Oceanography* 28: 2345–2361.

Whitehead, P. J. & Blomquist, G. O. (1991) Measuring contingent values for wetlands: effects of information about related environmental goods. *Water Resources Research* 27: 2523–2531.

Whitfield, P. E., Kenworthy, W. J., Hammerstrom, K. K. & Fonseca, M. S. (2002) The role of a hurricane in the expansion of disturbances initiated by motor vessels on seagrass banks. *Journal of Coastal Research* Special Issue 37: 85–98.

Whitlock, J. E., Jones, D. T. & Harwood, V. J. (2002) Identification of the sources of fecal coliforms in an urban watershed using antibiotic resistance analysis. *Water Research* 36: 4273–4282.

Whittaker, R. H. (1975) *Communities and Ecosystems*, 2nd edn. New York, USA: Macmillan.

WHO (1996) Water and Sanitation, Fact Sheet No. 112, World Health Organization [www document]. URL http:// www.who.int/inffs/en/fact112html

Wickens, P. A. & Field, J. G. (1986) The effects of water transport on nitrogen flow through a kelp-bed community. *South African Journal of Marine Sciences* 4: 79–92. Wiegers, J. (1990) Forested wetlands in western Europe. In: *Ecosystems of the World*, vol. 15, *Forested Wetlands*, ed.
A. E. Lugo, M. M. Brinson & S. Brown, pp. 407–436.
Amsterdam, the Netherlands: Elsevier.

Wiggins, B. A., Andrews, R. W., Conway, R. A., et al. (1999) Use of antibiotic resistance analysis to identify nonpoint sources of fecal pollution. *Applied and Environmental Microbiology* 65: 3483–3486.

Wigham, B. D., Tyler, P. A. & Billett, D. S. M. (2003) Reproductive biology of the abyssal holothurian *Amperima rosea*: an opportunistic response to variable flux of surface derived organic matter? *Journal of the Marine Biological Association of the United Kingdom* 83: 175–188.

Wigley, T. M. L. (1989) Measurement and prediction of global warming. In: Ozone Depletion: Health and Environmental Consequences, ed. R. R. Jones & T. Wigley, pp. 85–97. Chichester, UK: John Wiley.

Wigley, T. M. L. (1995) Global-mean temperature and sea level consequences of greenhouse gas stabilization. *Geophysical Research Letters* 22: 45–48.

Wilcock, H. R., Hildrew, A. G. & Nichols, R. A. (2001) Genetic differentiation of a European caddisfly: past and present gene flow among fragmented larval habitats. *Molecular Ecology* **10**: 1821–1834.

Wilcock, H. R., Nichols, R. A. & Hildrew, A. G. (2003) Genetic population structure and neighbourhood population size estimates of the caddisfly *Plectrocnemia conspersa*. *Freshwater Biology* **48**: 1813–1824.

Wilcove, D. S., Rothstein, D., Dubow, J., Phillips, A. & Losos, E. (1998) Quantifying threats to imperiled species in the United States. *BioScience* 48: 607–615.

Wild, S. R. & Jones, K. C. (1995) Polynuclear aromatic hydrocarbons in the United Kingdom environment: a preliminary source inventory and budget. *Environmental Pollution* 88: 91–108.

Wildlife Institute of India (2003) Homepage [www document]. URL http://www.wii.gov.in/

Wilkens, H., Culver, D. C. & Humphreys, W. F., eds. (2000) *Ecosystems of the World*, vol. 30, *Subterranean Ecosystems*. Amsterdam, the Netherlands: Elsevier.

Wilkie, M. J. & Fortuna, S. (2003) Status and Trends in Mangrove Area Extent Worldwide, Forest Resources Assessment Working Paper No. 63. Rome, Italy: Food and Agriculture Organization of the United Nations.

Wilkinson, C. (1996) Global change and coral reefs: impacts on reefs, economies and human cultures. *Global Change Biology* 2: 547–558. Wilkinson, C. R. & Sammarco, P. W. (1983) Effects of fish grazing and damselfish territoriality on coral reef algae. II. Nitrogen fixation. *Marine Ecology Progress Series* 13: 15–19.

William, P. B. & Orr, M. K. (2002) Physical evolution of restored breached levee salt marshes in the San Francisco Bay estuary. *Restoration Ecology* 10: 527–542.

Williams, D. M. (2001) Impacts of terrestrial run-off on the Great Barrier Reef World Heritage Area. Unpublished report. Townsville, Queensland, Australia: CRC Reef Research Centre.

Williams, G. D. & Zedler, J. B. (1999) Fish assemblage composition in constructed and natural tidal marshes of San Diego Bay: relative influence of channel morphology and restoration history. *Estuaries* 22: 702–716.

Williams, I. D. & Polunin, N. V. C. (2001) Large-scale associations between macroalgal cover and grazer biomass on mid-depth reefs in the Caribbean. *Coral Reefs* 19: 358–366.

Williams, I. D., Polunin, V. C. & Hendrick, V. J. (2001) Limits to grazing by herbivorous fishes and the impact of low coral cover on macroalgal abundance on a coral reef in Belize. *Marine Ecology Progress Series* 222: 187–196.

Williams, M., ed. (1990) Wetlands: A Threatened Landscape. Oxford, UK: Blackwell Scientific Publications.

Williams, W. D. (1981) Problems in the management of inland saline lakes. Verhandlungen Internationale Vereinigung für Limnologie 21: 688–692.

Williams, W. D. (1993a) Conservation of salt lakes. *Hydrobiologia* 267: 291–306.

Williams, W. D. (1993b) The worldwide occurrence and limnological significance of falling water-levels in large, permanent saline lakes. *Verhandlungen Internationale Vereinigung für Limnologie* 25: 980–983.

Williams, W. D. (1995) Lake Corangamite, Australia, a permanent saline lake: conservation and management issues. *Lakes and Reservoirs: Research and Management* 1: 55–64.

Williams, W. D. (1996) The largest, highest and lowest lakes of the world: saline lakes. *Verhandlungen Internationale Vereinigung für Limnologie* 26: 61–79.

Williams, W. D. (1998a) Management of Inland Saline Waters, United Nations Environment Programme/International Lake Environment Committee Guidelines of Lake Management Series No. 6. Kusatsu, Japan: ILEC/UNEP.

Williams, W. D. (1998b) Salinity as a determinant of the structure of biological communities in salt lakes. *Hydrobiologia* 381: 191–201.

Williams, W. D. (1998c) Dryland wetlands. In: Wetlands for the Future, ed. A. J. McComb & J. A. Davis, pp. 33–47. Adelaide, SA, Australia: Gleneagles Publishing. Williams, W. D. (1999) Salinization: a major threat to water resources in the arid and semi-arid regions of the world. *Lakes and Reservoirs: Research and Management* 4: 85–91.

Williams, W. D. (2000) Biodiversity in temporary wetlands of dryland regions. Verhandlungen Internationale Verieinigung für Limnologie 27: 141–144.

Williams, W. D. (2001) Salinization: unplumbed salt in a parched landscape. *Water Science and Technology* 43(4): 85–91.

Williams, W. D. (2002a) Environmental threats to salt lakes and the likely status of inland saline ecosystems in 2025. *Environmental Conservation* 29: 154–167.

Williams, W. D. (2002b) Anthropogenic salinization of inland waters. *Hydrobiologia* 466: 329–337.

Williams, W. D. & Aladin, N. V. (1991) The Aral Sea: recent limnological changes and their conservation significance. *Aquatic Conservation: Marine and Freshwater Ecosystems* 1: 3–23.

Williamson, M. (1999) Invasions. Ecography 22: 5-12.

Williamson, M. & Fitter, A. (1996) The varying success of invaders. *Ecology* 77: 1661–1666.

Willott, E. (2004) Restoring Nature, without mosquitoes? *Restoration Ecology* 12: 147–153.

Willrich, T. L. & Smith, G. E., ed. (1970) Agricultural Practices and Water Quality. Ames, IA, USA: Iowa State University Press.

Wilsey, B. J. & Potvin, C. (2000) Biodiversity and ecosystem functioning: the importance of species evenness in an old field. *Ecology* 81: 887–892.

Wilson, R. R. & Kaufmann, R. S. (1987) Seamount biota and biogeography. In: *Seamounts, Islands and Atolls*, ed.
B. H. Keating, P. Fryer, R. Batiza & G. W. Boehlert, pp. 355–

377. Washington, DC, USA: American Geophysical Union.

Wilson, S. K., Graham, N. A. J., Pratchett, M. S., Jones, G. P. & Polunin, N. V. C. (2006) Multiple disturbances and the global degradation of coral reefs: are reef fishes at risk or resilient? *Global Change Biology* 12: 1–15.

Wilson, W. D., Fatemi, S. M. R., Shokri, M. R. & Claereboudt, M. R. (2002) Status of coral reefs of Persian/ Arabian Gulf and Arabian Sea Region. In: *Status of Coral Reefs of the World: 2002*, ed. C. Wilkinson, pp. 53–62. Townsville, Queensland, Australia: Australian Institute of Marine Science.

Wiltshire, J. (2001) Future prospects for the marine minerals industry. Underwater 13: 40–44.

Windom, H. L. (1992) Contamination of the marine environment from land-based sources. *Marine Pollution Bulletin* 25: 32–36. Winger, P. V. (2002) Toxicological assessment of aquatic ecosystems: application to watercraft contaminants in shallow water environments. *Journal of Coastal Research* Special Issue 37: 178–190.

Winn, P. J. S., Young, R. M. & Edwards, A. M. C. (2003) Planning for the rising tides: the Humber Estuary Shoreline Management Plan. *Science of the Total Environment* 314–316: 13–30.

Winsor, P. (2001) Arctic sea ice thickness remained constant during the 1990s. *Geophysical Research Letters* 28: 1039–1041.

Winter, T. C. (1988) A conceptual framework for assessing cumulative impacts on hydrology of nontidal wetlands. *Environmental Management* 12: 605–620.

Winter, T. C. (1992) A physiographic and climatic framework for hydrologic studies of wetlands. In: Aquatic Ecosystems in Semi-Arid Regions: Implications for Resource Management, ed. R. D. Robarts & M. L. Bothwell, pp. 127–148. Saskatoon, Alberta, Canada: Environment Canada.

Winter, T. C. (1999) Relation of streams, lakes, and wetlands to groundwater flow systems. *Hydrogeology Journal* 7: 28–45.

Winter, T. C. (2001) The concept of hydrological landscapes. Journal of the American Water Resources Association 37: 335–349.

Wiseman, R., Taylor, D. & Zingstra, H., ed. (2003) Wetlands and agriculture, Proceedings of the Workshop on Agriculture, Wetlands and Water Resources, 17th Global Biodiversity Forum, Valencia, Spain, November 2002. International Journal of Ecology and Environmental Sciences Special Issue 29: 1–122.

Wishner, K., Levin, L., Gowing, M. & Mullineaux, L. (1990) Involvement of the oxygen minimum in benthic zonation on a deep seamount. *Nature* 346: 57–59.

Wittmann, F. (2002) Species distribution, community structure, and adaptations of Amazonian Várzea forests depending on annual flood-stress, using remote sensing methods. Unpublished Ph.D. thesis, University of Mannheim, Germany.

Woinarski, J. C. Z., Tidemann, S. C. & Kerin, S. (1988) Birds in a tropical mosaic: the distribution of bird species in relation to vegetation patterns. *Australian Wildlife Research* 15: 171–196.

Wolanski, E. (1989) Measurements and modelling of the water circulation in mangrove swamps. Unpublished report, RAF/87/038, Série Documentaire No. 3, pp. 1–43, COMARF Regional Project for Research and Training in Coastal Marine Systems in Africa. Paris, France: UNESCO and Coastal Marine Project in Africa.

Wolanski, E., Mazda, Y. & Ridd, P. (1992) Mangrove hydrodynamics. In: *Tropical Mangrove Ecosystems*, ed.
A. I. Robertson & D. M. Alongi, pp. 43–62. Washington, DC, USA: American Geophysical Union.

Wolff, W. J. (1997) Case studies and reviews: development of the conservation of Dutch coastal waters. *Aquatic Conservation: Marine and Freshwater Ecosystems* 7: 165–177.

Wolff, W. J. (2000) The south-eastern North Sea: losses of vertebrate fauna during the past 2000 years. *Biological Conservation* 95: 209–217.

Wong, Y. S., Lan, C. Y., Chen, G. Z., et al. (1995) Effect of wastewater discharge on nutrient contamination of mangrove soils and plants. *Hydrobiologia* 295: 243–354.

Wong, Y. S., Tam, N. F. Y. & Lan, C. Y. (1997) Mangrove wetlands as wastewater treatment facility: a field trial. *Hydrobiologia* 352: 49–59.

Wood, C. M. & McDonald, D. G., eds. (1997) Global Warming: Implications for Freshwater and Marine Fish. Cambridge, UK: Cambridge University Press.

Wood, P. J. & Petts, G. E. (1999) The influence of drought on chalk stream macroinvertebrates. *Hydrological Processes* 13: 387–399.

Woodroffe, C. D. (1987) Pacific island mangroves: distribution and environmental settings. *Pacific Science* 41: 166–185.

Woodroffe, C. D. (1990) The impact of sea-level rise on mangrove shorelines. *Progress in Physical Geography* 14: 483–520.

Woodroffe, C. D. (1992) Mangrove sediments and geomorphology. In: *Tropical Mangrove Ecosystems*, ed.
A. I. Robertson & D. M. Alongi, pp. 7–41. Washington, DC, USA: American Geophysical Union.

Woodroffe, C. D., Thom, B. G. & Chappell, J. (1985) Development of widespread mangrove swamps in mid-Holocene times in northern Australia. *Nature* 317: 711–713.

Woodwell, G. M., Craig, P. P. & Johnson, H. A. (1971) DDT in the biosphere: where does it go? *Science* 174: 1101–1107.

Wooster, M. (2000) Niagara River progress (?). Great Lakes News 15: 17–18.

World Bank (2002) *Bangladesh: Climate Change and Sustainable Development*, Report No. 21104-BD. Dhaka, Bangladesh: The World Bank.

World Commission on Environment and Development (1987) *Our Common Future: The Brundtland Report.* Oxford, UK: Oxford University Press.

Worm, B. & Myers, R. A. (2003) Meta-analysis of cod–shrimp interactions reveals top-down control in oceanic food webs. *Ecology* 84: 162–173.

- Worm, B., Lotze, H. K., Bostrom, C., et al. (1999) Marine diversity shift linked to interactions among grazers, nutrients and propagule banks. *Marine Ecology Progress* Series 185: 309–314.
- Worthington, S. & Worthington, E. B. (1933) Inland Waters of Africa. London, UK: Macmillan.
- Wren, C. D., Harris, S. & Harttrup, N. (1995) Ecotoxicology of mercury and cadmium. In: *Handbook of Ecotoxicology*, ed.
 D. J. Hoffman, B. A. Rattner, G. A. Burton Jr & J. Cairns Jr, pp. 392–423. Boca Raton, FL, USA: Lewis Publishers.
- WRI (1998) *World Resources 1998–1999*. New York, USA: Oxford University Press.
- Wright, L. D. & Short, A. D. (1984) Morphodynamic variability of surf zones and beaches: a synthesis. *Marine Geology* 58: 93–118.
- Wright, S. (1955) Limnological Survey of Western Lake Erie, Special Scientific Report: Fisheries No. 139. Washington, DC, USA: US Fish and Wildlife Service.
- WTO (2003) World Trade Organization: Doha Declarations. Geneva, Switzerland: World Trade Organization.
- Wu, J. F., Sunda, W., Boyle, E. A. & Karl, D. M. (2000) Phosphate depletion in the western North Atlantic Ocean. *Science* 289: 759–762.
- Wu, M. Y., Hacker, S., Ayres, D. & Strong, D. A. (1999) Potential of *Prokelesia* spp. as biological control agents of English cordgrass, *Spartina anglica. Biological Control* 16: 267–273.
- Wu, R. S. S. (1999) Eutrophication, water borne pathogens and xenobiotic compounds: environmental risks and challenges. *Marine Pollution Bulletin* **39**: 11–22.
- Wubben, D. L. (2000) UV-induced mortality of zoea I larvae of the brown shrimp *Crangon crangon* (Linnaeus 1758). *Journal of Plankton Research* 22: 2095–2104.
- Wurtsbaugh, W. A. & Gliwicz, Z. M. (2001) Limnological control of brine shrimp population dynamics and cyst production in the Great Salt Lake, Utah. *Hydrobiologia* 466: 119–132.
- WWC (2000) World Water Vision: Making Water Everybody's Business. Commission report, World Water Council [www.document]. URL http://:www.watervision.org
- WWC (2002) UN consecrates water as public good, human right [www document]. URL http://www. worldwatercouncil.org/download/UM_water_publicgood. pdf
- WWC (2003a) Water management [www document]. URL http://www.worldwatercouncil.org/wwa_contents_fr.shtml
- WWC (2003b) World Water Council 3rd World Water Forum. Marseille, France: World Water Council.

- WWF (1993) Living rivers. Unpublished report. Zeist, the Netherlands: World Wide Fund for Nature.
- WWF (1999) *Living Planet Report*, ed. J. Loh, J. Randers, A. MacGillivray, *et al.* Gland, Switzerland: World Wide Fund for Nature.
- WWF (2006) Living Planet Report 2006, ed. J. Loh & S. Goldfinger. Gland, Switzerland: World Wide Fund for Nature.
- WWF & EU (2001) Implementing the EU Water Freshwater Directive: Seminar 2 – The role of wetlands in river basin management. Unpublished report, 9–10 November 2000. Brussels, Belgium: European Union.
- Wyatt, G. (2003) Proposed introduction of *Atriplex pedunculata* (annual sea purslane) to a tidally-influenced site on the Essex coast. *BSBI News* 92: 19–22.
- Wyllie-Echeverria, S. & Ackerman, J. D. (2003) The seagrasses of the Pacific Coast of North America. In: *World Atlas of Seagrasses*, ed. E. P. Green & F. T. Short, pp. 217–224. Berkeley, CA, USA: University of California Press.
- Wyllie-Echeverria, S., Olson, A. M., Hershman, M. J., eds. (1994) Seagrass Science and Policy in the Pacific NorthWest: Proceedings of a Seminar Series, (SMA 94-1) EPA Report No. 910/R-94-004. Washington, DC, USA: US Environmental Protection Agency.
- Wyman, R. L. (1990) What's happening to the amphibians? Conservation Biology 4: 350–353.
- Wynberg, R. P. & Branch, G. M. (1997) Trampling associated with bait-collection for sandprawns *Callianassa kraussi* Stebbing: effects on the biota of an intertidal sandflat. *Environmental Conservation* 24: 139–148.
- Xiao, X., Melillo, J. M., Kicklighter, D. W., et al. (1998) Transient climate change and net ecosystem production of the terrestrial biosphere. *Global Biogeochemical Cycles* 12: 345–360.
- Xu, C. Y. (1999) From GCMs to river flow: a review of downscaling methods and hydrological modelling approaches. *Progress in Physical Geography* 23: 229–249.
- Yang, H. & Zehnder, A. J. B. (2001) China's regional water scarcity and implications for grain supply and trade. *Environment and Planning A* 33: 79–95.
- Yang, H. & Zehnder, A.J. B. (2002) Water scarcity and food import: a case study for southern Mediterranean countries. *World Development* **30**: 1413–1430.
- Yap, H. T. (1992) Marine environmental problems: experiences of developing regions. *Marine Pollution Bulletin* 25: 37–40.

Yarnell, B. (1998) Integrated regional assessment and climate change impacts in river basins. *Climate Research* 11: 65–74.

Yendo, K. (1902) Kaiso Isoyake Chosa Hokoku. Suisan Chosa Hokoku 12: 1–33 (in Japanese).

Yendo, K. (1903) Investigations on 'Isoyake' (decrease of seaweed). *Journal of the Imperial Fisheries Bureau* 12: 1–33 (in Japanese).

Yendo, K. (1914) On the cultivation of seaweeds, with special accounts of their ecology. *Economic Proceedings of the Royal Dublin Society* 2: 105–122.

Yentsch, C. S., Yentsch, C. M., Cullen, J. J., et al. (2002) Sunlight and water transparency: cornerstones in coral research. Journal of Experimental Marine Biology and Ecology 268: 171–183.

Yin, K. D., Harrison, P. J., Chen, J., Huang, W. & Qian, P. Y. (1999) Red tides during spring 1998 in Hong Kong: is El Niño responsible? *Marine Ecology Progress Series* 187: 289–294.

Yon, D. & Tendron, G. (1981) Alluvial Forests of Europe. Brussels, Belgium: European Committee for the Conservation of Nature and National Resources.

Yuan, X. & Martinson, D. G. (2000) Antarctic sea ice extent variability and its global connectivity. *Journal of Climate* 13: 1697–1717.

Yuan, X. & Martinson, D. G. (2001) The Antarctic dipole and its predictability. *Geophysical Research Letters* 28: 3609–3612.

Zachos, J., Pagani, M., Sloan, L., Thomas, E. & Billups, K. (2001) Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292: 686–693.

Zahir, H. (2002) Status of coral reefs of Maldives. In: Coral Reef Degradation in the Indian Ocean: Status Report 2002, ed. O. Linden, D. Souter, D. Wilhelmsson & D. Obura, pp. 119–124. Kalmar, Sweden: CORDIO.

Zedler, J. B. (1993) Canopy architecture of natural and planted cordgrass marshes: selecting habitat evaluation criteria. *Ecological Applications* 3: 123–138.

Zedler, J. B. (1996) Costal mitigation in southern California: the need for a regional restoration strategy. *Ecological Applications* 6: 84–93.

Zedler, J. B. & Callaway, J. C. (1999) Tracking wetland restoration: do mitigation sites follow desired trajectories? *Restoration Ecology* 7: 69–73.

Zedler, J. B., Callaway, J. C., Desmond, J., et al. (1999) Californian salt marsh vegetation: an improved model of spatial pattern. *Ecosystems* 2: 19–35. Zedler, J. B., Callaway, J. C. & Sullivan, G. (2001) Declining biodiversity: why species matter and how their functions might be restored. *BioScience* 51: 1005–1017.

Zedler, J. B., Nordby, C. S. & Kus, B. E. (1992) The Ecology of Tijuana Estuary: A National Estuarine Research Reserve. Washington, DC, USA: NOAA Office of Coastal Resource Management, Sanctuaries and Reserves Division.

Zedler, J. B., Paling, E. & McComb, A. (1990) Differential salinity responses to help explain the replacement of native *Juncus kraussii* by *Typha orientalis* in Western Australian salt marshes. *Australian Journal of Ecology* 15: 57–72.

Zedler, P. H. (1987) The Ecology of Southern California Vernal Pools: A Community Profile, Biological Report No. 85 (7.11). Washington, DC, USA: US Fish and Wildlife Service.

Zedler, P. H. (2003) Vernal pools and the concept of 'isolated wetlands'. *Wetlands* 23: 597–607.

Zehnder, A. J. B., Yang, H. & Schertenleib, R. (2003) Water issues: the need for action at different levels. *Aquatic Sciences: Research across Boundaries* 65: 1–20.

Zenkevitch, L. (1963) *Biology of the Seas of the USSR*. London, UK: Allen & Unwin.

Zepp, R. G., Callaghan, T. V. & Erickson, D. J. (2003) Interactive effects of ozone depletion and climate change on biogeochemical cycles. *Photochemical and Photobiological Sciences* 2: 51–61.

Zhang, J. D., Rothrock, D. A. & Steele, M. (2000) Recent changes in Arctic sea ice: the interplay between ice dynamics and thermodynamics. *Journal of the American Meteorological Society* 13: 3099–3114.

Zieman, J. C. (1976) The ecological effects of physical damage from motor boats on turtle beds in Southern Florida. *Aquatic Botany* 2: 127–139.

Zimmerman, R. C., Kohrs, D. G., Steller, D. L. & Alberte, R. S. (1997) Impacts of CO₂ enrichment on productivity and light requirements of eelgrass. *Plant Physiology* 115: 599–607.

Zinke, A. & Gutzweiler, K.-A. (1990) Possibilities for regeneration of floodplain forests within the framework on the flood-protection measures on the Upper Rhine, West Germany. *Forest Ecology and Management* 33/34: 13–20.

Zobell, C. E. (1971) Drift seaweeds on San Diego county beaches. Nova Hedmigia 32: 269–314.

Zoltai, S. C. (1988) Wetland environment and classification. In: *Wetlands of Canada*, ed. National Wetlands Working Group, pp. 3–26. Montreal, Quebec, Canada: Polyscience.

- Zong, Y. & Chen, X. (2000) The 1998 flood on the Yangtze, China. *Natural Hazards* 22: 165–184.
- Zwally, H. J., Comiso, J. C., Parkinson, C. L., Cavalieri, D. J. & Gloersen, P. (2002) Variability of Antarctic sea ice 1979–1998. *Journal of Geophysical Research* 107(C5): 10, doi: 1029/2000JC000733.
- Zwally, H. J., Parkinson, C. L. & Comiso, J. C. (1983) Variability of Antarctic sea ice and changes in carbon dioxide. *Science* 220: 1005–1012.
- Zwiers, F. W. & Kharin, V. V. (1998) Changes in the extremes of the climate simulated by CCC GCM2 under CO₂doubling. *Journal of Climate* 11: 2200–2222.
- Zwiers, F. W. (2002) The 20-year forecast. *Nature* **416**: 690–691.