

SOUTHERN

DNA extraction

- freeze and homogenize 50 mg – 1g of plant material (mortar). Transfer ground plant material into 2 ml tube (plant material should fill tube slightly over 0.5 ml marking)
- add 1ml extraktion buffer, mix (do not vortex! dissolve material by “shaking in circles”, inverting....) inkubate 10 min at 65 °C.
- add 300 µl 5M potassiumacetate (**no** pH adjustment)
- inkubate 10 min on ice
- spin 10 min at 15000 rpm, 4°C
- transfer supernatant to new tube, add 800 µl phenol/chloroform/isoamylalcohol, mix gently by inverting tube 2-3x, incubate 1 h on shaker (gently) - (min. 15 min!!)
- spin 5 min at 12000 rpm
- **aequous phase** in new tube, add 800 µl isopropanol, mix gently by inverting tube 2-3x, inkubate 10 min at RT
- spin 10 min bei 15000 rpm
- wash pellet with 1 ml ETOH 70%
- spin 10 min bei 15000 rpm
- remove ETOH, dry pellet
- resuspend pellet in 16-20 µl sterile dH₂O

Restriction (adjust protocol if necessary)

- 15 µl DNA
- 2 µl Puffer (10 x)
- 1µl Enzym
- 1 µl RNase

37 °C o.N.

Gelelektrophoresis

- Restriktionsansatz komplett auftragen
- 0.8 - 1%iges TAE Gel, ü.N. bei 30 V
- Gel Foto mit Linieal!

Extraction buffer

final conc.		stock conc.	100ml buffer vol.	500 ml buffer vol.
100 mM	Tris-HCl pH 8.0	1 M	10 ml	50 ml
50 mM	EDTA	0.5 M	10 ml	50 ml
500 mM	NaCl	5 M	10 ml	50 ml
1.5 %	SDS	10 %	15 ml	75 ml
	H2O		55 ml	275 ml

SOUTHERN Blot

- Place gel in tray and add 200-300 ml denaturing buffer, gently shake 45 min at room temp.
- wash gel 2x with dH₂O
- add 200 – 300 ml neutralizing buffer, gently shake 45 min at room temp.
- wash gel 2x with dH₂O
- build blot (use Nylon Membrane positively charged (Roche # 11417240001: € 415.15)
- transfer with 20 x SSC over night
- briefly wash membrane in 3x SSC, dry on filter paper
- UV-Crosslink
- prehybridize membrane: ~ 50 ml at 42°C min. 2-4h or over night
- label probe (Amersham Megaprime™ DNA Labeling System (GE Healthcare, #RPN1604: €447) and add to prehyb “setup”, incubate over night at 42°C
- wash membrane: 3x at 65 °C with 3xSSC + 0.1% SDS
- measure radioactivity (– when needed: add another wash step)
- place membrane between foil and place in phospho imager box + screen
- phosphoimager quantification of signals after 4h – 3d

Buffer:

20 x SSC:

3M NaCl + 0.3 M Na-citrate

denaturing buffer (Southern):

1.5 M NaCl (87.6 g/L) + 0.5 M NaOH (20.05 g/L)

neutralizing buffer (Southern):

1.5 M NaCl (87.6 g/L) + 0.5 M Tris (60.7 g/L)

(pre-)hybridizing buffer:

for 500 ml: 1g PVP (polyvinylpyrrolidone)
 1g ficoll
 1g BSA
 240ml buffer “N”

when completely dissolved add

240ml formamide

6 ml herring sperm DNA (heat denature prior to adding!)

store buffer at 4°C

buffer “N”:

1.8 M NaCl
100 mM NaH₂PO₄
10 mM EDTA
1.2 % SDS
adjust pH to pH 7.0