

# MERCURY

By William E. Brooks

**Domestic tables were prepared by Subina W. Pandey, statistical assistant, and the world production table was prepared by Linder Roberts, international data coordinator.**

In 2003, mercury was recycled in-plant, as home scrap, to supply the chlorine-caustic soda industry, which is the largest domestic end use for mercury in the United States. Global human health and environmental concerns about anthropogenic mercury releases brought about an overall market shift toward nonmercury alternatives for chlorine-caustic soda production and other mercury end uses. Some mercury was imported from Germany and Peru. Even though the primary ore of mercury, cinnabar, is found in several States in the United States, there is no primary production of mercury, and the last mine to produce mercury as its chief commodity, the McDermitt Mine in Nevada, closed in 1992. Mercury may also be obtained as a byproduct of gold or lead mining; production data, however, are not available. In 1971, the U.S. Environmental Protection Agency (EPA) indicated that mercury was a hazardous air pollutant, which added to the concern about the toxic effects of mercury on human health. The United States had 4,436 metric tons (t) of mercury stockpiled by the Defense Logistics Agency (DLA) at yearend 2003.

## Health, Legislation, and Government Programs

Mercury is known to broadly affect human health, and anthropogenic sources of mercury releases include chlorine-caustic soda production, coal-fired powerplants and industrial boilers, hazardous waste disposal, municipal and medical waste disposal, and recycling (Clean Air Network, 2000§;<sup>1</sup> Mercury Instruments, 2003§). Volcanic activity, artisanal gold mining, and mercury mining are other sources of mercury releases on a global scale (Hylander and Meili, 2002). Trace amounts of mercury may be present in minerals in barite drilling mud used in offshore oil and gas exploration (Neff, 2002§). The nervous system, brain, kidneys, and fetal development may be adversely affected by exposure to mercury, and a link between autism in children and mercury in the environment seems likely (Hightower, 2001§). In the 1960s, the effects of mercury on human health in Japan, or “Minamata Disease,” caused headlines worldwide (Bunce and Hunt, 2003§), and recently, the United Nations Environmental Programme (2002§) completed a Global Mercury Assessment in which issues that concern mercury in the environment, health effects in less developed regions, the global fishing industry, and sources of mercury releases were examined. Mercury is now studied and regulated by environmental and government health agencies worldwide (Porcella and others, 1995; Hylander, 2002§). The environmental impact of mercury in a variety of global settings, such as wetland sediments in California and Connecticut, black

shales in Finland, mine waste in the Philippines, mercury in the soil at an emission source, and mercury accumulation in snow were some of the featured articles in a special issue of *Environmental Geology* (2003) on mercury.

In the United States, the Omnibus Mercury Emissions Reduction Act of 2003 mandates reduced mercury emissions from all major sources, directs the EPA to issue new standards, and sets a timetable so that mercury emissions are reduced (Leahy and Snowe, 2003§). This Act addresses emission standards from a variety of sources and supports mercury research, encourages international collaboration, and prohibits sale of the 4,436 t of mercury held by the DLA on the global market.

In July 2002, the EPA specifically addressed mercury emissions from chlorine-caustic soda plants by using standards given in the Clear Air Act that reflect the maximum achievable control technology which would reduce emissions by 4,100 kilograms per year (kg/yr) (U.S. Environmental Protection Agency, 2002§). In 2000, the Mercury Policy Project, which is a public interest group, urged zero mercury use and urged the U.S. Department of Defense and the U.S. Department of Energy to accept 130 t of mercury from a closed chlorine-caustic soda plant in Maine into the DLA stockpile; the effort, however, apparently failed, and the mercury may have reached the global market (Bender, 2000§).

The EPA set a limit of 2 parts per billion (ppb) mercury in drinking water; the U.S. Food and Drug Administration set a maximum of 1 part per million of methylmercury (a toxic bacterial transformation of mercury) in seafood; and the U.S. Occupational Safety and Health Administration set a limit of 0.05 milligram of metallic mercury vapor per cubic meter in workplace air for an 8-hour shift and 40-hour workweek (Agency for Toxic Substances and Disease Registry, 2003b§). The American Medical Association resolved that the mercury content of foods, especially fish, be labeled and that physicians educate their patients as to the potential dangers of mercury ingestion (McCaffree, 2003§).

Research presented at a recent EPA National Forum on Contaminants in Fish showed that for pregnant women mercury in the umbilical cord blood may be 70% higher than the mercury content of the mother’s blood (Mahaffey, 2004§). Because one in six women of childbearing age in the United States has unsafe blood mercury levels, more than 600,000 babies are born every year at risk of mercury-related health problems (Physicians Committee for Responsible Medicine, 2004). Mercury may be traced to bioaccumulation of the metal from atmospheric mercury pollution that drifts worldwide from anthropogenic sources. For example, Canadian government studies showed elevated levels of pollution in the Arctic and high levels of mercury in the blood and breast milk of Inuit mothers

<sup>1</sup>References that include a section mark (§) are found in the Internet References Cited section.

in Nunavut (Brown, 2004). A number of regulations have been proposed that would require coal-fired powerplants, which may contribute 48 t of mercury to the atmosphere, to cut emissions by target dates (Gugliotta, 2004).

U.S. Geological Survey (USGS) research on mercury in coal, mercury mines in Alaska and California, mercury contamination from hydraulic mining in California, and volcanic emissions of mercury was included in Gray (2003). The environmental impact of mercury associated with mineral districts worldwide and byproduct mercury from gold-silver and massive sulfide deposits was described by Rytuba (2003). Also, USGS research contributed to understanding the abundance of mercury in domestic coal (Tewalt and others, 2001), the health impacts of coal combustion (Finkelman, 2000), and the mercury content of coal in the world market (Finkelman and others, 2001). The USGS collaborates with other Federal and State agencies, universities, and tribal nations in the National Atmospheric Mercury Deposition Network, which is a national program that monitored mercury deposition through a network of standardized monitoring sites (Nilles, 2000, 2003§). USGS scientists also examined glacial ice cores as part of studies focused on the history of global mercury contamination (Krabbenhoft and Schuster, 2002). Media reports of mercury contamination from minerals in barite drilling mud used by oil and gas drill rigs in the Gulf of Mexico prompted USGS research into bottom sediment geochemistry of Lake Ponchartrain, LA; the data showed only a few samples with elevated mercury content (Manheim, 2002§). The USGS National Mercury Project indicates that effective policy decisions regarding mercury emissions require research that will provide better understanding of its cycle in the environment (Pinsker, 2003; Krabbenhoft and Herrmann, 2000§).

Academic, tribal, and research organizations and other Government agencies collaborate in periodic USGS/EPA Mercury Roundtable teleconferences (Krabbenhoft and Herrmann, 2000§). Topics included the bioaccumulation of methylmercury in fish, quantifying methylmercury in fish with regard to species and size, sources of mercury in the Everglades, and the toxic legacy of mercury mining in California.

The USGS Toxic Substances Hydrology Program and the Mine Drainage Interest Group characterized mercury contamination, bioaccumulation, and environmental problems related to mining across the country (Suchanek and Marvin-DiPasquale, 2003§). USGS study sites in Virginia include closed pyrite mines near Contrary Creek and Prince William Forest Park (Cravotta and others, 2003§).

The EPA and National Wildlife Federation studies showed that rain falling on Chicago's North Shore may contain up to 32 times the amount of mercury that is considered to be safe. This is of concern because of the bioaccumulation of mercury in fish (Pierre, 2003; McCann, 1999§). Coal-fired power plants in the region are considered to be the source of the mercury (Pennsylvania Department of Environmental Protection, 2003§). Warnings about mercury content of fish have been posted in some parts of Florida (Mercury Roundtable, oral commun., June 17, 2003). In 1989, a consumption advisory for mercury and organic chemicals for fish caught in the Anacostia and Potomac Rivers was issued by the District of Columbia (Agency for

Toxic Substances and Disease Registry, 2003a§). The Virginia Department of Health issued warnings to limit the amount of fish consumed from several Virginia rivers because of elevated levels of mercury (Washington Post, 2004). Forty-four States have posted warnings about mercury contamination in lakes, streams, and rivers (Physicians Committee for Responsible Medicine, 2004). In the Pacific Northwest, USGS research indicated that the mercury content of fish in North Cascades National Park is approaching human health warning levels of 300 ppb and may pose a potential health risk (Pynn, 2004; Moran and Black, 2003§).

Proposed Clear Skies legislation in Congress will require powerplants to reduce overall pollution levels by 70% by 2018 (Planin, 2003a; U.S. Environmental Protection Agency, 2000a§). The Clean Air Planning Act, which is an alternative plan, would cut powerplant emissions of mercury, sulfur dioxide, and other pollutants by larger amounts by 2012 (Gugliotta and Planin, 2003). Mercury emissions from coal-burning plants are not regulated under Federal law and attempts to classify mercury as a hazardous pollutant, similar to asbestos and lead, were met with counter arguments that mercury be considered under a less stringent part of the Clean Air Act that includes pollutants that cause smog and acid rain and are not as toxic to humans (Lee, 2004). Approximately 48 metric tons per year (t/yr) of mercury were emitted annually by the Nation's 11,000 coal-fired powerplants, and emissions of mercury by some industrial sources pose an increasing health danger to children. Pollution from powerplants, which includes mercury, sulfur dioxide, and nitrogen oxides, may result in the premature deaths of thousands in the United States (Eilperin, 2004b). Legislation has been proposed that would cut emissions to 34 t in 2010 and 15 t by 2018 (Fialka, 2003). The EPA is also concerned about the approximately 65 t of mercury that is unaccounted for by the nine domestic chlorine-caustic soda plants that still rely on 1950s mercury cell technology (Eilperin, 2004a).

Mercury emission guidelines for hospital and medical waste incinerators are given by the U.S. Environmental Protection Agency (2003a§). Similar guidelines for electric steam generating units were also provided in the Federal Register (U.S. Environmental Protection Agency, 2000b§).

Approximately 6 million mercury thermometers, each of which contains approximately 0.7 gram of mercury, were sold in the United States during a single year (Goldstein, 2000). Continued use of such thermometers presents a health risk if broken in the user's home or if the thermometers are dumped in landfills or incinerated. Environmental and health organizations have petitioned Congress to limit the use of mercury fever thermometers and to improve the collection and management of mercury (Bill Summary, 2002§; Lyon, 2001§). Therefore, sale of mercury thermometers has been banned and the use of nonmercury thermometers has been encouraged (U.S. Environmental Protection Agency, 2003b§). For example, some local environmental agencies have recently offered to replace mercury thermometers with digital thermometers (Washington Post, 2003b). Mercury may still be found in some school laboratories. Recently, 250 milliliters of mercury was taken from a high school lab in Washington, DC, and was scattered around the school. This caused evacuation, closure, and EPA testing and

decontamination of the school (Fahrenthold, 2003). At another school in Maryland, a thermometer was accidentally broken. This caused decontamination of the classroom, school bus, and testing of the student's clothing (Washington Post, 2003a).

Dental amalgam also uses mercury, and how much mercury is released into the environment by removal or replacement of mercury amalgam fillings is unknown (District of Columbia Dental Society, 2003§). Approximately 12,000 kilograms of amalgam were sold in the United States in 2001 (Maine Department of Environmental Protection, 2003§). The Watson-Burton bill (H.R. 1680: Mercury in Dental Filling Disclosure and Prohibition Act) seeks to prohibit the introduction of mercury for dental fillings into interstate commerce after 2008 (Burton, 2004§); some public health organizations, however, require that dental amalgam be used (Carlton, 2004§). The use of amalgam is declining, and composite resin substitutes are available.

Soap and cosmetics that contain mercury are manufactured by some European companies and sold as skin-lighteners in Africa, Asia, and the Caribbean. Both manufacturers and users may be subject to skin damage or mercury-related health problems (Mercury Soap Campaign, 2004§). Some Caribbean groups may use mercury as a folk remedy (Goldstein, 2003). Mercury use as a fungicide in paints has been discontinued. The use of mercury batteries, which were once widely used in watches, cameras, and hearing aids, has declined because of new technologies and concern for the environment (Battery-Index, 2003§).

Volcanic activity also contributes mercury to the global environment, and USGS research shows that mercury is lost from the forest floor during forest fires (Woodruff and Cannon, 2001).

At yearend 2003, the DLA had an inventory of 4,436 t of mercury at several sites in the United States and until 1994, this mercury was available for sale. Mercury sales have been suspended in response to environmental concerns. In early 2004, the DLA indicated that the mercury would be consolidated at one site (Joseph Johnson, Specialist, Defense Logistics Agency, written commun., April 30, 2004).

## Production

Mercury has not been mined in the United States since closure of the McDermitt Mine in Nevada in 1992 (O'Driscoll, 2002§). The mine is now included on a list of EPA Superfund Sites (ToxicAlert, 2003§). Byproduct mercury is generated as a part of some domestic gold mining and smelting operations (Rogers, 2000§). In addition, mercury is recovered from domestic and imported scrap or byproduct mercury by recycling plants in the United States.

There are mercury occurrences in California (Dickson and Tunell, 1968; Linn, 1968), Idaho (Dickson and Tunell, 1968), Nevada (Dickson and Tunell, 1968; Fisk, 1968; Bateman, 1988), Texas (Bauer, 2000), and Utah. The geoenvironmental impact of mercury in Alaska and California is an ongoing USGS (2003§) research project. Occurrences of cinnabar were found in the Mercury District of Southwest Arkansas, and the USGS presented geochemical data on this area as a part of a national geochemical database (Moravec and Smith, 1997§).

In the United States, the largest amount of byproduct mercury comes from lode gold mining in Nevada. Data on the amount

of byproduct mercury produced are not available. Sales of mercury retort systems permit the inference that byproduct mercury is recovered from ore processed at six mines in Nevada (Summit Valley Equipment and Engineering, Inc., 2004§). Mine owners in Nevada have voluntarily begun to reduce mercury emissions by 50% within a 3-year period (Gold, 2002).

Byproduct mercury from Peru's Pierina gold mine is recovered, carefully packed, transported, and shipped to the United States for recycling. All handlers of this mercury, even customs agents, receive training in the safe handling of mercury (Michael Merry, logistics superintendent, Barrick Peru, oral commun., May 3, 2004). Byproduct mercury from Peru's Yanacocha gold mine is sent to Spain for processing.

Mercury is used in artisanal placer gold mining worldwide to amalgamate gold flakes and mercury may be released during mining and amalgam treatment in some countries. In Brazil, burning the mercury-laden amalgam led to elevated levels of mercury in the urine of gold shop workers that was 20 times greater than the limit of 50 micrograms/liter established by the World Health Organization (Malm, 1998). In Venezuela, processing centers have been established where artisanal miners can take their gold-mercury concentrates for retorting and recovery of the gold and mercury. This reduces health risks to miners, who commonly burn their amalgam in the open air to drive off the mercury, and limits mercury releases to the environment (Viega, 1997§). Sediment samples taken in streams near artisanal gold workings or amalgamation sites in Peru gave high mercury content (Medina, 2001§). In Ghana, artisanal gold miners may inhale the vapors from the heated pots that are used to purify the gold-mercury amalgam. Any mercury discarded in the streams bioaccumulates in the fish, which is widely consumed, and may lead to physical and mental problems (Harkinson, 2003§).

The USGS led mercury and mining studies in California (Alpers, 2002§), Nevada (Gray, Crock, and LaSorsa, 2002), the Philippines (Gray and others, 2003), and Suriname (Gray and others, 2002). Use of mercury in Latin American mining operations was addressed at a conference in Vienna, Austria, in 1997. It was estimated that more than 200 t/yr of mercury are released through artisanal mining in Latin America (Veiga, 1997§). A spill of byproduct mercury recovered from gold processing in Peru in 2000 gained worldwide attention (Drillbits and Tailings, 2000§; Griffin, 2000§).

## Consumption

Chlorine production, which is the major end use of mercury, provides an important industrial chemical for use in the pulp and paper industry. It is also used for water treatment, defense against biological attacks, antibiotics, microprocessors and computer housings, flak jackets, helmets, and bullet-proof glass (Chlorine Institute, 2003§). Elevated mercury concentrations were found in mud and biota in streams that received discharges from mercury cell chlorine-caustic soda plants (Georgia State Water Quality Control Board, 1971). Losses of mercury in the chlorine production process were initially as high as 200 grams of mercury per metric ton of chlorine output in the 1960s, but only 0.2 gram of mercury was reported lost per ton of chlorine

produced today (Bunce and Hunt, 2003§). Domestic chlorine-caustic soda plants use about 100 t/yr of mercury to replace mercury lost in the manufacturing process. But what happens to the mercury that is replaced is unknown (Planin, 2003b). The EPA is now concerned that as much as 65 t of mercury cannot be accounted for and may have been released into the environment (Eilperin, 2004b).

A limited amount of mercury was imported from Germany (19 t) and Peru (19 t) during 2003. The total amount of mercury imported in 2003 was 46 t which is much less than the 209 t of mercury imported in 2002. The change may be in response to a shutdown of mercury cell chlorine-caustic soda plants in Australia in 2001 (ACTED Consultants, 2004§). Closure of these plants may have been the source of the 107 t of mercury brought into the United States from Australia in 2002. The domestic chlorine-caustic soda industry has set a goal of reducing mercury use by 50% before 2005, in comparison with the base years of 1990 to 1995. Domestic mercury consumption is estimated to be 33% to 50% for chlorine manufacture, 33% to 50% for electronics, and the balance for other uses, such as dental applications, fluorescent lamps, and switches.

## Recycling

The chief end use of mercury has been as a cathode in the electrolytic production of chlorine and caustic soda from brine. Sodium is removed from the sodium-mercury amalgam, and the mercury is recycled within the chlorine-caustic soda plant as home or in-plant scrap. Chlorine and sodium are important industrial chemicals with a variety of uses (Amato, 1994). Diaphragm and membrane cells are alternative methods for chlorine-caustic soda production and do not use mercury (Roskill Information Services Ltd., 1990). Closure of mercury-based chlorine-caustic soda plants and subsequent retirement, recycling, sale, or storage of the approximately 3,000 t of mercury contained in these domestic plants is a serious concern (Raloff, 2003§).

In previous years, approximately 46 t of mercury was used by the chlorine-caustic soda industry followed by 60 t for switches, 44 t for dental amalgam, 28 t for lighting, and 22 t for measuring instruments (Bender, 2002§). Mercury may be recycled from these products.

Mercury used in tilt switches in cars (McCann, 2002; Natural Resources Council of Maine, 2003§), fluorescent lamps (Abernathy, 2003), and thermostats (Massachusetts Department of Environmental Protection, 2004§) is of environmental concern because of the potential for mercury releases during scrapping, waste treatment, or demolition. The EPA has required foundries that shred vehicles to create policies on how mercury is minimized in those vehicles before the furnace becomes contaminated (American Metal Market, 2003). Mercury may also be recovered and recycled from unbroken fluorescent lamps.

In 2003, mercury recycling companies included AERC Recycling, VA; Bethlehem Apparatus Company, PA; D.F. Goldsmith Chemical and Metal Corporation, IL; Mercury Waste Solutions, MN; and Onyx Environmental Services, WI (Bethlehem Apparatus Company, 2003§; D.F. Goldsmith

Chemical and Metal Corporation, 2003§; Mercury Waste Solutions, 2003§; AERC Recycling, 2004§; Onyx Environmental Services, 2004§). A list of more than 50 individuals and companies that recycle mercury is available (Mercury Recyclers, 2002§).

## World Review

In 2003, world mercury production was estimated to be 1,530 t/yr, which was less than the 2,200 t/yr average world mercury production from 1990 to 2000. Production estimates have a high degree of uncertainty because most countries do not report primary or byproduct production data because of environmental and health concerns. Algeria, China, Kyrgyzstan, and Spain were the apparent leaders in world production of mercury; production from China estimated to be 610 t in 2003. Minas de Almaden (Spain) can produce 50,000 flasks<sup>2</sup> per year and new production was reported from Spain (Metal Bulletin, 2003). Both Algerian and Spanish producers were negotiating prices in the range of \$175 to \$195 per flask (Metal-Pages, 2003b). The U.S. domestic price per flask was \$345 in June, 2004 (Platts Metals Week, 2004). Demand was strong in China and the Asian market, but information on use is limited. Chinese mercury production began to rise in 2002 (Metal-Pages, 2003a). Becton, Dickinson and Co. indicated that it will stop production of glass mercury thermometers at its plant in Brazil and will stop purchasing thermometers from China (Goldstein, 2000). Environmental concern caused a thermometer plant in India to close and the waste material to be transported to the United States for recycling (Marley, 2003).

Mercury releases in India were estimated to range between 60 to 70 t/yr (Bahuguna, 2003§) and 150 to 200 t/yr (Mercury in India, 2003§) as a result of chlorine production. The obligatory dismantling of 47 chlorine plants in Europe by 2007 will result in 15,000 t of mercury that must be managed (Faversham House Group, 2003§).

The Swiss company Batrec, founded in 1989, was Europe's leading recycler of materials containing mercury and produces mercury, zinc, and other metals from processing 3,300 t of batteries and sludges annually (Beck, 2004).

## Outlook

Mercury demand and production will continue to be affected by ongoing global environmental and human health concerns. Technological advances and changes in the chlorine-caustic soda industry will result in a decline in mercury use as alternative production systems are used and mercury cell plants are closed in Europe (2007), the United States, and India, thereby releasing large amounts of mercury for storage or sale. Voluntary reduction of mercury emissions by copper, gold, and zinc mining companies, stricter controls of mercury use in artisanal gold mining, and careful monitoring of mercury releases by

<sup>2</sup>Some international data and dealer prices are reported in flasks. Since 1927, mercury has commonly been measured and priced by the flask (approximately 76 pounds), a measure that is related to early mercury mining and processing in Spain (Myers, 1951). One metric ton of mercury contains approximately 29 flasks.

the chlorine-caustic soda industry will lessen releases to the environment and may affect mercury prices. Recycled mercury from dismantled mercury cell chlorine-caustic soda plants, byproduct mercury from gold mining, and mercury contained in the National Defense Stockpile will be more than adequate to meet domestic needs.

## References Cited

- Abernathy, P.W., 2003, Following the light: Resource Recycling, February, p. 24-27.
- Amato, Ivan, 1994, Crusade to ban chlorine: Garbage, summer, p. 32-39.
- American Metal Market, 2003, Foundries told to keep note of anti-mercury procedures: American Metal Market, v. 111, no. 46-5, November 21, p. 7.
- Bateman, P.W., 1988, Geological and geochemical characteristics of selected mercury deposits in Nevada and their relationship to precious metal deposits: Golden, CO, Colorado School of Mines, Ph.D. dissertation T-3504, 275 p.
- Bauer, E.M., 2000, Old Texas mercury mines attract tourists and danger: Washington Post, November 20, p. A3.
- Beck, Manfred, 2004, Batrec, recovering metals: Recycling International, May, no. 4, p. 24-26.
- Brown, D.L., 2004, Poisons from afar threaten Arctic mothers and traditions: Washington Post, April 11, p. A25.
- Dickson, F.W., and Tunell, George, 1968, Mercury and antimony deposits associated with active hot springs in the western United States, in Ridge, J.D., ed., The Graton-Sales volume, Ore deposits of the United States, 1933-1967: New York, NY, American Institute of Mining, Metallurgical and Petroleum Engineers, p. 1673-1701.
- Eilperin, Juliet, 2004a, EPA to probe missing mercury: Washington Post, May 30, p. A14.
- Eilperin, Juliet, 2004b, Tougher regulations urged on power plant emissions: Washington Post, June 10, p. A3.
- Environmental Geology, 2003, Special issue, Mercury: Environmental Geology, v. 43, no. 3, January, 363 p.
- Fahrenthold, D.A., 2003, Mercury is stolen, spread at Ballou High: Washington Post, October 3, p. B5.
- Fialka, J.J., 2003, Mercury threat to children rising says an unreleased EPA report: Wall Street Journal, February 20, p. A1.
- Finkelman, R.B., 2000, Health impacts of coal combustion: U.S. Geological Survey Fact Sheet FS-094-00, 2 p.
- Finkelman, R.B., Warwick, P.D., and Pierce, B.S., 2001, The world coal quality inventory: U.S. Geological Survey Fact Sheet FS-155-00, 4 p.
- Fisk, E.L., 1968, Cordero Mine, Opalite mining district, in Ridge, J.D., ed., The Graton-Sales volume, Ore deposits of the United States, 1933-1967: New York, NY, American Institute of Mining, Metallurgical and Petroleum Engineers, p. 1574-1591.
- Georgia State Water Quality Control Board, 1971, Mercury pollution investigation in Georgia: Atlanta, GA, Georgia State Water Quality Control Board, 35 p.
- Gold, 2002, US mine is praised for green issues: Gold, The Magazine of the World Gold Council, issue 8, p. 6.
- Goldstein, Avram, 2000, District joins effort to liquidate mercury: Washington Post, September 27, p. B2.
- Goldstein, Avram, 2003, Blood tests find 10 people with elevated mercury: Washington Post, October 15, p. B1.
- Gray, J.E., ed., 2003, Geologic studies of mercury by the U.S. Geological Survey: U.S. Geological Survey Circular 1248, 41 p.
- Gray, J.E., Labson, V.F., Weaver, J.N., and Krabbenhoft, D.P., 2002, Mercury and methylmercury contamination related to artisanal gold mining, Suriname: Geophysical Research Letters, v. 29, no. 23, p. 2105-2113.
- Gray, J.E., Crock, J.G., and Lasorsa, B.K., 2002, Mercury methylation at mercury mines in the Humboldt River Basin, Nevada, USA: Geochemistry-Environment, Exploration, Analysis, v. 2, p. 143-149.
- Gray, J.E., Greaves, I.A., Bustos, D.M., and Krabbenhoft, D.P., 2003, Mercury and methylmercury contents in mine-waste calcine, water, and sediment collected from the Palawan Quicksilver Mine, Philippines: Environmental Geology, v. 43, p. 298-307.
- Gugliotta, Guy, and Planin, Eric, 2003, EPA withholds air pollution analysis: Washington Post, July 1, p. A3.

- Hylander, L.D., and Meili, Markus, 2002, 500 years of mercury production—Global annual inventory by region until 2000 and associated emissions: The Science of the Total Environment, v. 304, p. 13-27.
- Krabbenhoft, D.P., and Schuster, P.F., 2002, Glacial ice cores reveal a record of natural and anthropogenic atmospheric mercury deposition for the last 270 years: U.S. Geological Survey Fact Sheet FS-051-02, 2 p.
- Lee, Jennifer, 2004, Questions about a market system for mercury: The New York Times, February 24, p. A1.
- Linn, R.K., 1968, New Idria mining district, in Ridge, J.D., ed., The Graton-Sales volume, Ore deposits of the United States, 1933-1967: New York, NY, American Institute of Mining, Metallurgical and Petroleum Engineers, p. 1623-1649.
- Malm, Olaf, 1998, Gold mining as a source of mercury exposure in the Brazilian Amazon: Environmental Research, v. 77, p. 73-78.
- Marley, Michael, 2003, US taking mercury waste from India: American Metal Market, v. 3, no. 21, May 26, p. 11.
- McCann, Joseph, 2002, Long road ahead to mercury solution: American Metal Market, v. 110, no. 52, March 18, p. 14.
- Metal Bulletin, 2003, Minas de Almaden works toward restart: Metal Bulletin, May 12, p. 9.
- Myers, D.K., 1951, History of the mercury flask: Journal of Chemical Education, v. 28, March, p. 127.
- Nilles, M.A., 2000, Atmospheric deposition program of the U.S. Geological Survey: U.S. Geological Survey Fact Sheet 112-00, 2 p.
- Physicians Committee for Responsible Medicine, 2004, Mercury is such a potent neurotoxin: Washington Post, May 18, p. A17.
- Pierre, R.E., 2003, High mercury levels are measure of poor rain quality in Chicago: Washington Post, June 1, p. A2.
- Pinsker, L.M., 2003, In search of the mercury solution: Geotimes, v. 48, no. 8, August, p. 16-21.
- Platts Metals Week, 2004, Weekly prices: Platts Metals Week, June 7, v. 75, no. 23, p. 19.
- Planin, Eric, 2003a, Crackdown urged on coal pollution: Washington Post, April 22, p. A4.
- Planin, Eric, 2003b, EPA issues new rule for mercury: Washington Post, December 20, p. A7.
- Porcella, D.B., Huckabee, J.W., and Wheatley, Brian, eds., 1995, Mercury as a global pollutant: London, United Kingdom, Kluwer Academic Publishers, 703 p.
- Pynn, Larry, 2004, Mercury levels in park fish are a health risk: Vancouver [British Columbia, Canada] Sun, May 3, p. B1.
- Roskill Information Services Ltd., 1990, The economics of mercury (7th ed.): London, United Kingdom, Roskill Information Services Ltd., 128 p.
- Rytuba, J.J., 2003, Mercury from mineral deposits and potential environmental impact: Environmental Geology, v. 43, no. 3, January, p. 326-338.
- Tewalt, S.J., Bragg, L.J., and Finkelman, R.B., 2001, Mercury in U.S. coal—Abundance, distribution, and modes of occurrence: U.S. Geological Survey Fact Sheet FS-095-01, 2 p.
- Washington Post, 2003a, Mercury spills on student, bus: Washington Post, November 23, p. C3.
- Washington Post, 2003b, Old thermometers exchanged for digitals: Washington Post, December 25, p. B3.
- Washington Post, 2004, High mercury, PCB levels in some fish: Washington Post, October 30, p. B3.
- Woodruff, L.G., and Cannon, W.F., 2001, The effect of fire on mercury and carbon in forest soils—Results from northern Michigan and Minnesota: Geological Society of America Abstracts with Programs, v. 33, no. 6, p. 186.

## Internet References Cited

- ACTED Consultants, 2004, Chlorine and caustic soda in Australia, accessed June 10, 2004, at URL [http://www.chemlink.com.au/caustic\\_oz.htm](http://www.chemlink.com.au/caustic_oz.htm).
- AERC Recycling, 2004, AERC recycling solutions, accessed June 4, 2004, at URL <http://www.aercrecycling.com/home>.
- Agency for Toxic Substances and Disease Registry, 2003a, Health consultation, Anacostia River Initiative, accessed May 30, 2003, at URL [http://www.atsdr.cdc.gov/HAC/PHA/anacostia/ana\\_pl.html](http://www.atsdr.cdc.gov/HAC/PHA/anacostia/ana_pl.html).
- Agency for Toxic Substances and Disease Registry, 2003b, ToxFAQs for mercury, accessed June 16, 2003, at URL <http://www.atsdr.cdc.gov/tfacts46.html>.

- Alpers, C.N., 2002, Mercury and arsenic contamination associated with abandoned mine lands—South Yuba Watersheds, accessed April 9, 2003, at URL <http://ca.water.usgs.gov/projects00/ca553.html>.
- Bahuguna, N.J., 2003, Mercury—Danger signals, accessed, June 2, 2003, at URL <http://www.boloji.com/wfs/wfs/090.htm>.
- Battery-Index, 2003, Mercury batteries, accessed June 2, 2003, at URL <http://battery-index.com/mercury-batteries.shtml>.
- Bethlehem Apparatus, 2003, Recovery/recycling service for hazardous and universal mercury waste, accessed June 3, 2003, at URL <http://www.bethlehemapparatus.com>.
- Bender, M.T., 2000, Retirement of mercury, A presentation to the Great Lakes Binational Toxics Strategy Stakeholders Forum, accessed June 1, 2004, at URL <http://www.mercurypolicy.org/new/documents/3mbretjg.pdf>.
- Bender, M.T., 2002, Dentist the menace?, accessed August 18, 2003, at URL <http://www.mercurypolicy.org/new/documents/DentistTheMenace.pdf>.
- Bill Summary, 2002, S.351 bill summary, accessed June 18, 2003, at URL <http://www.mercurypolicy.org/new/documents/S351Summary62502.pdf>.
- Bunce, Nigel, and Hunt, Jim, 2003, Mercury pollution, accessed June 2, 2003, at URL <http://www.physics.uoguelph.ca/summer/scor/articles/scor104.htm>.
- Burton, Dan, 2004, Congressman Dan Burton, accessed February 17, 2004, at URL <http://www.house.gov/burton/dan.htm>.
- Carlton, E.H., 2004, NAACP support of H.R. 1680, accessed February 23, 2004, at URL [http://mercurypoisoned.com/hearings/carlton\\_statement.html](http://mercurypoisoned.com/hearings/carlton_statement.html).
- Chlorine Institute, 2003, Benefits of chlorine, accessed July 11, 2003, at URL <http://www.c12.com/benefits/index.html>.
- Clean Air Network, 2000, Options and strategies for eliminating mercury pollution, accessed June 2, 2003, at URL <http://www.ems.org/mercury/solutions.pdf>.
- Cravotta, C.D., Nimick, D.E., Tewalt, S.T., Eppinger, R.R., and Finger, S.T., 2003, USGS Mine Drainage Interest Group, accessed May 14, 2003, via URL <http://mine-drainage.usgs.gov>.
- D.F. Goldsmith Chemical and Metal Corp., 2003, D.F. Goldsmith Chemical and Metal Corp., accessed June 3, 2003, at URL <http://www.dfgoldsmith.com>.
- District of Columbia Dental Society, 2003, Amalgam waste, accessed May 30, 2003, at URL <http://www.dcdental.org/amalgam.html>.
- Drillbits and Tailings, 2000, Mercury spill poisons villages near the Yanacocha mine in Peru, accessed December 3, 2002, at URL [http://www.moles.org/ProjectUnderground/drillbits/5\\_11/1.html](http://www.moles.org/ProjectUnderground/drillbits/5_11/1.html).
- Faversham House Group, 2003, Chlorine industry should stop using mercury and refrain from passing it on, accessed June 2, 2003, at URL <http://www.edie.net/news/Archive/6695.cfm>.
- Griffin, Greg, 2000, Newmont's Peru gold mine a mixed blessing, accessed June 11, 2003, at URL [http://www.ems.org/banks/yanacocha\\_denver\\_post.pdf](http://www.ems.org/banks/yanacocha_denver_post.pdf).
- Harkinson, Josh, 2003, Illegal gold mining in Ghana shafts locals' health and the environment, accessed March 30, 2004, at URL <http://www.gristmagazine.com/maindish/harkinson062403asp>.
- Hightower, Jane, 2001, Mercury and autism in children, San Francisco Medical Society, accessed May 8, 2003, at URL <http://www.sfms.org/sfm/sfm401b.htm>.
- Hylander, Lars, 2002, European perspectives on management of mercury stockpiles and mercury-bearing waste, accessed March 23, 2004, at URL <http://www.newmoa.org/NEWMOA/htdocs/prevention/mercury/breakingcycle/compendium/Hylander.ppt>.
- Krabbenhoft, David, and Herrmann, Jonathan, 2000, USGS/EPA Mercury Roundtable, accessed June 19, 2003, at URL <http://minerals.usgs.gov/mercury/roundtable.html>.
- Leahy, Patrick, and Snowe, Olympia, 2003, The omnibus Mercury Emissions Reduction Act of 2003, accessed June 1, 2003, at URL <http://leahy.senate.gov/press/200302/022703c.html>.
- Lyon, Jim, 2001, Support for S. 351 letter, accessed June 5, 2003, at URL <http://www.mercurypolicy.org/new/documents/finalcollins/tr.pdf>.
- Mahaffey, K.R., 2004, Methylmercury and epidemiology update, accessed August 8, 2004, at URL <http://www.epa.gov/waterscience/fish/forum/2004/presentations/Monday/mahaffey.pdf>.
- Maine Department of Environmental Protection, 2003, A strategy to reduce mercury content of products, accessed July 9, 2003, at URL <http://www.state.me.us/dep/mercury/productsweb.pdf>.
- Manheim, Frank, 2002, New environmental research publication has timely release for mercury concern in the Gulf of Mexico, accessed June 16, 2004, at URL <http://soundwaves.usgs.gov/2002/06/pubs2.html>.
- Massachusetts Department of Environmental Protection, 2004, Mercury in heating, ventilation, and air conditioning (HVAC) laboratories in voc-tech schools, accessed March 16, 2004, at URL <http://www.newmoa.org/newmoa/htdocs/prevention/mercury/schools/HVAC.pdf>.
- McCaffree, D.R., 2003, Mercury in food as a human health hazard—Resolution 516 (A-03), accessed July 24, 2003, at URL <http://www.ama-assn.org/ama1/upload/mm/annual103/e516a03.doc>.
- McCann, H.G., 1999, Mercury found in Midwest rain, accessed May 30, 2003, at URL [http://healthandenergy.com/mercury\\_in\\_the\\_rain.htm](http://healthandenergy.com/mercury_in_the_rain.htm).
- Medina, Guillermo, 2001, Mitigacion del mercurio en la mineria artesanal del Peru, accessed November 21, 2003, at URL <http://www.gama-peru.org/journada-hg/medina.pdf>.
- Mercury in India, 2003, Usage and releases, accessed June 21, 2004, at URL <http://www.toxiclink.org/backend/images/publications/publications-2-33.pdf>.
- Mercury Instruments, 2003, Mercury vapor monitor on top of Europe's highest volcano, accessed June 2, 2003, at URL [http://www.mercury-instruments.de/NEWS\\_Startseite.html](http://www.mercury-instruments.de/NEWS_Startseite.html).
- Mercury Recyclers, 2002, Mercury—A resource fact sheet, accessed December 10, 2002, at URL <http://pasture.ecn.purdue.edu/~mercury/src/recyclers.htm>.
- Mercury Soap Campaign, 2004, A modern day economic and health scandal, accessed June 7, 2004, at URL [http://www.anamed.net/English\\_Home\\_Page/Mercury\\_Soap\\_Campaign](http://www.anamed.net/English_Home_Page/Mercury_Soap_Campaign).
- Mercury Waste Solutions, 2003, Managing your risk and liability, Mercury Waste Solutions, accessed June 23, 2003, via URL <http://www.mwsi.com>.
- Metal-Pages, 2003a, China's output figures—January-October 2002, accessed January 16, 2003, at URL <http://www.metal-pages.com>.
- Metal-Pages, 2003b, Selenium springs into life as cobalt consolidates on high, accessed June 11, 2003, at URL <http://www.metal-pages.com>.
- Moran, Patrick, and Black, Robert, 2003, Persistent organic pollution and heavy metals in glacial fed lakes and aquatic biota in national parks and forests of Puget Sound Basin, accessed May 3, 2004, at URL <http://www.nps.gov/noca/scidaysabs03.htm>.
- Moravec, B.G., and Smith, S.S., 1997, Little Rock quadrangle, Arkansas, NURE HSSR study, U.S. Geological Survey National Geochemical Database Open-File Report 97-492, accessed April 12, 2004, at URL [http://pubs.usgs.gov/of/1997/ofr-97-0492/quad/q\\_litlrk.htm](http://pubs.usgs.gov/of/1997/ofr-97-0492/quad/q_litlrk.htm).
- Natural Resources Council of Maine, 2003, An act to prevent mercury emissions when recycling and disposing of motor vehicles (LD 1921), accessed July 11, 2003, at URL [http://www.maineenvironment.org/legislature/mercury\\_reduction\\_FS.htm](http://www.maineenvironment.org/legislature/mercury_reduction_FS.htm).
- Neff, Jerry, 2002, Influence of offshore oil and gas platforms on environmental risks of mercury in the Gulf of Mexico, accessed June 16, 2004, at URL <http://www.masgc.org/mercury/abs-neff.html>.
- Nilles, M.A., 2003, NADP—Mercury deposition network, accessed June 2, 2003, at URL [http://bqs.usgs.gov/acidrain/mercury/nadpmdn\\_files/slide0002.htm](http://bqs.usgs.gov/acidrain/mercury/nadpmdn_files/slide0002.htm).
- O'Driscoll, Bill, 2002, Border town would gain from development of mine, Reno Gazette-Journal, accessed June 2, 2003, at URL <http://www.gcu-vse.com/06NewsLetterArticles/border%20town.PDF>.
- Onyx Environmental Services, 2004, Preferred solutions, services and technologies, accessed February 23, 2004, at URL <http://onyxes.com/index.jsp>.
- Pennsylvania Department of Environmental Protection, 2003, Monitoring pollutants in rain, accessed May 30, 2003, at URL <http://www.dep.state.pa.us/dep/deputate/airwaste/aq/acidrain/acidrain/htm>.
- Raloff, Janet, 2003, Mercury retirement—The ultimate solution may be to store the metal, not sell it, Science News Online, accessed August 13, 2003, at URL <http://www.sciencenews.org/20030201/bob8.asp>.
- Rogers, Keith, 2000, EPA to work with state on toxic releases, Las Vegas Review Journal, accessed June 2, 2003, at URL [http://www.reviewjournal.com/lvrj\\_home/2000/May-13-Sat-2000/news/13568588.html](http://www.reviewjournal.com/lvrj_home/2000/May-13-Sat-2000/news/13568588.html).
- Suchanek, T.H., and Marvin-DiPasquale, M., 2003, The influence of natural and anthropogenic processes on mercury cycling in mine-dominated aquatic ecosystems, accessed June 2, 2003, via URL <http://toxics.usgs.gov/meetings/agu2001anthropogenic.html>.
- Summit Valley Equipment and Engineering, Inc., 2004, Client list, retorts, accessed June 4, 2004, at URL <http://www.summit-valley.com/client.html>.
- ToxicAlert, 2003, Nevada superfund sites, accessed June 2, 2003, at URL [http://www.cqs.com/super\\_nv.htm](http://www.cqs.com/super_nv.htm).
- United Nations Environmental Programme, 2002, Global mercury assessment, accessed June 23, 2003, at URL <http://www.chem.unep.ch/mercury/Report/Final%20report/final-assessment-report-25nov02.pdf>.

U.S. Environmental Protection Agency, 2000a, EPA decides mercury emissions from power plants must be reduced, Press Release, accessed May 30, 2003, at URL <http://Yosemite.epa.gov/opa/admpress.nsf>.

U.S. Environmental Protection Agency, 2000b, Regulatory finding on the emissions of hazardous air pollutants from electric utility steam generating units, Federal Register, accessed June 2, 2003, at URL [http://www.epa.gov/ttn/oarp/t3/fr\\_notices/utilfind.pdf](http://www.epa.gov/ttn/oarp/t3/fr_notices/utilfind.pdf).

U.S. Environmental Protection Agency, 2002, National emission standards for hazardous air pollutants—Mercury emissions from mercury cell chlor-alkali plants, Federal Register, accessed June 1, 2004 at URL <http://www.epa.gov/fedrgstr/EPA-AIR/2002/July/Day-03/a15873.htm>.

U.S. Environmental Protection Agency, 2003a, Hospital/medical/infectious waste incinerators, accessed May 30, 2003, at URL <http://www.epa.gov/ttnatw01/129/hmiwi/rihmiwi.html>.

U.S. Environmental Protection Agency, 2003b, Mercury fever thermometers, accessed May 30, 2003, at URL <http://www.epa.gov/glnpo/bnsdocs/hg/thermometers.html>.

U.S. Geological Survey, 2003, Geoenvironmental impacts of mercury and arsenic, accessed June 2, 2003, at URL <http://minerals.usgs.gov/west/projects/hgas.shtml>.

Veiga, M.M., 1997, Mercury in artisanal gold mining in Latin America, accessed May 14, 2003, at URL [http://www.facome.uqam.ca/facome/pdf/viega\\_02.pdf](http://www.facome.uqam.ca/facome/pdf/viega_02.pdf).

## GENERAL SOURCES OF INFORMATION

### U.S. Geological Survey Publications

Historical Statistics for Mineral and Material Commodities in the United States. Open-File Report OF-01-06, 2003.

Materials Flow of Mercury in the Economies of the United States and the World, The. Circular 1197, 2000.

Mercury. Ch. in Mineral Commodity Summaries, annual.

Mercury. Ch. in United States Mineral Resources, Professional Paper 820, 1973.

Mercury in the Environment. Professional Paper 713, 1970.

### Other

Materials Flow of Mercury in the United States, The. U.S. Bureau of Mines Information Circular 9412, 1994.

Mercury Process for Making Chlorine. Euro Chlor, 2003.

Economics of Mercury, The (7th ed.). Roskill Information Services Ltd., 1990.

Sixth Annual Report to EPA. The Chlorine Institute, Inc., 2002.

TABLE 1  
SALIENT MERCURY STATISTICS<sup>1</sup>

(Metric tons unless otherwise specified)

	1999	2000	2001	2002	2003
United States:					
Secondary production, industrial	NA	NA	NA	NA	NA
Imports for consumption	62	103	100	209	46
Exports	181	182	108	201	287
Industry stocks, yearend <sup>2</sup>	NA	NA	NA	NA	NA
Industrial consumption	NA	NA	NA	NA	NA
Price, average per flask, free market <sup>3</sup>	140	140	140	140	140
World, mine production	1,320 <sup>r</sup>	1,360 <sup>r</sup>	1,500 <sup>r</sup>	1,510 <sup>r</sup>	1,530

<sup>r</sup>Revised. NA Not available.

<sup>1</sup>Data are rounded to no more than three significant digits, except prices.

<sup>2</sup>Stocks at consumers and dealers not available. Mine stocks withheld to avoid disclosing company proprietary data.

<sup>3</sup>Source: Platts Metals Week.

TABLE 2  
U.S. IMPORTS AND EXPORTS OF MERCURY BY COUNTRY<sup>1</sup>

Country	2002		2003	
	Quantity, gross weight (metric tons)	Value (thousands)	Quantity, gross weight (metric tons)	Value (thousands)
<b>Imports:</b>				
Australia	107	\$187	--	--
Canada	11	14	6	\$8
Chile	75	53	--	--
Germany	15	614	19	861
India	--	--	1	5
Peru	--	--	19	23
Spain	1	12	--	--
United Kingdom	(2)	9	(2)	18
Total	209	889	46	914
<b>Exports:</b>				
Brazil	2	23	7	50
Canada	4	26	5	40
Ecuador	--	--	2	32
France	8	124	3	42
Germany	21	54	4	26
Guyana	--	--	6	32
Hong Kong	--	--	17	75
India	21	68	25	80
Japan	2	59	--	--
Korea, Republic of	7	34	--	--
Malaysia	--	--	2	19
Mexico	33	237	35	399
Netherlands	73	281	57	270
Peru	--	--	51	185
Singapore	18	62	22	205
United Kingdom	--	--	2	21
Vietnam	--	--	40	172
Other	12	77	9	47
Total	201	1,050 <sup>r</sup>	287	1,690

<sup>r</sup>Revised. -- Zero.

<sup>1</sup>Data are rounded to no more than three significant digits; may not add to totals shown.

<sup>2</sup>Less than 1/2 unit.

Source: U.S. Census Bureau.

TABLE 3  
MERCURY: WORLD MINE PRODUCTION, BY COUNTRY<sup>1,2</sup>

(Metric tons)

Country	1999	2000	2001	2002	2003
Algeria	240	216	321 <sup>r</sup>	307 <sup>r</sup>	300 <sup>c</sup>
China <sup>c</sup>	200	200	190	435 <sup>r</sup>	610
Finland	40	76	71	70	65
Kyrgyzstan	300	257	300	300 <sup>r</sup>	300
Mexico <sup>c</sup>	15	15	15	15	15
Morocco <sup>c</sup>	10	10	10	9	9
Russia <sup>c</sup>	50	50	50	50	50
Spain	433	500	500	300	150
Tajikistan <sup>c</sup>	35	40	40	20	30
Ukraine	NA	NA	--	NA	--
United States <sup>3</sup>	NA	NA	NA	NA	NA
Total	1,320 <sup>r</sup>	1,360 <sup>r</sup>	1,500 <sup>r</sup>	1,510 <sup>r</sup>	1,530

<sup>c</sup>Estimated. <sup>r</sup>Revised. NA Not available. -- Zero.

<sup>1</sup>World totals and estimated data are rounded to no more than three significant digits; may not add to totals shown.

<sup>2</sup>Table includes data available through April 29, 2004.

<sup>3</sup>Data on byproduct mercury are not available.