

HLAMatchmaker: A Molecularly Based Algorithm for Histocompatibility Determination. V. Eplet Matching for HLA-DR, HLA-DQ, and HLA-DP

Rene J. Duquesnoy and Medhat Askar

ABSTRACT: This report describes the design of the eplet version of HLAMatchmaker to determine class II compatibility at the structural level. This matching algorithm is based on the hypothesis, developed from molecular modeling of crystallized antigen-antibody complexes, that functional epitopes are represented by patches of surface-exposed nonself-amino acid residues surrounded by residues within a 3-Å radius. Patch determinations with a molecular viewer of crystalline structural models downloaded from the Entrez Molecular Modeling Database Web site led to the identification of 44 DRB, 33DQB, 29 DQA, 20 DPB, and 9 DPA unique combinations of polymorphic positions. The residue compositions of these patches were then determined from amino acid sequences. This analysis resulted in a repertoire of 146 DRB, 74 DQB, 58 DQA, 45 DPB, and 19 DPA eplets. In many eplets, the residues are in short linear sequences, but many other eplets have discontinuous se-

quences of residues that cluster on or near the molecular surface. This analysis has also shown that all serologically defined DR and DQ antigens detectable by monospecific antibodies have unique eplets. Other eplets are present in groups of class II antigens, many of which appear as cross-reacting. The eplet version of HLAMatchmaker should be considered as a hypothetical model for the structural assessment of donor-recipient compatibility and the determination of mismatch acceptability for sensitized patients. This computer algorithm can be downloaded from the HLA Matchmaker Webside at <http://tpis.upmc.edu>. *Human Immunology* 68, 12–25 (2007). © American Society for Histocompatibility and Immunogenetics, 2007. Published by Elsevier Inc.

KEYWORDS: HLAMatchmaker; HLA; epitope structure; histocompatibility; eplet

ABBREVIATION

HLA human leukocyte antigens

INTRODUCTION

Class II human leukocyte antigens (HLA) play an important role in determining donor-recipient compatibility in solid organ and stem cell transplantation. Class II mismatching elicits strong alloimmune responses that impair transplant success. Preformed donor-specific anti-class II antibodies increase the risk of transplant failure [1–5], and the posttransplant development of anti-class II antibodies is associated with a higher incidence of acute and chronic rejection [6, 7].

Current class II matching strategies in kidney transplantation consider only the serologically defined HLA-DR antigens controlled by the DRB1 locus, although mismatching for HLA-DQ and HLA-DP appears associated with lower graft survivals [8–13] and the development of clinically relevant alloantibodies in transplant recipients [14]. Moreover, molecular typing has revealed a high degree of heterogeneity of HLA-DR antigens. Newer serum screening methods, such as enzyme-linked immunosorbent assay, flow cytometry, and Luminex, have greatly enhanced the detection and specificity analysis of anti-class II antibodies in sensitized patients.

The evaluation of HLA compatibility and the characterization of anti-HLA antibodies require a better understanding of the HLA epitope repertoire. HLAMatchmaker is a structurally based matching program that

From the Division of Transplantation Pathology, Thomas E. Starzl Transplantation Institute, University of Pittsburgh Medical Center, Pittsburgh, PA, USA.

Address reprint requests to: Dr. Rene J. Duquesnoy, Professor of Pathology, Immunology and Surgery, University of Pittsburgh Medical Center, Thomas E. Starzl Biomedical Science Tower, Room W1552, Pittsburgh, PA 15261; Tel: (412) 647-6148; Fax: (412) 647-1755; E-mail: duquesnoyr@msx.upmc.edu.

Received Aug 07, 2006; revised Sep 12, 2006; accepted Oct 02, 2006.

considers each HLA antigen as a string of epitopes represented by short sequences (originally referred to as triplets) involving polymorphic amino acid residues in antibody-accessible positions [15]. HLAMatchmaker determines which triplets are different between donor and recipient, and this algorithm is clinically useful for matching purposes [16–20] and in determining HLA mismatch acceptability for sensitized patients [21–31]. Although many triplets correspond to serologically defined private and public epitopes, they provide an incomplete description of the HLA epitope repertoire.

A recent study has led to a new version of HLAMatchmaker that considers the hypothesis, developed from molecular modeling of crystallized antigen-antibody complexes, that functional epitopes are represented by patches of surface-exposed nonself-amino acid residues surrounded by residues within a 3-Å radius [32]. These patches are referred to as “eplets,” and many of them are short linear sequences common to triplets, but others have residues in discontinuous sequence positions that cluster together on the molecular surface. The eplet version of HLAMatchmaker therefore considers a more complete repertoire of structurally defined epitopes.

This report describes how eplets are assigned in determining HLA-DR, -DQ, and -DP compatibility at the humoral immune level. This analysis considers 4-digit alleles encoded by not only DRB1 and DQB1, but also DRB3, DRB4, DRB5, DQA1, DPA1, and DPB1, because all of them have antigenic determinants that can induce specific antibodies. This paper will show how eplets correspond to serologically defined class II antigens and how the eplet version of HLAMatchmaker can be used to determine structurally based class II compatibility at the humoral immune level.

METHODS AND RESULTS

Topography of Polymorphic Amino Acid Residues

The construction of the eplet repertoire is based on polymorphic amino acid residues on the HLA molecular surface. Their locations are easily determined with three-dimensional models of class II molecules. The Entrez Molecular Modeling Database of the National Center for Biotechnology Information stores on its Web site (<http://www.ncbi.nlm.nih.gov/Structure>) an extensive collection of crystallographic structures of HLA molecules that can be viewed with the Cn3D structure and sequence alignment software program [33, 34].

Five DRB structures have been downloaded from the Molecular Modeling Database Web site, all of them have the same monomorphic DRA1*0101 sequence. They are DRB1*0101 (PDB Number 1KG0) [35], DRB1*0301 (1A6A) [36], DRB1*0401 (1D6E) [37], DRB1*1501 (1BX2) [38], and DRB5*0101 (1H15)




FIGURE 1 Visualization of polymorphic amino residue positions on HLA-DRB and HLA-DQ molecules. The following crystalline models are shown: DRA1*0101, DRB1*0101 (PDB # 1KG0), and DQA1*0301, DQB1*0302 (PDB No. 1JK8). Left = top view; middle = β -chain side view right = α -chain side view.

[39]. There are three models of HLA-DQ heterodimers: DQA1*0501, DQB1*0202 (1S9) [40], DQA1*0301, DQB1*0302 (1JK8) [41], and DQA1*0102, DQB1*0602 (1UVQ) [42].

Cn3D viewing of crystalline HLA-DR and HLA-DQ molecular models shows different patterns of surface expression of polymorphic residues (Figure 1). The structural polymorphisms of HLA-DR are restricted to the β chains. They are readily visible on the top of the molecule adjacent to the bound peptide, and many of them involve contiguous sequences. Polymorphic residues on the side of the molecule generally comprise distinct clusters in both $\beta 1$ and $\beta 2$ domains. A few polymorphisms are visible at the bottom part of the molecule that is nearby the cell membrane. DRB and DQB seem to show similar numbers of polymorphic positions. DQA displays somewhat contiguous polymorphic positions on the top of the molecule near the bound peptide and on the side of the $\alpha 1$ domain. The polymorphic positions in the $\alpha 2$ domain seem to be more in distinct clusters.

No crystalline structures of HLA-DP molecules are available in the Molecular Modeling Database. Predicted three-dimensional models for HLA-DP and other DR and DQ class II alleles can be generated by submitting amino acid sequences to the Geno3D online molecular modeling server [43]. This web-based service is located at <http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>. The Vector NTI-3D Molecular Viewer software (Invitro-

gen Life Science Software, Frederick, MD) can visualize the locations of polymorphic amino acid residues and also measure intermolecular distances. These prediction models yield incomplete information about molecular surface expression of polymorphic residues because they deal only with isolated α or β chains without any peptide in the peptide-binding groove.

Determination of Patches on HLA Class II Molecules

Each surface-exposed polymorphic residue represents an essential element of a potential epitope recognized by antibody. The “select by distance” command of the Cn3D molecular viewer has been used to identify residues within a 3-Å radius. A patch defined this way seems to provide the best estimate of the size of a functional epitope [32].

Sequence comparisons of 381 most common DRB1, 3, 4, 5 alleles have identified 49 polymorphic positions on the molecular surface. Cn3D viewing has shown that 13 of them are on the top (*i.e.*, the α -helices) and 26 are on the side of the molecule (Table 1). Six positions have “underside” locations (*i.e.*, underneath the groove) and four are at the “bottom” near the cell membrane; they become more readily visible if the molecule has been turned upside down. These positions seem less antibody-accessible if the HLA antigen is anchored in the cell membrane like in the lymphocytotoxicity test, but they might react with antibody if the HLA molecule is fixed to a different surface like in a solubilized antigen-binding assay. Molecular surface expression of polymorphic residues has been graded as prominent (++), readily visible (+), and somewhat visible (\pm).

Table 1 lists the sequence positions clustered within a 3-Å radius of each exposed polymorphic position. These patches are combinations of monomorphic and polymorphic positions (marked in bold, underlined font), and they have an average of 4.2 residues. About one third are continuous sequences, and the remaining are discontinuous sequences including several patches (*e.g.*, positions 12, 14, and 16) with residues far removed in sequence. Three patches in positions 6, 34, and 57 have monomorphic DRA residues (prefixed with A), and five α helix patches (positions 57, 70, 74, 77, and 81) include residues of peptides bound to the groove; their positions have the prefix “P.” Exposed peptide residues might contribute to the functional epitope recognized by alloantibody. Several studies have shown the influence of HLA-bound peptides on the reactivity of class I and class II specific antibodies [44–47].

Sequence comparisons of 43 DQB1 alleles and Cn3D viewing of DQ molecules have identified patches for 36 polymorphic positions on the DQB chain surface (Table

2), 13 of them are on the top and 18 are on the side of the DQB molecule. There are three underside and two bottom locations. DQB patches have fewer residues than DRB patches (3.8 vs 4.2, $p = 0.04$ by two-tailed Student’s *t*-test), and one half of them are continuous sequences. Only one patch (in position 30) has a peptide residue and another patch (position 53) has a monomorphic α -chain residue.

The 21 DQA1 alleles have almost the same number of polymorphic surface positions as the 43 DQB1 alleles, namely 37 including 12 on the top and 20 on the side of the molecule (Table 3). DQA patches have an average of 4.1 residues. One contains a peptide residue and two have monomorphic β -chain residues.

The determination of patches on DP molecules was more difficult because no crystalline structures are available for DP, and the Vector NTI-3D Molecular Viewer can visualize predicted structures of only single DP chains rather the whole molecule with a peptide inserted. Therefore, we included Cn3D viewing of DR and DQ molecules and considered positions equivalent to the polymorphic positions on DP sequences. These positions were identified with the Basic Local Alignment Search Tool (BLAST) program [48] available on the National Center for Biotechnology Information Web site <http://www.ncbi.nlm.nih.gov/BLAST>.

This patch analysis was limited to the first domain (positions 1–91) of 95 DPB alleles. Insufficient sequence information of the second DPB domain precluded any patch determination. The number of patches is lower for DPB than for DRB and DQB: 19 versus 32 and 26 patches in the first domains of DRB and DQB, respectively (Table 4). The DPB patches have an average of 3.9 residues. DPA1 alleles ($n = 13$) have fewer patches than DQA1, 11 versus 37. DPA patches have an average of 4.4 residues (Table 4).

These findings on HLA-DR, DQ, DP patches are comparable in size and location to those reported for HLA-A, B, C patches [32]. The polymorphic positions determine residue variability within each patch. Class II patches have between 1 and 4 polymorphic positions. A few patches have the same polymorphic positions although there are differences between the monomorphic positions. For instance, the DRB patches in positions 85 and 86 (see Table 1) have the same two polymorphic positions but different monomorphic positions. Such patches are considered equivalent.

The patch information in Tables 1–4 yielded the following numbers of unique combinations of polymorphic positions: DRB: 46, DQB: 33, DQA: 29, DPB: 20, and DPA: 9; they are considered the positional basis for the class II HLA epitope repertoire. The residue compositions of these polymorphic patches were determined from amino acid sequences retrieved from the IMGT/

TABLE 1 Polymorphic and monomorphic residue positions in three-Angstrom patches on HLA-DRB1, -DRB3, -DRB4 and -DRB5 alleles*

Class II Locus	Sequence position	Molecular location	Surface exposure	3.0 Angstrom patches													
DRB	4	Side	++	3	<u>4</u>	5											
DRB	6	Side	+	5	<u>6</u>	7	A15	A17									
DRB	12	Underside	±	11	12	13	29										
DRB	14	Underside	+	13	14	15	16	27	29								
DRB	16	Underside	+	15	16	17	18	25									
DRB	18	Side	+	17	18	19	23										
DRB	25	Side	++	16	24	25	26	43									
DRB	26	Side	±	25	26	27	42										
DRB	31	Underside	+	10	29	30	31	32									
DRB	32	Underside	+	31	32	33	35										
DRB	33	Underside	+	8	32	33	34										
DRB	34	Side	++	33	34	35	A83										
DRB	40	Side	+	28	39	40	41										
DRB	41	Side	+	40	41	42	43	44	45								
DRB	44	Side	+	41	43	44	45										
DRB	47	Side	+	28	46	47	48	62									
DRB	48	Side	++	47	48	49											
DRB	51	Side	+	37	50	51	52										
DRB	57	Top	+	56	57	58	61	A76	P13								
DRB	58	Top	+	54	57	58	59	62									
DRB	59	Top	++	58	59	60											
DRB	60	Top	++	59	60	61	63										
DRB	67	Top	+	66	67	68	71										
DRB	70	Top	+	67	69	70	71	73	P11								
DRB	71	Top	±	70	71	72	73										
DRB	73	Top	+	69	72	73	74	76	77								
DRB	74	Top	+	70	71	72	73	74	79	P8	P9						
DRB	76	Top	++	73	75	76	77										
DRB	77	Top	++	73	76	77	78	P6									
DRB	81	Top	+	80	81	82	85	P4									
DRB	85	Top	+	84	85	86											
DRB	86	Side	±	82	85	86	87	90									
DRB	96	Side	++	95	96	97	180										
DRB	98	Side	++	97	98	99	120										
DRB	104	Side	+	103	104	105	107	114									
DRB	105	Side	++	104	105	106	107										
DRB	108	Side	++	107	108	109											
DRB	112	Bottom	++	108	111	112	113										
DRB	120	Side	+	98	119	120	121										
DRB	133	Bottom	++	132	133	134											
DRB	135	Bottom	++	134	135	136											
DRB	140	Side	++	139	140	141											
DRB	142	Side	++	138	141	142	143										
DRB	149	Side	++	148	149	150											
DRB	180	Side	+	96	177	179	180	181									
DRB	181	Side	++	180	181	182											
DRB	183	Side	+	182	183	184											
DRB	187	Side	++	186	187	188											
DRB	189	Bottom	++	188	189	190											

* Polymorphic positions are marked in bold, underlined font.

TABLE 2 Polymorphic and monomorphic residue positions in three-Angstrom patches on HLA-DQB1 alleles*

Class II Locus	Sequence position	Molecular location	Surface exposure	3.0 Angstrom patches					
DQB	3	Side	++	<u>3</u>	4				
DQB	14	Underside	±	<u>13</u>	<u>14</u>	15	27		
DQB	23	Side	++	18	22	<u>23</u>	24		
DQB	26	Underside	+	25	<u>26</u>	27	42		
DQB	30	Side	±	29	<u>30</u>	31	<u>37</u>	<u>38</u>	P9
DQB	45	Side	++	41	44	<u>45</u>	<u>46</u>	72	
DQB	46	Side	++	<u>45</u>	<u>46</u>	<u>47</u>			
DQB	47	Side	+	<u>46</u>	<u>47</u>	48	62		
DQB	49	Side	++	48	<u>49</u>	50			
DQB	52	Side	++	51	<u>52</u>	<u>53</u>	54		
DQB	53	Top	++	<u>52</u>	<u>53</u>	54	A78		
DQB	55	Top	++	<u>52</u>	54	<u>55</u>	<u>56</u>		
DQB	56	Top	++	<u>55</u>	<u>56</u>	<u>57</u>			
DQB	57	Top	±	<u>56</u>	<u>57</u>	58			
DQB	66	Top	++	65	<u>66</u>	<u>67</u>			
DQB	67	Top	+	64	<u>66</u>	<u>67</u>	68	<u>71</u>	
DQB	70	Top	++	69	<u>70</u>	<u>71</u>			
DQB	71	Top	+	<u>67</u>	<u>70</u>	<u>71</u>	72		
DQB	74	Top	±	73	<u>74</u>	<u>75</u>			
DQB	75	Top	+	<u>74</u>	<u>75</u>	76	80		
DQB	77	Top	++	73	<u>76</u>	<u>77</u>	78	81	
DQB	84	Top	+	83	<u>84</u>	<u>85</u>			
DQB	85	Top	++	<u>84</u>	<u>85</u>	<u>86</u>	<u>89</u>		
DQB	87	Side	+	83	<u>86</u>	<u>87</u>	88	92	
DQB	89	Side	+	<u>85</u>	<u>86</u>	88	<u>89</u>	<u>90</u>	92
DQB	90	Underside	±	<u>86</u>	<u>89</u>	<u>90</u>	91		
DQB	116	Side	+	102	115	<u>116</u>	117		
DQB	125	Side	+	124	<u>125</u>	<u>126</u>			
DQB	126	Side	++	124	<u>125</u>	<u>126</u>	127		
DQB	130	Side	++	129	<u>130</u>	131	174		
DQB	135	Side	++	134	<u>135</u>	136			
DQB	140	Side	++	139	<u>140</u>	141			
DQB	167	Bottom	++	166	<u>167</u>	<u>168</u>	190		
DQB	168	Bottom	+	<u>167</u>	<u>168</u>	169			
DQB	182	Side	++	181	<u>182</u>	183			
DQB	185	Side	++	184	<u>185</u>	186			

* Polymorphic positions are marked in bold, underlined font.

HLA online database [49] with a Microsoft Excel macro (called HLA Patch Generator) developed by Grzegorz Dudek (Czestchowa University of Technology, Poland).

Assignments of Class II Eplets

DRB. An analysis of the most common four-digit DRB alleles (122 DRB1 and 18 DRB3, 4, 5 alleles) has yielded a total of 222 patches with different combinations of polymorphic residues, 91 of them are on the top of the molecule, and most of them have overlapping residues. The remaining 132 polymorphic patches are largely on the side of the molecule and include 58 at underside or bottom locations.

Further comparisons have shown that certain overlapping polymorphic patches can be grouped together because they belong to a single allele or a distinct group of alleles including cross-reacting antigens. The term eplet

is used to describe a given patch or an overlapping group of patches.

For example, DRB1*1501, DRB1*1502, DRB1*1503 and DRB5*0202 share the same three overlapping patches: 70IQAA, 71QAA, and 74QAAA. They are collectively referred to as one eplet assigned as 71QAA. In other cases, an eplet represents a single polymorphic patch. For instance, the 98QS patch shared between DRB3*0101, DRB3*0201, DRB3*0202, and DRB3*0301 has been assigned as the 98QS eplet. This eplet corresponds to the serologically defined DR52 determinant.

Several polymorphic patches are shared between either all DRB1 alleles or all DRB3, 4, 5 alleles, and they cannot be considered immunogenic. For example, there are two polymorphic patches in position 108. One is 100T found on DRB5 molecules, and specific antibody-

TABLE 3 Polymorphic and monomorphic residue positions in three-Angstrom patches on HLA-DQA1 alleles*

Class II Locus	Sequence position	Molecular location	Surface exposure	3.0 Angstrom patches							
DQA	2	Side	++	1	<u>2</u>	3					
DQA	18	Side	++	17	<u>18</u>	19	B7				
DQA	21	Side	+	20	<u>21</u>	22					
DQA	25	Side	±	14	24	<u>25</u>	<u>26</u>	27	37	P5	
DQA	40	Side	+	39	<u>40</u>	<u>41</u>					
DQA	41	Side	++	38	<u>40</u>	<u>41</u>	42				
DQA	44	Side	++	43	<u>44</u>	<u>45</u>					
DQA	45	Side	+	36	<u>44</u>	<u>45</u>	46				
DQA	47	Side	+	<u>34</u>	46	<u>47</u>	<u>48</u>				
DQA	48	Side	+	47	<u>48</u>	49	<u>51</u>				
DQA	50	Side	++	49	<u>50</u>	<u>51</u>					
DQA	51	Underside	+	<u>48</u>	<u>50</u>	<u>51</u>	<u>52</u>	<u>53</u>			
DQA	52	Side	+	49	<u>51</u>	<u>52</u>	<u>53</u>				
DQA	53	Top	++	<u>51</u>	<u>52</u>	<u>53</u>	<u>54</u>				
DQA	54	Top	++	<u>51</u>	<u>53</u>	<u>54</u>	<u>55</u>				
DQA	55	Top	+	<u>54</u>	<u>55</u>	<u>56</u>					
DQA	56	Top	++	<u>53</u>	<u>55</u>	<u>56</u>	57				
DQA	59	Top	+	58	<u>59</u>	60	<u>61</u>	62			
DQA	61	Top	++	58	<u>59</u>	60	<u>61</u>	62	65		
DQA	64	Top	++	60	63	<u>64</u>	65				
DQA	66	Top	+	62	65	<u>66</u>	67	70			
DQA	69	Top	±	65	68	<u>69</u>	70				
DQA	75	Top	++	74	<u>75</u>	<u>76</u>	<u>79</u>				
DQA	76	Top	±	72	<u>75</u>	<u>76</u>	77	80			
DQA	79	Top	++	<u>75</u>	78	<u>79</u>	<u>80</u>	B32			
DQA	80	Side	±	<u>75</u>	<u>79</u>	<u>80</u>	81				
DQA	107	Side	+	97	106	<u>107</u>	108				
DQA	129	Side	++	126	128	<u>129</u>	<u>130</u>				
DQA	130	Side	+	<u>129</u>	<u>130</u>	131					
DQA	138	Side	+	137	<u>138</u>	<u>139</u>	141				
DQA	139	Side	+	33	<u>138</u>	<u>139</u>	140	141			
DQA	153	Side	+	136	152	<u>153</u>	154	B150			
DQA	156	Bottom	+	106	155	<u>156</u>	157				
DQA	160	Bottom	+	159	<u>160</u>	<u>161</u>					
DQA	161	Bottom	++	<u>160</u>	<u>161</u>	162					
DQA	163	Bottom	+	127	162	<u>163</u>	164				
DQA	175	Side	++	174	<u>175</u>	176					

* Polymorphic positions are marked in bold, underlined font.

ies would react with the serologically defined DR51, which corresponds to DRB5. The other one is 108P present on all DRB1, DRB3, and DRB4 molecules. This patch cannot induce alloantibodies because it is always a self-sequence. The patches in position 34 represent another example: 34HQ is shared between all DR4 alleles and 34NQ is shared between all DRB3, DRB4, and DRB5 alleles plus all DRB1 alleles that are not DR4. Thus, 34NQ must be considered as a self-sequence and cannot be immunogenic. This analysis has yielded 17 self-DRB patches, and they have been deleted from the eplet repertoire.

This analysis yielded a total of 146 DRB eplets, 52 of them are on the α helices on the top of the molecule. There are 59 eplets on the side surface, including 8 at the bottom. A total of 38 eplets are beneath the pep-

tide-binding groove and they cluster in two underside locations, namely positions 12, 14, and 16 ($n = 19$) and positions 31, 32, and 33 ($n = 19$). Eplets in bottom and underside locations seem less antibody accessible if the HLA molecule is bound to the cell membrane.

Table 5 shows serologically defined DR antigens that have one or more corresponding eplets. These antigens can be readily identified with monospecific allosera and/or monoclonal antibodies as demonstrated during the 1984–1997 International Histocompatibility Workshops [50–54]. Seven serologically defined DR antigens, namely DR5, DR6, DR13, DR14, DR15, DR16, DR17, and DR18, do not have corresponding eplets. None of them except DR15 and DR17 can be identified with monospecific antibodies; their serologic determination is based on reactivity patterns of antibodies specific for

TABLE 4 Polymorphic and monomorphic residue positions in three-Angstrom patches on HLA-DPB1 and HLA-DPA1 alleles*

Class II Locus	Sequence position	Molecular location	Surface exposure	3.0 Angstrom patches						
DPB	8	Underside	±	7	<u>8</u>	9	31			
DPB	11	Side	±	10	<u>11</u>	12	A11			
DPB	28	Side	±	26	27	<u>28</u>	29	36	P8	
DPB	33	Underside	++	30	32	<u>33</u>	34			
DPB	35	Underside	+	7	34	<u>35</u>	<u>36</u>	48		
DPB	36	Underside	±	28	<u>35</u>	<u>36</u>	37			
DPB	43	Side	+	39	42	<u>43</u>	44	<u>70</u>		
DPB	56	Side	+	52	55	<u>56</u>	<u>57</u>	60		
DPB	57	Side	++	<u>56</u>	<u>57</u>	58				
DPB	64	Top	+	63	<u>64</u>	<u>65</u>				
DPB	65	Top	+	<u>64</u>	<u>65</u>	66	<u>69</u>			
DPB	69	Top	+	<u>65</u>	68	<u>69</u>	<u>70</u>			
DPB	70	Top	++	<u>43</u>	66	<u>69</u>	<u>70</u>	71		
DPB	76	Top	+	72	75	<u>76</u>	77			
DPB	84	Side	+	83	<u>84</u>	<u>85</u>	88			
DPB	85	Side	++	81	<u>84</u>	<u>85</u>	<u>86</u>			
DPB	86	Side	+	<u>85</u>	<u>86</u>	<u>87</u>				
DPB	87	Side	++	<u>86</u>	<u>87</u>	88				
DPB	91	Side	++	90	<u>91</u>	92				
DPA	18	Side	+	14	16	17	<u>18</u>	67		
DPA	28	Underside	+	25	26	27	<u>28</u>	29		
DPA	31	Underside	+	20	30	<u>31</u>	32			
DPA	50	Top	++	49	<u>50</u>	<u>51</u>				
DPA	51	Top	+	47	<u>50</u>	<u>51</u>	52			
DPA	72	Top	+	68	71	<u>71</u>	<u>73</u>	75	76	
DPA	73	Top	±	<u>72</u>	<u>73</u>	74	75	76	77	
DPA	83	Side	+	82	<u>83</u>	84	B37			
DPA	111	Side	++	86	110	<u>111</u>	112			
DPA	127	Side	+	121	126	<u>127</u>	128			
DPA	160	Bottom	+	124	159	<u>160</u>	161			

* Polymorphic positions are marked in bold, underlined font.

epitopes shared between different groups of antigens. For instance, serologic assignments of the serologic DR6 splits DR13 and DR14 can be deduced from antibodies reactive with different groups, such as DR2+6, DR3+6, DR3+5+6, DR5+13, DR3+8+13, and antigens, such as DR8 and DR11 [55]. The reaction patterns of these antibodies are often too complex for reliable serologic assignments of individual DR13 and DR14 alleles.

The DR2 split DR15 and the DR3 split DR17 can be defined serologically by monospecific antibodies. It is possible that the DR15 and DR17 antigens correspond to a pair of eplets whereby one is the specific recognition site and the other functions as a critical contact site for antibody. As described elsewhere [32, 56], certain HLA antibodies react with epitopes represented by combinations of two eplets separated 6–15 Å from each other. For DR17, the most likely pair of eplets appears to be 73GRDN, present on all DR3 molecules, and 48FR, present on DR17, DR11, DR13, and DR15. These eplets are about 9.5 Å apart. Similarly, the DR15 spec-

ificity might be represented by the combination of 71QAA (on DR15 and DRB5*0202) with 40DFD (on DR4, DR8, DR11, DR13, DR14, DR15, DR16, DR17, and DRB3*01) or 48FR (on DR17, DR11, DR13, and DR15).

Table 5 also lists unique eplets that correspond to the serologically defined antigens encoded by the DRB3, DRB4, and DRB5 loci. DR53 has a relatively large number of unique eplets in nonoverlapping positions. There are also unique eplets that distinguish DRB3*01 from DRB3*02 alleles. These eplets appear to correspond to the DR52 subtypes serologically defined as DR52a and DR52b [57, 58]. Similarly, the 31QDIY and 6C eplets can differentiate between DRB5*01 and DRB5*02.

DQB1. An HLA Patch Generator analysis of 43 *DQB1* alleles has yielded 106 *DQB1* patches with different combinations of polymorphic residues, 48 of them are on the top and 46 are on the side of the molecule; 12 are in underside or bottom locations.

TABLE 5 Serologically defined HLA-DR antigens with uniquely corresponding eplets

DR Antigen	Unique eplets						
DRB1							
DR1	12LKF	31QCIY					
DR2	12PKR	133L	142M				
DR3	73GRDN						
DR4	12VKH	34HQ	96YL	180LT			
DR7	14YKH	25HQF	31QLFY	71DRE			
DR8	25YRF	73ALDT					
DR9	12DKF	70FRRA					
DR10	12VKF	31ERVH	40EYD	74RRAA			
DR11	57DE						
DR12	25YRL	31YHFH	47EFR				
DRB3,4,5							
DRB3 (DR52)	98QS						
DRB3*01 (DR52a)	12RKS	183A					
DRB3*02 (DR52b)	31LHFH	51R					
DRB4 (DR53)	25HWN	44NL	48YQ	81YV	96QM	180MM	187Q
DRB5 (DR51)	12DKY	104AR	108T				
DRB5*01 (DR51a)	31QDIY						
DRB5*02 (DR51b)	6C						

As shown for DRB, we have assigned eplets from groups of overlapping DQB1 patches shared between combinations of DQB1 alleles. For instance, the 55RPL eplet represents the combination of overlapping 55RPL, 56 RLD, and 57LD patches on DQB1*0401 and DQB1*0402.

This analysis has resulted in a repertoire of 74 distinct DQB1 eplets. Table 6 shows which eplets correspond to DQ1-DQ7 specificities; all of them can be defined serologically with monospecific antibodies. The DQ3 subtype DQ8 has a unique 56PPA eplet that is structurally similar to the 55PPP eplets that correspond to DQ3. There is no unique eplet for the DQ3 subtype DQ9, and this is consistent with the experience that there are no monospecific antibodies against DQ9 [54].

DQA1. Complete sequence information (positions 6–180) is available for 18 DQA1 alleles. HLA Patch Generator identified 91 DQA patches with different polymorphic residue combinations, 38 are on the top and 41 are on the side of the molecule. There are 12 polymorphic patches on underside and bottom locations. Similar to DRB and DQB, eplets have been assigned to describe single polymorphic patches and overlapping patches that belong to the same group of DQA1 alleles. Altogether, there are 58 distinct DQA1 eplets.

Although DQA1 polymorphisms have never been defined serologically, several two-digit DQA1 alleles have multiple unique eplets (Table 6). Especially, DQA1*01 can be distinguished with seven nonoverlapping eplets. No unique eplets have been identified for DQA1*04 and DQA1*06.

DPA and DPB. An HLA Patch Generator analysis of DPB1*01 through DPB1*99 has generated 62 polymorphic patches, 31 of them are on the top and 23 are on the side of the DPB1 chain. Further comparisons of overlapping patches led to the assignment of a total of 45 DPB eplets, 15 of which are on one or few DPB1 alleles all of which appear to have low frequencies. The identification of DPB eplets is confined to the first domain (positions 8–91), because insufficient sequence information is available for the second domain of DPB chains.

The 13 DPA1 alleles have only nine polymorphic positions, and an HLA Patch Generator analysis has led to the assignment of 19 eplets, 5 of which are on the top

TABLE 6 Serologically defined DQ antigens and 2-digit DQA1 alleles with their corresponding eplets

DQ locus	Unique eplets					
DQ Antigen						
DQ1	52PQ					
DQ5	30HYV	71VGA	87AY	116I		
DQ6	87AF	125GQ				
DQ2	30SIV	45GE	52LL	70RK		
DQ3	55PPP					
DQ7	45EV					
DQ8	56PPA					
DQ4	55PRL					
DQA1 Allele						
DQA1*01	18F	48WF	56KGG	69A	76IM	175Q
DQA1*02	47EKL	52FHR				
DQA1*03	25YS	47EQL	52FRR	75IVR		
DQA1*05	76SL	107I	156L	163S	175K	

and 10 are on the side of the DPA chain. DQA1*02 has a unique 111R, and DQA1*04 has two unique eplets, 18T and 73IA. There are no unique eplets for DQA1*01 and DQA1*03.

Although no DPA1 or DPB1 allele-specific antibodies have been identified [59], there are several murine and human monoclonal antibodies that recognize DP epitopes that appear to eplets, such as 31M and 51RA on DPA1, and 56ED and 85DEA on DPB chains [59].

Example of Eplet-Based Determination of Class II Compatibility

Table 7 represents an illustration of eplet mismatches for HLA-DR and HLA-DQ. This analysis was done for the phenotype DRB1*1101, 1501; DRB3*0202; DRB5*0101; DQA1*0102, 0501; DQB1*0301, 0602, which is serologically equivalent to DR11,15; DR51,52; DQ6,7. The incompatible DR antigens in the left columns of Table 7 show varying numbers of mismatched eplets. The highest range of 9–12 eplet mismatches occurs with DR7, DR9, DR10, DR12, and DR14. The lowest range of 1–3 eplet mismatches is seen for DR3, DR13, and DR16. These low numbers are not surprising because DR3 and DR13 cross-react with DR11 since they share several epitopes that correspond to eplets. Moreover, DR15 and DR16 are serologic splits of DR2, whereas DRB1*1601 and DRB1*1602 have one and three mismatched eplets, respectively. It should be noted however, that DR12—which together with DR11 is a split of DR5—has nine mismatched eplets.

As expected, all eplets that are unique for the antigens listed in Table 7 are mismatches for this phenotype. It may also be seen that many mismatched eplets have overlapping sequences. For instance, DR1 has this pair of overlapping eplets: 12LKF, which is unique on DR1 and 14FEH, which is present on DR1, DR9, and DR10. They may represent distinct epitopes because specifically reactive antibodies have been identified [54]. On the other hand, it is also possible that the combination of 12LKF and 14FEH comprises a single structural determinant that could induce an antibody that reacts with only DR1.

The right columns of Table 7 list examples of eplet mismatches for class II alleles with the same two-digit types as in the above phenotype. DRB1*1104 and DRB1*1502 are zero-eplet mismatches. The other DRB1*11 and DRB1*15 alleles have one or two—but not necessarily the same—mismatched eplets.

Antigens controlled by the DRB3, 4, 5 loci have varying numbers of mismatched eplets. DRB4*0401 (DR53) has the most ($n = 16$), and this is partially due to the large number of eplets unique to DR53 (Table 5). It should be noted that the other two-digit DRB3 and DRB5 alleles have mismatched eplets. Especially

DRB3*0101, which corresponds to the serologically defined DR52a specificity [57, 58], has seven mismatched eplets, including two that are unique for this allele. These eplet differences are clinically relevant because our experience has shown several cases whereby a DRB3*0202- or DR52b-positive patient makes antibodies reactive with DRB3*0101 or DR52a (unpublished data). Good matches can be present for DRB3 and DRB5 alleles with the same two-digit types. For instance, DRB3*0201, DRB3*0203, and DRB5*0105 are zero-eplet mismatches, and others have one or two mismatched eplets.

DQ antigens seem structurally more mismatched than DRB antigens because they have two polymorphic chains: DQB antigens have 7–10 mismatched eplets and DQA alleles have 4–7 mismatched eplets (Table 7). Although DQ5 (DQB1*0501) and DQ6 (DQB1*0602) are serologic splits of DQ1, there is a high degree of structural incompatibility of DQ5 as indicated by 10 mismatched eplets. The phenotype of this example contains the DQ3 split DQ7 (DQB1*0301). With one mismatched eplet, the DQ3 split DQ9 (DQB1*0303) mismatches seems structurally the most compatible among the DQB1*03 mismatches. DQB1*0603, DQA1*0101, DQA1*0502, and DQA1*0503 are zero or one eplet mismatches.

DISCUSSION

This report describes the design of the eplet-based class II version of HLAMatchmaker and how this algorithm can be used to determine structural compatibility for antigens encoded by HLA-DR, HLA-DQ, and HLA-DP. For each of these loci, an eplet repertoire has been developed from patches of residues within a 3-Å radius of each polymorphic residue exposed on the molecular surface. In many eplets, the residues are in short linear sequences, but many other eplets have discontinuous sequences of residues that cluster on or near the molecular surface. This analysis has identified eplets that correspond to serologically defined DR and DQ antigens recognized by monospecific antibodies. Other eplets are present on groups of class II antigens, many of which are considered cross-reacting.

The class II version of HLAMatchmaker considers all polymorphisms in the HLA-D regions that lead to alloantibody responses, and they should be defined by DNA-typing methods, preferably at the four-digit allelic level. Most clinical laboratories type for DRB and DQB antigens and other class II alleles are not considered. Interestingly, DQA1 alleles can be putatively assigned to a given phenotype on the basis of common DRB1-DQB1-DQA1 haplotypes reported in various ethnic groups [60–62]. HLA-DP typing is almost never done

TABLE 7 Example of an HLAMatchmaker Analysis to Determine HLA-DR and HLA-DQ Compatibility at the Eplet Level

Mismatched Antigen	Number of mm Eplets	Mismatched Eplets	Same 2-Digit Allele	Number of mm Eplets	Mismatched Eplets
DRB1*0101 (DR1)	8	12LKF, 14FEH, 25HRL, 26RL, 31QCIY, 67LR, 71QRA, 74QRAA	DRB1*1102	2	67IE, 71DEA
DRB1*0301 (DR3)	3	25HRY, 31YYFH, 73GRDN	DRB1*1103	2	67IE, 71DEA
DRB1*0401 (DR4)	6	12VKH, 34HQ, 47DYR, 71QKA, 96YL, 180LT	DRB1*1104	0	
DRB1*0701 (DR7)	9	4Q, 14YKH, 25HGF, 31QLFY, 59AES, 67IR, 71DRG, 98ES, 180VM	DRB1*1105	2	12STG, 14GEH
DRB1*0802 (DR8)	5	12STG, 14GEY, 25YRF, 47DYR, 73ALDT	DRB1*1106	1	81HA
DRB1*0901 (DR9)	12	4Q, 12DKF, 14FEH, 25HRY, 31QGIY, 59AES, 70FRA, 71RRA, 73AEDT, 74RRAE, 98ES, 180VM			
DRB1*1001 (DR10)	11	12VKF, 14FEH, 25HRL, 26RL, 31ERVH, 40EYD, 67LR, 70LRRA, 71RRA, 74RRAA, 180VM	DRB1*1502	0	
DRB1*1201 (DR12)	10	12STG, 14GEY, 25YRL, 26RL, 31YHFH, 47EFR, 59AES, 67IR, 70IDRA, 81HA	DRB1*1503	1	31QHFY
DRB1*1301 (DR13)	3	31YYFH, 67IE, 71DEA	DRB1*1504	1	67FA
DRB1*1401 (DR14)	9	31 YYFH, 47DYR, 57AA, 67LR, 70LRRA, 71RRA, 73AEDT, 74RRAE, 112Y			
DRB1*1601 (DR16)	1	47DYR			
DRB1*1602 (DR16)	3	47DYR, 67LR, 70LDRA			
DRB3*0101 (DR52a)	7	12RKS, 25HRY, 31LYFH, 47DYR, 59AES, 73GRDN, 183A	DRB3*0201	0	
DRB3*0301 (DR52c)	2	31LYFH, 59AES	DRB3*0203	0	
DRB4*0101 (DR53)	16	4Q, 16HLW, 25HWN, 31QYIY, 40IYN, 44NL, 48YQ, 67LR, 70LRRA, 71RRA, 73AEDT, 74RRAE, 81YV, 96QM, 180MM, 187Q	DRB3*0204	1	73GRDN
DRB5*0202 (DR51b)	3	6C, 31QGIY, 81HA	DRB5*0102	1	31QGIY
			DRB5*0103	2	31QGIY, 67FT
			DRB5*0104	1	73ALDT
			DRB5*0105	0	
DQB1*0201 (DQ2)	8	30SIV, 45GE, 52LL, 56LPA, 57PA, 66DI, 70RK, 77DR	DQB1*0302 (DQ8)	3	56PPA, 57PA, 185I
DQB1*0401 (DQ4)	7	23L, 26G, 55PRL, 66DI, 70ED, 74SV, 185I	DQB1*0303 (DQ9)	1	185I
DQB1*0501 (DQ5)	10	14GL, 26G, 30HYV, 57PV, 70GA, 74SV, 77DR, 87AY, 116I, 125SQ	DQB1*0304	2	56PPA, 57PA
			DQB1*0305	4	26G, 56PPA, 57PA, 185I
			DQB1*0601	4	3P, 30YDV, 66DI, 67DIT
			DQB1*0603	1	30HYA
			DQB1*0604	4	30HYA, 57PV, 87GY, 130Q
			DQB1*0609	3	57PV, 87GY, 130Q
DQA1*0201	7	25FT, 47EKL, 48LF, 52FHR, 76IL, 79IRS, 175E	DQA1*0101	1	47ERW
DQA1*0301	7	25YS, 47EQL, 48LF, 52FRR, 75IVR, 79IRS, 175E	DQA1*0103	3	25FT, 40EK, 129HA
DQA1*0401	4	69T, 76IL, 79IRS, 175E	DQA1*0104	2	2G, 47ERW
DQA1*0601	5	25FT, 69T, 76IL, 79IRS, 175E	DQA1*0502	0	
			DQA1*0503	1	160SE

Recipient Type: DRB1*1101, 1501; DRB3*0202; DRB5*0101; DQA1*0102, 0501; DQB1*0301, 0602

in the clinical setting, although there is evidence that HLA-DP matching affects kidney transplant survival [12, 13] and that transplant patients can produce anti-HLA-DP antibodies [14, 63–65]. Such antibodies are specific for epitopes defined by short sequences and shared between groups of DP alleles [59, 64].

For the time being, the clinical use of HLAMatchmaker will primarily focus on determining structural compatibility for the HLA-DR and HLA-DQ loci. As illustrated in Table 7, certain antigens have many more mismatched eplets than others, and this might correlate to their ability of inducing specific antibodies. Applying the original HLAMatchmaker program, Dankers *et al.* have demonstrated that the frequency of HLA class I antibody production during pregnancy and after kidney transplantation correlates with the number of mismatched triplets on exposed HLA-A and HLA-B antigens [23]. No data are available on such correlations for mismatched class II eplet numbers.

This study has also generated information about the location, the surface expression, and the amino acid composition of each eplet. These factors undoubtedly play an important role in the immunogenicity of an eplet, *i.e.*, its ability to induce a specific antibody response. Recent studies have shown considerable differences in eplet (or triplet) immunogenicity [21, 28, 66]. Highly immunogenic eplets will increase sensitization and the risk for antibody-mediated rejection. An international collaborative study is underway to determine class I and class II eplet immunogenicity in kidney transplant patients [66].

Eplet versions of HLAMatchmaker for antibody analysis are slightly different from the matching programs because they incorporate the notion that antibody reactivity patterns cannot determine the differential recognition of individual eplets within a group of eplets unique for a given antigen (see Tables 5 and 6) or a combination of alleles. In such cases, the program uses a single eplet that represents a group of eplets. HLA typing differences between antibody producer and immunizer will define the mismatched eplet repertoire, and this information facilitates the interpretation of antibody reactivity patterns with HLA panels. The identification of reactive and nonreactive eplets permits a determination of HLA mismatch acceptability for sensitized patients.

At present, the eplet version of HLAMatchmaker should be considered as a hypothetical model for determining structural HLA compatibility and mismatch acceptability. Its validation depends on the outcome of clinical data.

The eplet versions of HLAMatchmaker and the HLA Patch Generator can be downloaded from the website <http://tpis.upmc.edu>. There is also an Eplets and Patches

file that shows how eplets have been assigned from polymorphic patches.

ACKNOWLEDGMENT

This study is supported by grant AI-55933 from the National Institutes of Health.

REFERENCES

1. Schoenemann C, Groth J, Leverenz S, May G: HLA class I and class II antibodies: monitoring before and after kidney transplantation. *Transplantation* 65:1519, 1998.
2. Mahoney RJ, Taranto S, Edwards E: B-Cell Crossmatching and kidney allograft outcome in 9031 United States transplant recipients. *Hum Immunol* 63:324, 2002.
3. Gebel H, Bray R, Nickerson P: Pre-transplant assessment of donor-reactive, HLA-specific antibodies in renal transplantation: contraindication vs. risk. *Amer J Transplant* 3:1488, 2003.
4. Iniotaki-Theodoraki AG, Boletis JN, Trigas G, Kalogeropoulou HG, Kostakis AG, Stavropoulos-Giokas CG: Humoral immune reactivity against human leukocyte antigen (HLA)-DQ graft molecules in the early posttransplantation period. *Transplantation* 75:1601, 2003.
5. Itescu S, Tung TC, Burke EM, Weinberg A, Moazami N, Artrip JH, Suciu-Foca N, Rose EA, Oz MC, Michler RE: Preformed IgG antibodies against major histocompatibility complex class II antigens are major risk factors for high-grade cellular rejection in recipients of heart transplantation. *Circulation* 98:786, 1998.
6. McKenna RM, Takemoto S, Terasaki PI: Anti-HLA antibodies after solid organ transplantation. *Transplantation* 69:319, 2000.
7. Terasaki PI: Humoral theory of transplantation. *Amer J Transplant* 3:665, 2003.
8. Duquesnoy RJ, Annen K, Marrari M, Kauffman HM Jr: Association of MB compatibility with successful intrafamilial kidney transplantation. *N Engl J Med* 302:821, 1980.
9. Matsuno N, Hidetoshi I, Ando A, T. N, Sato T, Ichikawa S, Sonoda T, Tsuji K: Importance of DQB as indicator in living related kidney transplant. *Transplantation* 49:208, 1990.
10. Tong JY, Hsia S, Parris GL, Nghiem DD, Cottingham EM, Rudert WA, Trucco M: Molecular compatibility and renal graft survival—the HLA DQB1 genotyping. *Transplantation* 55:390, 1993.
11. Middleton D, Mytilineos D, Savage D, Ferrara GB, Angelini G, Ameroso A, Trainor F, Gaweco A, Mazzola G, Delfino L, Berrino M, Oplez G: Matching for HLA-DPB1 alleles in zero mismatched HLA-A, -B and -DR Renal Transplants. *Transpl Proc* 24:2439, 1992.
12. Mytilineos J, Deufel A, Opelz G: Clinical relevance of HLA-DPB locus matching for cadaver kidney retrans-

- plants: a report of the Collaborative Transplant Study. *Transplantation* 63:1351, 1997.
13. Laux G, Mansmann U, Deufel A, Opelz G: A new epitope-based HLA-DP matching approach for cadaveric kidney transplantation. *Transplantation* 75:1527, 2003.
 14. Arnold M-L, Pei R, Spriewald B, Wassmuth R: Anti-HLA class II antibodies in kidney retransplant patients. *Tissue Antigens* 65:370, 2005.
 15. Duquesnoy RJ: HLA-Matchmaker: a molecularly based algorithm for histocompatibility determination. I. Description of the algorithm. *Hum Immunol* 63:339, 2002.
 16. Duquesnoy RJ, Takemoto S, De Lange P, Doxiadis IIN, Schreuder GMT, Claas FHJ: HLA-Matchmaker: a molecularly based algorithm for histocompatibility determination III. Effect of matching at the HLA-A,B amino acid triplet level on kidney transplant survival. *Transplantation* 75:884, 2003.
 17. Boehringer D, Reinhard T, Duquesnoy R, Boehringer S, Enczmann J, de Lange P, Claas F, Sundmacher R: Beneficial effect of matching at the HLA-A and B amino-acid triplet level on rejection free survival in penetrating keratoplasty. *Transplantation* 77:417, 2004.
 18. Laux G, Mytilineos J, Opelz G: Critical evaluation of the amino acid triplet-epitope matching concept in cadaver kidney transplantation. *Transplantation* 77:902, 2004.
 19. Duquesnoy R, Claas F: Is the application of HLA-Matchmaker relevant in kidney transplantation? *Transplantation* 79:250, 2005.
 20. Haririan A, Fagoaga O, Daneshvar H, Morawski K, Sillix D, El-Amm J, West M, Garnick J, Migdal S, Gruber S, Nehlsen-Cannarella S: Predictive value of HLA epitope matching using HLA-Matchmaker for graft outcomes in a predominantly African-American renal transplant cohort. *Clin Transplant* 20:226, 2006.
 21. Duquesnoy RJ, Marrari M: HLA-Matchmaker: a molecularly based algorithm for histocompatibility determination. II. Verification of the algorithm and determination of the relative immunogenicity of amino acid triplet-defined epitopes. *Hum Immunol* 63:353, 2002.
 22. Lobashevsky AL, Senkbeil RW, Shoaf JL, Stephenson AK, Skelton SB, Burke RM, Deierhoi MH, Thomas JM: The number of amino acid residues mismatches correlates with flow cytometry crossmatching results in high PRA renal patients. *Hum Immunol* 63:364, 2002.
 23. Dankers MKA, Witvliet MD, Roelen DL, De Lange P, Korfage N, Persijn GG, Duquesnoy RJ, Doxiadis IIN, Claas FHJ: The number of amino acid triplet differences between patient and donor is predictive for the antibody reactivity against mismatched HLA antigens. *Transplantation* 128:1236, 2004.
 24. Duquesnoy RJ, Witvliet MJ, Doxiadis IIN, de Fijter H, Claas FHJ: HLA-Matchmaker-based strategy to identify acceptable HLA class I mismatches for highly sensitized kidney transplant candidates. *Transpl Int* 7:31, 2004.
 25. Claas FHJ, Witvliet M, Duquesnoy RJ, Persijn G, Doxiadis IIN: The acceptable mismatch program as a fast tool to transplant highly sensitized patients awaiting a post-mortal kidney: short waiting time and excellent graft outcome. *Transplantation* 78:190, 2004.
 26. Iniotaki-Theodoraki A, Kalogeropoulou E, Apostolaki M, Doxiadis I, Stavropoulos-Giokas C: Humoral sensitization against rejected grafts: specific antibodies to graft immunogenic amino acid triplets. *Transplant Proc* 36:1728, 2004.
 27. Varnavidou-Nicolaidou A, Doxiadis I, Iniotaki-Theodoraki A, Patargias T, Stavropoulos-Giokas C, Kyriakides G: HLA class I donor-specific triplet antibodies detected after renal transplantation. *Transplant Proc* 36:1732, 2004.
 28. Adeyi O, Girnita A, Awadalla Y, Askar M, Shapiro R, Howe J, Martell J, Zeevi A, Nalesnik M, Rhandawa P, Demetris AJ, Duquesnoy RJ: Serum analysis after kidney transplant nephrectomy reveals restricted antibody specificity patterns against donor HLA class I antigens. *Transpl Immunol* 14:53, 2005.
 29. Doxiadis IIN, Duquesnoy RJ, Claas FHJ: Extending options for highly sensitized patients to receive a suitable kidney graft. *Curr Opin Immunol* 17:536, 2005.
 30. Goodman R, Taylor C, O'Rourke C, Lynch A, Bradley A, Key K: Utility of HLA-Matchmaker and single-antigen HLA-antibody detection beads for identification of acceptable mismatches in highly sensitised patients awaiting kidney transplantation. *Transplantation* 81:1331, 2006.
 31. Nambiar A, Duquesnoy RJ, Adams S, Oblitas J, Leitman S, Stroncek D, Marincola F: HLA-Matchmaker-driven analysis of response to HLA matched platelet transfusions in alloimmunized patients. *Blood* 107:1680, 2006.
 32. Duquesnoy R: A structurally based approach to determine HLA compatibility at the humoral immune level. *Human Immunol* 67:847-862, 2006.
 33. Chen J, Anderson JB, DeWeese-Scott C, Fedorova ND, Geer LY, He S, Hurwitz DI, Jackson JD, Jacobs AR, Lanczycki CJ, Liebert CA, Lin C, Madej T, Marchler-Bauer A, Marchler GH, Mazumder R, Nikolskaya AN, Rao BS, Panchenko AR, Shoemaker BA, Simonyan V, Song JS, Thiessen PA, Vasudevan S, Wang Y, Yamashita RA, Yin JJ, Bryant SH: MMDB: Entrez's 3D-structure database. *Nucleic Acids Res* 31:474, 2003.
 34. Hogue C: Cn3D: a new generation of three-dimensional molecular structure viewer. *Trends Biochem Sci* 22:314, 1997.
 35. Mullen MM, Haan KM, Longnecker R, Jardetzky TS: Structure of the Epstein-Barr virus gp42 protein bound to the MHC class II receptor HLA-DR1. *Mol Cell* 9:375, 2002.
 36. Ghosh P, Amaya M, Mellins E, Wiley DC: The structure of an intermediate in class II MHC maturation: CLIP bound to HLA-DR3. *Nature* 378:457, 1995.
 37. Bolin DR, Swain AL, Sarabu R, Berthel SJ, Gillespie P,

- Huby NJ, Makofske R, Orzechowski L, Perrotta A, Toth K, Cooper JP, Jiang N, Falcioni F, Campbell R, Cox D, Gaizband D, Belunis CJ, Vidovic D, Ito K, Crowther R, Kammlott U, Zhang X, Palermo R, Weber D, Guenet J, Nagy Z, Olson GL: Peptide and peptide mimetic inhibitors of antigen presentation by HLA-DR class II MHC molecules. Design, structure-activity relationships, and X-ray crystal structures. *J Med Chem* 43:2135, 2001.
38. Smith KJ, Pyrdol J, Gauthier L, Wiley DC, Wucherpfennig KW: Crystal structure of HLA-DR2 (DRA1*0101, DRB1*1501) complexed with a peptide from human myelin basic protein. *J Exp Med* 188:1511, 1998.
39. Lang H, Jacobsen H, Ikemizu S, Andersson C, Harlos K, Madsen L, Hjorth P, Sondergaard L, Svejgaard A, Wucherpfennig K, Stuart D, Bell J, Jones EY, Fugger L: A functional and structural basis for TCR cross-reactivity in multiple sclerosis. *Nat Immunol* 3:940, 2002.
40. Kim CY, Quarsten H, Bergseng E, Khosla C, Sollid LM: Structural basis for HLA-DQ2-mediated presentation of gluten epitopes in celiac disease. *Proc Natl Acad Sci U S A* 101:4175, 2004.
41. Lee KH, Wucherpfennig KW, Wiley DC: Structure of a human insulin peptide-HLA-DQ8 complex and susceptibility to type 1 diabetes. *Nat Immunol* 2:501, 2001.
42. Siebold C, Hansen BE, Wyer JR, Harlos K, Esnouf RE, Svejgaard A, Bell JI, Strominger JL, Jones EY, Fugger L: Crystal structure of HLA-DQ0602 that protects against type 1 diabetes and confers strong susceptibility to narcolepsy. *Proc Natl Acad Sci U S A* 101:1999, 2004.
43. Combet C, Jambon J, Deleage G, Geourjon C: Geno3D: automatic comparative molecular modeling of protein. *Bioinformatics* 18:213, 2002.
44. Viken HD, Paulsen G, Drover S, Marshall WH, Sollid LM, Gaudernack G, Thorsby E: Influence on antibody recognition of amino acid substitutions in the cleft of HLA-DQ2 molecules. Suggestive evidence of peptide-dependent epitopes. *Hum Immunol* 44:63, 1995.
45. Smith KD, Mace BE, Valenzuela A, Vigna JL, McCutcheon JA, Barbosa JA, Huczko E, Engelhard VH, Lutz CT: Probing HLA-B7 conformational shifts induced by peptide-binding groove mutations and bound peptide with anti-HLA monoclonal antibodies. *J Immunol* 157:2470, 1996.
46. Takamiya Y, Sakaguchi T, Miwa K, Takiguchi M: Role of HLA-B*5101 binding nonamer peptides in formation of the HLA-Bw4 public epitope. *Int Immunol* 8:1027, 1996.
47. Mulder M, Eijssink C, Kester MGD, Franke MEI, Kardol MJ, Heemskerk MHM, van Kooten C, Verreck FA, Drijfhout JW, Koning F, Doxiadis IN, Claas FJH: Impact of peptides on the recognition of HLA class II molecules by human HLA antibodies. *J Immunol* 175:5950, 2005.
48. Altschul S, Madden T, Schäffer A, Zhang J, Zhang Z, Miller W, Lipman D: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389, 1997.
49. Robinson J, Waller M, Parham P, de Groot N, Bontrop R, Kennedy LJ, Stoer P, Marsh SG: IMGT/HLA and IMGT/MHC: sequence databases for the study of the major histocompatibility complex. *Nucleic Acids Res* 31: 311, 2003.
50. Albert ED, Baur MP, Mayr WR: Histocompatibility Testing 1984. In *Class II Antigen Reports*. Berlin, Springer-Verlag, 1984.
51. Dupont B (Ed): Immunobiology of HLA. In *Histocompatibility Testing 1987*. Vol. I. New York, Springer-Verlag, 1987.
52. Tjuji K, Aizawa M, Sasazuki T (Eds): HLA1991. Proceedings of the Eleventh International Workshop and Conference. Vol. 1. Oxford, Oxford University Press, 1991.
53. Charron D: Allele and Haplotype Society Reports #10-18. In *Proceedings of the Twelfth International Histocompatibility Workshop and Conference*. Sevres, EDK Medical and Scientific International Publisher, 1997.
54. Navarrete C, Brown C, De Lange P, Schreuder GMT: 12th International Histocompatibility Workshop HLA Class II Monoclonal Antibodies Study. In Charron D (ed): *HLA Genetic Diversity of HLA. Functional and Medical Implication*. Vol. 1, Paris, EDK Publisher, 1997.
55. Schreuder GMT: Serological Definition of HLA-DR and -DQ Polymorphisms. The Hague, CIP-Data Koninklijke Bibliotheek, 1989.
56. Duquesnoy RJ, Mulder A, Askar M, Fernandez-Vina M, Claas FJH: HLAMatchmaker-based analysis of human monoclonal antibody reactivity demonstrates the importance of an additional contact site for specific recognition of triplet-defined epitopes. *Hum Immunol* 66:749, 2005.
57. Fuggle SV, Carter C, Watts F, Kirkley J, Morris PJ: Monoclonal antibody definition of multiple polymorphic epitopes on HLA-DRw52. *Human Immunol* 19:255, 1987.
58. Taylor CJ, Ugozoli L, Tanigaki N, Tosi G, Bunce M, Ting A, Ferrara GB: Antigen Society #29 Report (DRw52). In Dupont B (ed): *Immunobiology of HLA*. Vol. I. Histocompatibility Testing, New York, Springer Verlag, 1987:273, 1987.
59. Tongio MM, van der Berg-Loonen E, Bignon JD, Chandanayingyong D, Dormoy A, Eiermann T, Marshall W, Park MS, Scheuder GMT: HLA-DP Detected by Serology. Sevres, EDK Medical and Scientific International Publisher, 1997.
60. Fernandez-Vina M, Moraes JR, Moraes ME, Miller S, Stastny P: HLA class II haplotypes in Amerindians and in black North and South Americans. *Tissue Antigens* 38: 235, 1991.
61. Clayton J, Lonjou C, Whittle D: DRB1-DQA1-DQB1 Haplotype Frequencies. In Charron D (ed): *HLA Genetic Diversity of HLA. Functional and Medical Implication*. Paris, EDK Publisher, 1997.

62. Klitz W, Maiers M, Spellman S, Baxter-Lowe LA, Schmeckpeper B, Williams TM, Fernandez-Vina M: New HLA haplotype frequency reference standards: high-resolution and large sample typing of HLA DR-DQ haplotypes in a sample of European Americans. *Tissue Antigens* 62:296, 2003.
63. Pfeffer K, Vogeler U, Albrecht KH, Eigler FW, Buchholz B, Grosse-Wilde H: HLA-DP antibodies in patients awaiting renal transplantation. *Transpl Internat* 8:180, 1995.
64. Youngs D: DP alloantibodies. *American Society for Histocompatibility and Immunogenetics Quarterly* 2:60, 2004.
65. Qiu J, Cai J, Terasaki PI, El-Awar N, Lee JH: Detection of antibodies to HLA-DP in renal transplant recipients using single antigen beads. *Transplantation* 80:1511, 2005.
66. Duquesnoy RJ, Claas FHJ: Progress report of 14th International Histocompatibility Workshop Project on the structural basis of HLA compatibility. *Tissue Antigens In Press*, 2006.