


Oilseeds and Roots





- Soil Fertility
- Stand Establishment
- Winter Survival
- Rhizoctonia s. Root Imagery



Pie in the sky VS. Down to earth



Soil Fertility

- N-Rate Calculator for Spring Canola
- The 4Rs for Nutrient Management
- Micronutrients in Canola




N Rate Calculator

Data used to create an N calculator

- Multiple small plot experiments
- Multiple locations
- Multiple N rates including 0 fertilizer application




Unit Nitrogen Requirements





Unit Nitrogen Requirement as Effected by Depth of Sampling

Optimum Yield vs. UNR for different sampling







N Rate Calculator Exercise





Screen shots Share and Make a Copy








| Box B Credit From Previous Grain and Forage Legume Crops | |
|--|--------------------|
| Preceding crop and yield | Credit (lb N/acre) |
| Peas > 2500 lb/acre | 20 |
| Peas 1500 to 2500 lb/acre | 15 |
| Peas < 1500 lb/acre | 10 |
| Lentils >1000 lb/acre | 10 |
| Alfalfa | 50 |

Koenig Dryland Winter Wheat EB1987E 2005



Yield Response to N in Winter Canola





Calculator Sources of Error


- % Organic Matter Mineralization
- Yields greater than 2500 lbs/A can push UNR very low
- Legume credits and wheat debits are based on older data



What additional features would you like to see?



Effects of additional N on seed quality



Porter et al 2019



Considering the correct placement, source, and timing of N for canola




Ammonia toxicity from the fertilizer band





Increasing Urea






Urea

AS


UAN






Tap Root Survival

Urea = 4.7 AS = 9.7 UAN = 20.6





Relationship between in-furrow concentration and row spacing



In Furrow Concentration Calculator






Micronutrients in Canola Tissue Levels and Application Methods




Tissue Tests For Different Varieties in the Same Field






Tissue Tests For Different Varieties in the Same Field





Zn and B Application Trials








Does Canola Variety Effect Responses to Micronutrient Applications?







Future of Micronutrient Research

- Establish 'norms' for high yielding canola fields.
- Examine intra- and inter- field variation
- Examine intra- and inter- region variation



And that's all on fertility for now



Stand Establishment

- Standard Practice
 - -Planting dates, depths, moisture
- Pie in the sky
 - -In-furrow moisture
 - -Seed Priming
 - -Companion Cropping




Planting Dates, Depths, and Moisture

Water and Temperature



Planting Dates and Tillage in The Great Plains





Seed Priming and In-Furrow Moisture

- On farm strip trial and small plot tests to date have not shown and improvement
- Lab experiments being conducted to develop conditions which these practices are 'likely' to work in.




In-Furrow Water Application






Relationship between in-furrow concentration and row spacing






Nurse Crops For Winter Canola Establishment








Winter Survival








Winter Survival

- Weather severity is the most crucial component
- Tillage and crown height appear to have some effect




Dual Purpose Crops







Timeline for Dual Purpose Companion Planted Canola



Applicable to WW-F and Preventative Plant Scenarios







2017-2018 Grazing Trial






| Treatments | Yield (lbs/a) | Grazing pressure (46 head) |
|------------|---------------|----------------------------------|
| Pasture 1 | 2464 | Heavy |
| Pasture 2 | 2143 | Severe |
| Pasture 3 | 3322 | Light |
| Ungrazed | 3384 | None |

Table 1: Seed yield and cattle gain from canola grazing.



2019-2020 Crop Year

Oat-Canola

















Improved Forage From Intercropping Oats

| Table 2 | Canola | Oat & Canola |
|---------------------------------|--------|--------------|
| Dry Matter | 11.6 | 19 |
| Crude Protein | 21.3 | 14.2 |
| Acid Detergent Fiber (ADF) | 16.9 | 20.2 |
| Neutral Detergent Fiber (NDF) | 17.3 | 29.5 |
| Relative Feed Value (Estimated) | 407 | 231 |




Monitoring Fall Water Usage





Reduced Water Usage in Grazed vs Ungrazed Plots





Additional Insect Challenges





Rhizoch



Lesion Formation









WOCS annual winter workshops

Jan. 28th Almira, WA

Jan. 30th Clarkston, WA



WASHINGTON OILSEED CROPPING SYSTEMS

Part of the Washington State Biofuels Initiative



Thank you!






Intercrops as and opportunity to stack diversity in the absence of cover crops



Complementary Architectures










Theoretical Advantages of peaola
















Complimentary Architectures

























Logistical Concerns

