

## HS4440-WU01 COVID-19: Working Safely at LLNL During the COVID-19 Pandemic REVISION 3

Version 1.2.0 / April 23, 2020

## **Authors/Instructional Designers:**


C. I. Hall M. F. Hagerty

### **Auspices Statement:**

Lawrence Livermore National Laboratory is operated by Lawrence Livermore National Security, LLC, for the U.S. Department of Energy, National Nuclear Security Administration under Contract DE-AC52-07NA27344.

## Working Safely at LLNL During the COVID-19 Pandemic

### 1 Welcome




### 2 Announcement



## 3 COVID-19: Working Safely at LLNL During Minimum Safe Operations



### 4 Overview



### **5 Reduced Minimum Safe Operations:**



## **Reduced Mission-Critical Operations:**

"In the face of the coronavirus pandemic, the Lab has reduced its staffing at sites 200 and 300 to that required for safety, security and the minimal maintenance of facilities.

Small groups of employees have made themselves available at the Lab to respond without delay to questions about the stockpile that arise in the course of surveillance and dismantling, to maintain critical national response activities like NEST and NARAC and to engage in projects that have spun up to respond directly to the pandemic. I want to extend my thanks to all those who have been on-site keeping the Lab, and the nation safe and secure during these difficult days.

Our planning for what we have been calling minimum safe operations always included the continuation and resumption over time of a set of mission critical activities on-site, a posture NNSA deems "Reduced Mission-Critical Operations." Doing so is consistent with local shelter-in-place orders, which exempt critical defense activities.

The Laboratory's top priority is protecting the health and safety of the workforce during any on-site operations, including those resuming this week. The number of people on-site will remain small to minimize contact and enable "social distancing." Protocols for working on-site include the precautions we have all been practicing, as well as access to PPE and the maintenance of clean work areas. These protocols have been formalized into a Site-wide Work Control Document that spell out the controls designed to protect employees from the potential spread of the coronavirus in the workplace."

Director Bill Goldstein, 3/30/2020


### 6 Work Planning & Control




## Work Planning & Control \*\*UPDATED - 4/23/2020\*

- All workers on-site are required to wear a face covering when interacting with, or near, other individuals on-site. Face coverings will be provided by the Laboratory.
- If you are on-site performing AL-0 work, such as inspections, walk-throughs, and office
  work, this presentation serves as your guidance for implementing appropriate controls
  for your work.
- If you are on-site performing work under a WCD or Competent Worker program, your RI will conduct a pre-start review (WCDs) or a pre-job brief (Competent Workers), prior to commencing work. The Laboratory has approved a site-wide Work Permit with controls for the COVID-19 hazard. During the pre-start review/pre-job brief, your RI will discuss which controls for COVID-19 will be applied, based upon the work controls in your JHA, and whether social distancing can be maintained.
  - If the controls in your JHA are more protective than the COVID-19 controls, the more
    protective controls will be used.

### 7 What is COVID-19?




## 8 COVID-19 is responsible for a world-wide pandemic



### 9 What are the Symptoms of COVID-19?

## What are the Symptoms of COVID-19?

- · Symptoms will vary from person to person.
- Reported illnesses have ranged from no symptoms to mild symptoms to severe illness.
- The clinical picture (spectrum of symptoms) is evolving as more information becomes available from across the world.
  - Other symptoms that have been reported include fatigue, sore throat, headache, loss of sense of taste, chills and diarrhea.
- Symptoms usually develop 5 days after exposure but this can range between 2-14 days.
- Percentage of cases by illness severity (based on current data):
  - ~81% of confirmed cases are mild (no pneumonia)
  - ~14% of confirmed cases are severe (clinical symptoms include shortness of breath)
  - ~5% of confirmed cases are critical (pneumonia, respiratory failure)
- Those individuals age ≥65 years, or with a history of other illnesses, like diabetes, lung disease, or immunocompromising conditions, have a higher risk of severe illness.



### 10 What do we know about how COVID-19 is spread?


### \*\*UPDATED - 4/13/2020\*\*

## What do we know about how COVID-19 is spread?

Current evidence suggests that there may be <u>three</u> potential routes of transmission of COVID-19. The relative role each route plays in community spread of the disease remains somewhat unclear.

| relative role each route plays in community spread of the disease remains somewhat unclear. |  |  |
|---|--|--|
| Route of<br>Transmission  | Definition   | Relative Role in COVID-19<br>transmission  |
| Droplet<br>transmission   | Respiratory secretions containing the virus are expelled when an infected individual coughs, sneezes, breathes and/or speaks and land in the mouth, nose or eyes of someone nearby (generally within 6 feet).                                      | Believed to be a <u>major driver</u> of<br>community spread.   |
| Contact<br>transmission   | Respiratory droplets containing the virus settle on surfaces. Someone touches that surface and then touches their mouth, nose or eyes which can potentially result in transmission of the virus to themselves.                                     | <ul> <li>Believed to <u>contribute</u> to<br/>community spread, but not believed<br/>to be the primary driver.</li> </ul>  |
| Aerosol<br>transmission   | Small particles (generally defined as smaller than 5µm; sometimes referred to as droplet nuclei) containing virus are inhaled into the lungs. Aerosol transmission can occur over short ranges (within 6 ft) as well as potentially longer ranges. | <ul> <li>Believed to play an important role in spread within healthcare system resulting from aerosol-generating medical procedures.</li> <li><u>UNCLEAR role</u> in community spread, not believed to be a major driver of community spread.</li> </ul> |


### 11 Understanding Transmission




### 12 Direct Transmission



# 13 Contact Transmission: how long can the SARS-CoV-2 virus survive on surfaces?



### 14 What about aerosol transmission



### 15 How long does the virus last?

Short-range aerosol transmission appears to play an important role in the spread of COVID-19 among healthcare workers performing aerosol-generating medical procedures (e.g., intubations, removing patients from ventilator).


Recent reports indicate that SARS-CoV-2, the virus that causes COVID-19 can survive as an aerosol in the air for up to 3 hours.

- This study used an aerosol generating device (nebulizer) to generate the aerosol in this study which <u>does not</u> mimic respiratory secretions generated when an individual coughs or sneezes.
- The nebulizer does mimic aerosol-generating procedures conducted in a <u>healthcare setting</u> (e.g., intubation, disconnecting a patient from a ventilator).




The half-life of the virus is in air is approximately 1 hour- meaning that after about an hour, about 50% of the virus particles are no longer infectious.


## 16 Aerosol transmission in the community



# 17 We know that People Can Transmit COVID-19 even without showing any symptoms:



## 18 The Unfortunate Answer is: We Just Don't Know Yet



### 19 The Unfortunate Answer is: We Just Don't Know Yet

## Here's What We Don't Know!

\*\*NEW-4/13/2020\*\*

Key Outstanding Questions that Make it Difficult to Determine the Relative Role of Aerosol Transmission in Community Spread of COVID-19:

- · Is the virus-laden aerosol capable of traveling long distances?
  - If it is able to travel long distances is the virus still present in sufficient number to be able to cause an infection? Or does the virus get diluted to a relatively harmless concentration?
- The most recent <u>study</u>, published on April 10, 2020, in which researchers performed air sampling in hospital wards housing COVID-19 patients – indicated that SARS-CoV-2 RNA was detected up to 4 meters (~13 ft) from patients.
  - Importantly though- the researchers did not assess virus viability by attempting to culture the virus from the air samples. Therefore, they cannot conclude whether the air samples contained viable virus.
  - In addition, these were patients that were symptomatic; therefore, conclusions cannot be drawn for asymptomatic spreaders.
- Virus dilution in the air and inactivation due to desiccation of the virus as it travels long distances over longer periods of time may reduce the potential for long-distance aerosol transmission, but this remains unclear.
- The CDC believes that airborne transmission from person-to-person over long distances is unlikely.



### 20 Key Take Away Messages

#### \*\*NEW-4/13/2020\*

### Key Take Away Messages:



- People likely emit virus-containing particles in a range of sizes doing a number of everyday activities such as breathing, speaking, singing, coughing, sneezing.
- Some evidence suggests that small viruscontaining particles (infectious aerosols) can be important in short-range (<6 ft) aerosol transmission of COVID-19.
- The role of long range (>6 ft) aerosol transmission of COVID-19 remains unclear. However, it is not currently believed to be a major driver of the COVID-19 pandemic.

### 21 So What DO We Know?

Household Contacts Provide Evidence that Close Contact Plays an Important Role in COVID-19 Transmission

\*\*NEW-4/13/2020\*\*

We know that persons sharing a residence with an individual that has COVID-19 have a higher probability (risk) of getting infected than people who do not have a member of their household that is infected with COVID-19.

#### What does this tell us?


This tells us that being in  $\underline{\text{close contact}}$  with someone who has COVID-19 is one of the most important risk factors.




### 22 OK - But What Do We Mean by "Close Contact"?



### 23 How long is too long?



# 24 Unfortunately- there is no magic formula for determining how long or how far apart constitutes safe!




### 25 With So Much Uncertainty – What Are We Supposed to Do??

### \*\*NEW-4/13/2020\*\* With So Much Uncertainty – What Are We Supposed to Do?? The idea that so many everyday activities can potentially transmit COVID-19 can be scary! Not knowing the actual risks associated with each route of transmission can be even scarier! And not having clear, executable guidance on how close is too close or how long is too long can be frustrating! So What Can We Do? 1. Realize that this is a new virus that we don't fully understand and acknowledge that we will have to be flexible enough to adjust to a rapidly evolving global health crisis with new information coming at us faster than we can process. 2. Consider all potential routes of exposure, assess the relative risk based on the best available data, and identify appropriate controls to minimize those risks. The Lab has a team of subject matter experts dedicated to searching for and reviewing new information and data on a daily basis and assessing the Lab's current controls to ensure we are protecting our workers based on the best available current data.

### 26 How Do I Work Safely During Reduced Mission Critical Operations?



# 27 LLNL has Implemented Controls to Reduce Risk to Personnel Working Onsite



### 28 No Single Control Alone is Protective:



# 29 The MOST IMPORTANT Things You Can Do, Both at Work and At Home, To Minimize Your Risk of Contracting or Spreading COVID-19:



and inconsequential, the following four behaviors, discussed in more detail on the following slides, <u>are the most important things you can do to protect yourself and others.</u>





Stay home if you are sick; Go home if you become sick at work.



Wash your hands frequently with soap and water!



### 30 If You Are Sick - STAY HOME!

## IPDATED-4/13/2020\*\* If You Are Sick — STAY HOME!



If you have been told by your Personal Healthcare Provider (HCP) that you have or may have COVID-19 – either by testing or symptoms alone – follow the guidelines provided by your provider and contact Health Services.

Contact Health Services: By phone: 2-6843 (COVID)

By email: <a href="mailto:covid19@llnl.gov">covid19@llnl.gov</a>
Information is passed to Health Services via an established confidential process

If you are sick but have **NOT** been told by your HCP that you have or may have COVID-19- stay home until you are symptom-free for <u>72-hours</u> without the use of symptom altering medications (e.g., fever-reducers, cough-suppressants).

 If you have used 5 or more consecutive days of sick leave – contact Health Services PRIOR to returning to work.


If you get sick while you are at work – leave work IMMEDIATELY – do not linger or attempt to complete your shift.

- · Ensure you leave your work area in a safe state.
- Notify your supervisor (or a co-worker) and send an email to covid19@IInl.gov.

### 31 If You Are Sick – STAY HOME!



## 32 COVID-19 Case Management Aims to Prevent Transmission of COVID-19 at the Workplace



### 33 Social Distancing: What is it?

## Social Distancing: What Is It?

**Purpose**: Social distancing reduces the rate of disease transmission and can stop an outbreak.

What is Social Distancing? Social distancing is a <u>group of measures</u> taken by Public Health Officials to limit the spread of a disease.

Social Distancing is one of the most powerful tools to combat all possible routes of exposure!

#### How do we implement social distancing at work?


Maintain physical separation from others to the maximum extent possible!

#### MORE DISTANCE IS BETTER:

- · Stay home if you are sick.
- · Telecommute to the maximum extent possible.

#### WHEN YOU HAVE TO COME TO WORK:

- · Limit the number of people in one place to the extent possible.
  - Don't congregate in groups in for example your office, lunch/break rooms, conference rooms, hallways.
  - At a minimum maintain 6 feet separation from others to the extent possible.
  - · Minimize the amount of time spent with others.



Social distancing deliberately increases the physical space between people to prevent sick people from coming in contact with healthy people to reduce the opportunity for spreading COVID-19.

### 34 Wash your Hands Frequently

## Wash Your Hands Frequently

- Wash your hands frequently with soap and water for at least 20 seconds.
- If soap and water are not immediately available, use an alcohol-based hand-sanitizer containing 260% alcohol.
  - · If hands are visibly dirty, wash with soap and water!

#### Wash your Hands or Use Hand Sanitizer:


- · Before touching your face for any reason
- · Before preparing food/eating
- After coughing or sneezing
- · After touching high-touch surfaces
- Before and after putting on, adjusting and taking off your N95 respirator.

Note: Handwashing is more effective at <u>removing</u> germs than hand sanitizer.




Watch the video
proper hand-washing technique


### 35 Handwashing



### 36 Avoid Touching Your Face – Especially Your Mouth, Nose and Eyes



## 37 Cover your Cough and Sneeze:



# 38 Frequent Cleaning and Disinfection of High-Touch Surfaces Augments Infection Control Measures




### 39 Important Considerations Regarding Cleaning and Disinfection:



### 40 Frequently Clean High-Touch Surfaces:



### 41 Use PPE to Augment – But Not Replace – Infection Control Measures



### 42 To Mask or Not to Mask- That is the Question!



### 43 What is the Purpose of Wearing a Face Covering?

#### \*NEW - 4/23/2020\*\*

What is the Purpose of Wearing a Face Covering?

### COVID-19 is believed to be primarily spread through respiratory droplets.

- · Droplets are expelled by you when you talk, breath, cough, sneeze, laugh and sing.
- Individuals can be infected with the virus and expelling potentially infectious droplets without showing any symptoms.
- People can also be infected and contagious up to 48 hours before developing symptoms.
- · As such, infected people can unintentionally spread the virus without recognizing that they are infected.

Wearing a face covering reduces the chance that the mask wearer spreads potentially infectious respiratory droplets to others or to surfaces.

When combined with social distancing (maintaining at least 6 feet separation between yourself and others) — wearing a face covering when around others can reduce the risk of transmitting the virus to others.

Wearing a face covering is NOT a substitute for social distancing and frequent handwashing!

### 44 Am I Required to Wear a Face Covering While I am On-site?

#### \*\*NEW = 4/23/2020\*\*

Am I Required to Wear a Face Covering While I am On-site?

In line with the newly issued order from the Alameda County Public Health Department, the Laboratory now requires that all on-site personnel use face coverings.

Face coverings <u>are required</u> while on-site whenever you are interacting with or are near others. Examples may include:

- · In hallways, stairwells and elevators.
- · In common areas such as restrooms and kitchens.
- · In shared vehicles when more than one person is present.

### Face coverings are not required when:

- · You are alone in your individual office,
- When engaging in <u>outdoor</u> activities such as walking, biking, running or eating lunch
   when you can maintain the recommended 6 feet of separation from others.

Note: employees are encouraged to have a face covering with them and available to put on in case they encounter a situation where they cannot maintain social distancing.




### 45 Can I wear a Face Covering with an Exhalation Valve?

### Can I wear a Face Covering with an Exhalation Valve? Face coverings should not have a one-way exhalation valve. The Alameda County Public Health Order does not recognize WHY? masks that incorporate a one-way valve (exhalation valve) as face coverings. The intent of the face covering is to protect **Exhalation valve** others from respiratory droplets released by Makes it easier to the wearer when they cough, sneeze or talk. breathe Helps keep your face One-way (exhalation) valves allow some cooler portion of respiratory droplets expelled by the wearer to be released through the Reduces moisture build up inside the facepiece exhalation valve.


### 46 Are Face Coverings Being Provided by the Lab?




### 47 Can I Bring my Own Face-Covering from Home for Use On-Site?




### 48 How to Properly Don (put on) and Doff (take off) Face Coverings




### 49 Certain Job Duties Will Require the Use of an N95 Respirator




### 50 Why is the Lab Issuing Valved N95 Respirators for Some Job Duties?



## 51 Key Factors for Effective Use of an N95 Respirator:



### 52 Ensuring a Proper Fit/Seal: Facial Hair Matters



### 53 Properly donning and doffing an N95

\*\*UPDATED - 4/23/2020\*\*

## Know How to Properly Put On and Take Off Your N95 Respirator:

- How you put on and take off your respirator will affect how well it protects you from potentially infectious droplets.
- · You must ensure a proper fit.
- Watch this video to learn how to properly put on your respirator and perform a seal check:

#### Watch the Video

Always wash your hands before and after putting on, adjusting and taking off your N95 respirator!



#### IMPORTANT VIDEO NOTES:

- The video shows a different style N95 than that provided at the laboratory. However, the overall procedure applies to donning/doffing all N95 respirators.
- The N95s being provided by the Lab do not come in different sizes, but users still need to be sure to perform a fit check!
- The video says to discard your respirator after use this is because
  it is geared toward healthcare workers. <u>DO NOT DISCARD YOUR</u>
  <u>RESPIRATOR AFTER A SINGLE USE!</u> When cared for properly,
  respirators used in most work settings can be re-used safely for an
  extended period of time. See next slides!

### 54 Store Your N95 Respirator Properly:


### Store Your N95 Respirator Properly:

- Store your respirator in a clean, resealable plastic bag OR a paper bag.
  - Discard the bag after each use (paper bag) or clean the interior of the bag regularly to keep the inside of the N95 respirator clean (plastic bag).
  - Allow your N95 to dry in a clean area prior to placing it in a bag.
  - Plastic bags: Leave a corner of the bag open during storage to prevent moisture build up inside the bag.
- Store your respirator in a way that minimizes the potential for the respirator to become damaged or deformed.
- · Label the bag with your name to prevent use by others.
- · Consider labeling your respirator with your name.
  - · Write your name on the straps or side of the respirator.
  - · Do not write across the filter of the respirator.







### 55 How Long Can I Re-Use my Respirator?



### 56 Where Can I Get an N95 Respirator?



### 57 Consider Wearing Gloves If You Can't Control the Surfaces You Touch:



### 58 Know How to Properly Use Gloves:



# 59 How Do We Navigate Protecting Ourselves When we Encounter Situations that may be Unique to Being at Work?




### 60 Can I Hold In-Person Meetings?



# 61 Things and Places that we Normally Take for Granted can Pose Unique Risks During the COVID-19 Pandemic:



### 62 How Should I Navigate Common Areas?



### 63 Social Distancing at Lunch

## Social Distancing At Lunch!

- Cafeterias remain closed during reduced mission critical work.
- Eat in your office, if you have one, whenever possible.
- If you do not have an office practice social distancing at lunch to the extent possible:
  - · Consider eating outside for some fresh air!
  - Limit the number of individuals in the break room to maintain social distancing.
  - Stagger your lunches with your co-workers whenever possible.



## 64 Performing Everyday Risk Assessments Using What we Have Learned about COVID-19:

## Performing Everyday Risk Assessments Using What We Have Learned about COVID-19:

A mental checklist, based on information we know about COVID-19 and the mechanisms available to protect ourselves from spreading and acquiring COVID-19, can help us navigate unique situations we may encounter at work.

Assume everyone is potentially infected, and every surface is potentially contaminated.

#### With this in mind- think through how you would approach the situation:

- ☐ Will there be close contact (consider time and distance)?
- ☐ Are there physical barriers in place that can provide some level of protection?
- ☐ Is there sufficient airflow or is airflow limited?
- ☐ Are there surfaces that I need to be concerned about?


### Based on the answers to these questions:

What controls should I apply to reduce the risk of exposure?

### 65 Elevators May Pose Unique Risks – Small Spaces with Limited Airflows

## Elevators May Pose Unique Risks – Small Spaces with Limited Airflow ■ Will there be close contact (consider time and distance)? YES (<6 ft distance)</p> Are there physical barriers in place that can provide some level of protection? NO ☐ Is there sufficient airflow or is airflow limited? LIMITED AIRFLOW ☐ Are there surfaces that I need to be concerned about? YES- elevator buttons, handrails Based on the answers to these questions: ■ What controls should I apply to reduce the risk of exposure? When possible, limit the number of riders in an elevator to ONE When not possible, limit the number of riders to the extent possible and all individuals should wear a face covering (e.g., KN95 mask, surgical-style mask or cloth mask). · Be cognizant of high-touch surfaces! · Use a barrier (e.g., paper towel, tissue) to press buttons Wash hands (or use hand sanitizer) after interacting with any high-touch surface! · Avoid touching your face (eyes, nose, mouth) until you have had an opportunity to wash your hands!


# 66 CAIN booths May Pose Unique Risks – Small Spaces with Limited Airflows




### 67 Public Restrooms – How to Approach During a Global Pandemic

## Public Restrooms – How to Approach During a Global Pandemic ■ Will there be close contact (consider time and distance)? Maybe (could be less than 6 feet and generally not longer than 10 minutes potential exposure) ☐ Are there physical barriers in place that can provide some level of protection? YES (stall walls and door provide a physical barrier between you and others). ☐ Is there sufficient airflow or is airflow limited? SUFFICIENT ☐ Are there surfaces that I need to be concerned about? YES-door handles, stall handles, faucets, paper towel dispensers Based on the answers to these questions: What controls should I apply to reduce the risk of exposure? · Consider not using adjacent stalls. · Consider not using adjacent sinks - wait for others to finish washing their hands. · Be cognizant of high-touch surfaces! · Wash hands (or use hand sanitizer) after interacting with any high-touch surface! Avoid touching your face (eyes, nose, mouth) until you have had an opportunity to wash your hands! Wipe down high-touch surfaces prior to and after use. · Wear a face covering when you are near others in a public setting.

### 68 Sharing a Car With a Coworker During the COVID-19 Pandemic



### 69 COVID-19 Hotline:



### 70 Resources:

